

PRODUCT
PRODUIT MANGANESE

PROVINCE OR
TERRITORY PROVINCE OU
TERRITOIRE

British Columbia

N.T.S. AREA 92 C/16
RÉGION DU S.N.R.C.

REF. Mn 2
RÉF.

NAME OF PROPERTY
NOM DE LA PROPRIÉTÉ

SHERK LAKE

OBJECT LOCATED - Mn symbol on Geological map.
OBJET LOCALISÉ

UNCERTAINTY 500 m
FACTEUR D'INCERTITUDE

Lat. 48°55'20" Long. 124°12'30"
Lat. Long.

Mining Division
Division minière Victoria

District
District

County
Comté Township or Parish
Canton ou paroisse

Lot
Lot Concession or Range
Concession ou rang

Sec
Sect. Tp. Ct. R. R.

OWNER OR OPERATOR/PROPRIÉTAIRE OU EXPLOITANT

DESCRIPTION OF DEPOSIT/DESCRIPTION DU GISEMENT

The host rocks are cherty tuffs of the Palaeozoic Sicker Group. Rhodochrosite and rhodonite occur as irregular masses in a bed of red jasper 2 to 3 feet thick. The jasper has been traced for more than 1 000 feet along the strike, but only locally does it contain manganese.

HISTORY OF EXPLORATION AND DEVELOPMENT
HISTORIQUE DE L'EXPLORATION ET DE LA MISE EN VALEUR

The occurrence is located at about 950 m elevation on the south side of Sherk Lake, at the head of Chemainus River, some 80 km northwest of Victoria.

121596

HISTORY OF PRODUCTION/HISTORIQUE DE LA PRODUCTION

REFERENCES/BIBLIOGRAPHIE

Fyles, J.T.; Geology of the Cowichan Lake Area; Bulletin
No. 37, p. 69, BC Dept. of Mines, 1955.

MAP REFERENCES/RÉFÉRENCES CARTOGRAPHIQUES

#Geological Map of the Cowichan Lake Area, Sc. 1": 1 mile,
accomp. Bulletin No. 37, BC Dept. of Mines 1955.

*Map 92 C/16, Cowichan Lake (Topo.), Sc. 1: 50 000.

REMARKS/REMARQUES

| | | | | | | | |
|---------------------------------|-------|--|--|--|--|--|--|
| Comp./Rev. By Comp./rév. par | DMacR | | | | | | |
| Date Date | 01-88 | | | | | | |