

PRODUCT ZINC
PRODUIT

PROVINCE OR PROVINCE OU
TERRITORY TERRITOIRE

British Columbia

N.T.S. AREA 94 D/7
RÉGION DU S.N.R.C.

REF. ZN 2
RÉF.

NAME OF PROPERTY
NOM DE LA PROPRIÉTÉ

ASITKA RIVER

OBJECT LOCATED
OBJET LOCALISÉ

UNCERTAINTY
FACTEUR D'INCERTITUDE

Lat. 56°21'

Long. 126°30'30"

Lat.

Long.

Mining Division **Omineca**

District

Cassiar

County
Comté

Township or Parish
Canton ou paroisse

Lot
Lot

Concession or Range
Concession ou rang

Sec
Sect.

Tp.
Ct.

R.
R.

OWNER OR OPERATOR/PROPRIÉTAIRE OU EXPLOITANT

DESCRIPTION OF DEPOSIT/DESCRIPTION DU GISEMENT

The formation consists of andesitic and basaltic tuffs of Upper Triassic or Jurassic age, belonging to the Takla group. A number of northwesterly and northerly trending faults occur in the vicinity. A 6 inch wide vein, exposed for a length of 2 feet, is mineralized with calcite, quartz, sphalerite, and chalcopryrite. A selected sample assayed: gold, trace; silver, 1.67 ounces a ton; copper, 5.41%; lead, none; zinc, 14.36%.

Associated minerals or products - **Copper, silver,**
Minéraux ou produits associés

HISTORY OF EXPLORATION AND DEVELOPMENT
HISTORIQUE DE L'EXPLORATION ET DE LA MISE EN VALEUR

This occurrence is located at the 6,000 foot elevation on the Sikanni Range, 3 miles southwest of the mouth of Quenada Creek.

HISTORY OF PRODUCTION/HISTORIQUE DE LA PRODUCTION

REFERENCES/BIBLIOGRAPHIE

Lord, C.S.; McConnell Creek Map-Area, Cassiar District, British Columbia; Memoir 251, p. 63, Geol. Surv. of Canada, 1948.

MAP REFERENCES/RÉFÉRENCES CARTOGRAPHIQUES

Map 962 A, McConnell Creek, B.C., (Geol.), Sc. 1":4 miles -
 accomp. Memoir 251.

Map 94 D, McConnell Creek, B.C., (Topo.), Sc. 1:250,000.

REMARKS/REMARQUES

Comp./Rev. By Comp./rév. par								
Date Date	02-67							