

NAME OF PROPERTY

NAHLIN FAULT

OBJECT LOCATED

UNCERTAINTY IN METRES (approx.) Lat. 59°05' Long. 133°15'

Mining Division Atlin District Cassiar

County Township or Parish

Lot Concession or Range

Sec Tp. R.

OWNER OR OPERATOR

DESCRIPTION OF DEPOSIT

High-grade magnesite occurs widely in the carbonatized zone along the southern contact of the Nahlin ultramafic body. The high-grade magnesite observed is very fine-grained, pure white material occurring as veins up to 4 feet wide in outcrops of carbonatized serpentinite.

HISTORY OF EXPLORATION AND DEVELOPMENT

Occurrences of high-grade magnesite have reportedly been traced from near the shore of Atlin Lake southeastward for a considerable distance.

HISTORY OF PRODUCTION

REFERENCES

Aitken, J.D.; Atlin Map-Area, British Columbia;
Memoir 307, p. 79, Geol. Surv. of Canada, 1959.
Annual Reports, Dept. of Mines, B.C.: 1915, p. 65.

MAP REFERENCES

Map 1082 A, Atlin, (Geol.), Sc. 1":4 miles - accomp.
Mem. 307.

Map 104 N, Atlin, (Topo.), Sc. 1:250,000.

REMARKS

Comp./Rev. By							
Date	9-66						