



Province of British Columbia
Ministry of Energy, Mines and Petroleum Resources

Open File 1986-7

**OCCURRENCE AND DISTRIBUTION
OF
PLATINUM-GROUP ELEMENTS
IN
BRITISH COLUMBIA**

**Compiled by V. J. Rublee
June 1986**

**This is a contribution towards the British Columbia- Canada Mineral Development
Agreement**

This open file report is preliminary and
may not conform to normal British Columbia
Geological Survey Branch editorial standards
for publications.

TABLE OF CONTENTS

	PAGE
INTRODUCTION	3
ALASKAN-TYPE PLATINUM-GROUP ELEMENT OCCURRENCES AND THEIR DERIVED PLACERS ..	4
Tulameen Complex	6
ALPINE-TYPE PLATINUM-GROUP ELEMENT OCCURRENCES AND THEIR DERIVED PLACERS...	17
Scottie Creek	18
Keefers	21
Mastodon	22
CU-NI SULPHIDE PLATINUM-GROUP ELEMENT OCCURRENCE.....	25
Tofino Nickel	26
Kitkat	29
Giant Mascot	30
Swede	34
Dykes Option (Mineral Dyke)	36
MISCELLANEOUS PLATINUM GROUP ELEMENT OCCURRENCES	38
i) Hydrothermal Related Cable Claim	39
Nome Claim	40
Mt. Ida	42
Mother Lode (Burnt Basin).....	45
ii) Alkalic Hosted	47
Franklin Camp	48
Sappho	56
iii) Skarns, Porphyry Copper, Stratabound Lead Zinc	58
Sullivan	58
Nickel Plate and Copper Mountain	59
ACKNOWLEDGMENTS	73
BIBLIOGRAPHY	74

LIST OF TABLES

1. Noble metal contents by lithology (weighted mean values) in the Tulameen Complex (after St. Louis, 1984) 10
2. Assays of platinum obtained from "black lead" in the Franklin Camp and drainage channels (from Thomlinson, 1928) 51
3. Platinum-group element placer deposits in British Columbia61

LIST OF FIGURES

1. Alaskan and alpine-type ultramafic complexes in British Columbia5
2. Generalized geology of the Tulameen Mining Camp (after Findlay, 1969) ...8
3. Distribution of the noble metals in the Tulameen Complex by lithology (after St. Louis, 1984) 9
4. Generalized geology of the Franklin Camp (after Drysdale, 1915)50

OPEN FILE MAP 1986-7 1:2 000 000 SCALE
Platinum Group Element Occurrences in British Columbia

INTRODUCTION

Platinum-group occurrences in the Canadian Cordillera are poorly documented, and with the exception of the Tulameen Ultramafic complex, superficially studied. The main sources of information available on the platinum potential in British Columbia were investigations undertaken by the Munition Resources Commission in 1920, and the Canada Department of Mines in 1934. From this framework, publications by the Geological Survey of Canada, British Columbia Ministry of Energy, Mines and Petroleum Resources, United States Geological Survey, Industrial Sector, and unpublished thesis material aided in substantiating, and detailing the platinum-group mineral inventory of British Columbia.

The classification of platinum-group element deposits used in this report is given in the following Table:

Primary Classification of Platinum-Group and by-Product deposits (after Cabri, 1981):	
A. PGE-dominant deposits	B. PGE Co-product and bi-product deposits
1. Merensky-type	1. Alpine-type and derived placers
2. Layered Chromite	2. Ni-Cu magmatic sulphides
3. Alaskan-type and derived placers	3. Miscellaneous deposits

In British Columbia, platinum-group mineralization occurs within Alaskan-type and derived placers, Alpine-type and derived placer, Ni-Cu magmatic sulphide, and Miscellaneous deposits.

ALASKAN-TYPE PLATINUM-GROUP ELEMENT OCCURRENCES AND THEIR DERIVED PLACERS

Alaskan-type ultramafic complexes are crudely zoned, steeply dipping intrusions emplaced into either stable platform (hence 'Aldan'-type zoned complexes) or orogenic ('Ural'-type) environments. The bodies, roughly circular in cross section, consist of dunite to peridotite cores rimmed by a pegmatitic hornblendite, with successive intermediate shells of peridotite, olivine pyroxenite, hornblende pyroxenite and clinopyroxene hornblendite. The alkalic petrochemistry, highly calcic pyroxenite, abundance of hornblende, and absence of orthopyroxene and plagioclase, characterize and differentiate Alaskan-type bodies from Alpine ultramafic and stratiform intrusions (Crockett, 1971).

Two parallel north-northwest trending belts containing zoned ultramafic bodies occur in the northern Cordillera (Fig. 1). The most westerly belt, which occurs within the Alexander Terrane, consists of 52 zoned plutons which extend for 560 kilometres from Kodiak Island to south of Prince Rupert, British Columbia. The Turnagain, Polaris and Tulameen complexes within the Quesnel Terrane constitute the eastern, British Columbia Belt.

Worldwide, this type of ultramafic rock suite is a significant source of platinum-group elements (specifically platinum, osmium and iridium) and may carry up to 10 grams per tonne total PGE (Crockett, 1982). Pt-Fe alloys, platiniridium and osmiridium are associated with chromitiferous forsterite dunites and pyroxenites, as well as sulphide and oxide phases as in the case of the Alaskan panhandle bodies. Derived placers are a significant source of platinum, palladium, osmium, ruthenium and iridium, as exemplified by the Goodnews Bay (Alaska), and Tulameen Complexes (British Columbia) from which total PGE produced exceeded 647,500 ounces and 20,000 ounces respectively (Barker et. al.; 1982; Mertie, 1969). The Salt Chuck Mine on Duke Island in the Alaskan panhandle, produced in excess of 661,771 ounces of platinum and palladium as a by-product from copper ore (Barker et. al., 1982).




Figure 1. Alaskan and Alpine-Type Ultramafic Complexes in British Columbia (after McTaggart, 1971).

TULAMEEN COMPLEX

LOCATION

The Tulameen complex, exposed over an area of 60 square kilometres, is located on the Tulameen River, a tributary of the Similkameen River, approximately 24 kilometres east of Princeton, at the boundary between the Intermontane and Cascade Fold Belts.

REGIONAL GEOLOGY

The complex is a zoned, southeasterly elongated ultramafic-gabbroic body that has been emplaced into upper Triassic Nicola group metasedimentary and metavolcanic rocks. It is unconformably overlain by terrigenous sedimentary rocks and andesitic to basaltic flows of the Eocene Princeton Group. The Jurassic Eagle granodiorite, a phase of the Mount Lytton Batholith, lies to the west. The ultramafic complex has been assigned a mid Jurassic age on a basis of K-Ar determinations of 174 ± 4 Ma on hornblende by (Roddick, 1970), and of 186 Ma on biotite by (Leech, 1963).

LOCAL GEOLOGY

Ultramafic rocks within the Tulameen intrusive form assymmetrically zoned, steeply dipping plugs, enclosed by an older alkalic (potassium rich, silica undersaturated) gabbroic suite (Findlay, 1969). Findlay describes the distribution of the three areas of ultramafic rocks within the complex as follows:

"In the northern part of the complex, the ultramafic units display the characteristic zonal pattern of similar intrusions in Alaska and U.S.S.R., comprising a dunite core surrounded by shells of olivine pyroxenite and hornblende clinopyroxenite. South of the Olivine Mountain, where dunite is not exposed, the two main ultramafic zones contain a median zone of olivine clinopyroxenite bounded by hornblende clinopyroxenite. In the Tanglewood Hill area, hornblende clinopyroxenite is the ultramafic rock exposed."

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

(i) Lode Occurrences

A Study of the geochemistry of PGE within the Tulameen complex, undertaken by St. Louis (1984), demonstrated the distribution of PGE to be a function of the degree of differentiation (zoning) within the ultramafic intrusive. The highest platinum, iridium, osmium and rhodium values were obtained from dunites, peridotites, and their altered (serpentinized) equivalents; the pyroxenites and hornblendites were depleted (St. Louis, 1984). Palladium appears to be confined to the marginal phases (hornblende clinopyroxenite and hornblende) and enclosing gabbroic rocks (Fig. 3).

In addition to studying the relative abundance of the PGE in the various lithologies, St. Louis analysed the PGE. The analytical results of the quantitative analyses, presented in Table 1, clearly demonstrate Pt, Ir, Os and Rh enrichment in the chromite-rich and serpentinized samples.

Mineralogical study by (St. Louis,) of the chromite-rich rocks (dunite) defined three modes of occurrence of the PGE: 1) as inclusions (predominantly Pt-Fe alloys - Pt_xFe , tulameenite - Pt_2FeCu , platinum, and genkinite - $(Pt,Pd)_4Sb_3$) within chromite grains, 2) as anhedral grains (commonly sperrylite $PtAs_2$, irarsite- $IrAsS$, and associated Fe-Ni sulphides) interstitial to chromite grains and uncommonly 3) as platinum group metals exsolutions from chromite grains.

Platinum group elements and observed sulfide enrichment in serpentinites and serpentinized dunites may indicate remobilization and concentration of the PGE as well as Ni, Fe, As and S during the process of serpentinization (St. Louis, 1984). The restricted distribution/partitioning of palladium into the least mafic phases (later stage differentiates) may be attributed, as proposed by St. Louis, to the siderophile character of palladium or its affinity for more siliceous rocks. The significance of temperature control on the distribution of the platinum group elements in the Tulameen Complex is suggested by the association of Pt, Ir, Rh and Os with high temperature chromite and the partitioning of Pd into the lower temperature (less refractory) phases. As described by St. Louis, (1982, page 221):




Figure 2. Generalized geology and rivers and tributaries with placer platinum in the Tulameen area (after Mertie, 1969).




Figure 3. Distribution of the Noble Metals in the Tulameen Complex (after St. Louis ,1984).

TABLE 1

NOBLE METAL CONTENTS BY LITHOLOGY (WEIGHTED MEAN VALUES IN PPM AT 95% CONFIDENCE LEVEL) AFTER ST. LOUIS, 1984

Lithology	Pt	Ir	Au	Os	Pd	Rh
Dunite and peridotite	55.9+1.7 (30)	0.103+0.001 (33)	0.174+0.002 (10)	BLQ	BLQ	1.01+0.02 (31)
Serpentinite and serpentinite-dunite	150.4+3.6 (10)	2.62+0.01 (11)	0.22+0.001 (10)	2.6+1.1 (2)	BLQ	1.47+0.05 (11)
Dunite, peridotite, serpentinite, and serpentinite-dunite	72.8+1.5 (40)	0.105+0.001 (44)	0.209+0.002 (43)	2.6+1.1 (2)	BLQ	1.13+0.02 (42)
Olivine clinopyroxenite and clinopyroxenite	33.7+3.6 (2)	0.15+0./01 (1)	0.20+0.01 (2)	BLQ	BLQ	0.45+0.06 (2)
Hornblende clinopyroxenite and hornblendite	16.4+5.9 (1)	0.01+0.01 (1)	0.30+0.01 (1)	BLQ	24.5+15.9 (1)	0.1+0.1 (1)
Magnetite-rich	37.6+2.8 (7)	0.25+0.011 (4)	0.19+0.01 (7)	1.9+0.4 (5)	42.9+17.8 (1)	0.34+0.05 (5)
Syenogabbro and syenodiorite	24.4+5.1 (2)	0.10+0.01 (3)	0.10+0.01 (1)	BLQ	BLQ	0.2+0.1 (1)
Sulfide-rich	43.9+3.3 (4)	0.13+0.01 (3)	1.15+0.01 (5)	1.8+1.0 (1)	55.3+7.2 (5)	0.23+0.05 (3)
Chromite-rich	837.5+10.8 (6)	24.9+0.1 (7)	2.01+0.03 (7)	23.1+4.5 (3)	BLQ	47.8+0.1 (7)
Average ¹	8.79+0.92	0.032+0.002	0.039+0.002	0.39+0.11	14.47+3.29	0.09+0.02
Average ²	57.7+1.1	0.106+0.001	0.204+0.002	1.98+0.38	49.2+6.2	0.89+0.02

¹weighted on the volume percentages of the lithologies²weighted on the number of samples of each lithology

Number of samples given in parenthesis

"Overall, Findlay (1963) found the highest concentrations of Pt in dunite and peridotite. In rock samples, he found the highest Pt content to be 0.225 gram per short ton in dunite on Olivine Mountain. The background Pt content in the dunite is 0.08 to 0.09 gram per short ton (Findlay, 1963). Chromite segregations within the ultramafic rocks returned the highest value of 7.34 grams per short ton Pt".

OTHER PLATINUM LODE OCCURRENCES IN THE TULAMEEN DISTRICT

Platinum values of 4.29 grams per tonne (Kemp, 1902) were obtained from sheared granites along Siwash Creek, which enters the Tulameen River 4.8 kilometres upstream from Eagle Creek. An assay of 137 grams per tonne platinum (O'Neill and Gunning, 1934) was returned from a trial carload of copper ore mined from skarnified Nicola limestones at Law's Camp, due north of Grasshopper Mountain. Copper Sulphide mineralization in a 60 foot greenstone dyke on Newton Creek, a tributary of Granite Creek, was analysed and returned a value of 8.57 grams platinum per tonne of sulphide.

(ii) Placer Occurrences

From 1889 to 1891 the Tulameen District, with an average annual output of 1500 ounces, was recognized to be the most important producer of platinum in North America; all production came from placer operations. Production continued intermittently until 1936 with an estimated total output of 20,000 ounces (Uglow, 1920). Rice (1947) describes the occurrence of platinum in the placers as follows:

"The ratio of platinum to gold is about 1 to 4 in the Similkameen and lower reaches of the Tulameen River, but increases upstream to where, near the mouth of Olivine Creek, the platinum is about equal in amount to, or may even exceed, the gold. On the whole, however, the platinum is of lesser importance. Platinum nuggets are never flattened and rarely even sub-angular, but occur in small rounded grains resembling fine shot, and mostly of uniform size. These nuggets are smaller than the gold on the average, but at the same time the platinum does not occur in fine, flaky particles. Most of the pellet-like nuggets have a roughly pitted surface; many show adhering grains of chromite and magnetite; and olivine and pyroxene are occasionally attached to them."

Alloys of Pt and Fe comprise the major platinum group metals in the placer; with minor Ir, Rh, Os, Cu, Pd and Ni substitute for Pt (Raicevic and Cabri, 1976). Palladium contents are not significant. Figure 2 shows the location of platiniferous creeks and rivers in the Tulameen District.

TULAMEEN DISTRICT

SELECTED BIBLIOGRAPHY

- Berry, T. F. (1963): Concentration Tests on Tulameen River Sand from the Brigitte Mining and Consulting Company limited, Vancouver, British Columbia, Department of Mines and Technical Surveys Canada, Mines Branch Investigation Report IR 56-87, 7 pages.
- Bilquist, Ron (1979): Report on Prospecting Survey of Grasshopper 1 and 2 Mineral Claims B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 7944.
- Bilquist, Ron and Culbert, R. R. (1982): Report on the Geochemical Survey of Grasshopper 1 and 2 Mineral Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 10063.
- Cabri, L. J. (1973): Mineralogical Investigation of a Placer Sand Sample from the Tulameen Area, British Columbia, Department of Mines Technical Surveys, Canada, Mines Branch Investigation Report IR 73-29, 14 pages.
- Cabri, L. J., Owens, D. R. and LaFlamme, J. H. G. (1973): Tulameenite, a new platinum-iron-copper mineral from placers in the Tulameen River Area, British Columbia, Canadian Mineralogist, Volume 12, pages 21-25.
- Camsell, Charles, (1910): Platinum Mining in the Tulameen District, B.C. Canadian Mining Institute, Quarterly Bulletin 9, Pages 29-44.
-(1913). Geology and Mineral Deposits of the Tulameen District, British Columbia. Geological Survey of Canada, Memoir Number 26, 188 pages.
- Carvalan, F. R., (1984): Prospecting Report on the J and L 1, 2, 3 and 4 claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12121.
-(1984). Report on Geochemical and Geophysical Surveys, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12190.
- Findlay, D. C. (1963): Petrology of the Tulameen Ultramafic Complex, Yale District, British Columbia; Unpublished Ph.D. Thesis, Queens University, 415 pages.
-(1965). Platinum in the Tulameen Ultramafic Complex, B.C., Geological Survey of Canada, Paper 65-2, page 20.
-(1969). Origin of the Tulameen Ultramafic-Gabbro Complex, southern British Columbia, Canadian Journal of Earth Sciences, Volume 6, pages 399-425.

SELECTED BIBLIOGRAPHY

- Freeland, P. P. (1925): Olivine Mountain, Minister of Mines, B.C., Annual Report 1924, pages B170-176.
-(1927). Tulameen River Section, Minister of Mines, B.C., Annual Report 1926, pages A228-234.
-(1928). Placer Mining in the Southern District, Minister of Mines, B.C., Annual Report 1927, pages C256-257.
-(1929). Tulameen River Section, Minister of Mines, B.C., Annual Report, pages C268-272.
-(1930). Placer Mining in the Southern District, Minister of Mines, B.C., Annual Report 1929 pages C280-281.
-(1931). Southern Mineral Survey District Number 4, Minister of Mines, B.C., Annual Report 1930, pages A210-213.
- Galloway, J. D. (1914): Similkameen Mining Division, Minister of Mines, B.C., Annual Report 1913, pages K225-227, K234-235.
- Ryback-Hardy, V. (1983): Geochemical and Geological Report on the J and L Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 11666.
- Hunter, Hugh (1918): Similkameen Mining Division, Minister of Mines, B.C., Annual Report 1917, pages F216-217.
-(1919). Tulameen River, Minister of Mines, B.C., Annual Report 1918, pages K213-214.
- Jones, Harold M. (1983): Report on the Hand H Claim Group-Olivine Mountain, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 11763.
- Kemp, J. F. (1902): The Geological Relations and Distribution of the Platinum and Associated Metals, United States Geological Survey, Bulletin Number 193, Series A, Economic Geology, 14, pages 34, 38-51.
- Macaulay, R. M. (1919): The Source of Placer in the Tulameen District of British Columbia, Engineering and Mining Journal, Volume 107, pages 303-336.

SELECTED BIBLIOGRAPHY

- Mackenzie, G.C. (1920): Mineral Investigations - Platinum, Munition Resources Commission Canada, Final Report, pages 147-156.
- Maltby, P. D. R. (1961): Recovery of Iron, Gold and Platinum From a Sample Submitted by G. R. A. Dorphan Mining Company Limited, Tulameen, British Columbia, Department of Mines and Technical Surveys, Canada, Mines Branch Investigation Report IR 61-69.
- Naldrett, A. J. and Cabri, L. J. (1976): Ultramafic and Related Rocks; Their Classification and Genesis with Special Reference to the Concentration of Nickel, Sulfides and Platinum Group Elements, Economic Geology, Volume 71, pages 1131-1158~
- O'Neill, J. J. (1918): The Platinum Situation in Canada, Geological Survey of Canada, Summary Report 1918, Part G, pages 2G, 7G-12G.
- O'Neill, J. J. and Gunning H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, pages 89-98.
- Peto, Peter (1983): Prospecting Report on the Cameron and Reay's Claims (Olivine Mountain), B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12481.
- Poiterin, Eugene (1924): Platiniferous Rocks from Tulameen Map Area, Yale District, British Columbia and Ural Mountains, Russia, Geological Survey of Canada, Summary Report, 1923, Part A, pages 84-101.
- Raicevic, D. and Bruce, R. W. (1975): Determination of Gold and Platinum Contents in a Low-Grade Alluvial Gravel From the Tulameen River Area of British Columbia, Department of Mines and Technical Surveys Canada, Mines Branch Investigation Report, IR 74-49, 28 pages.
- Raicevic, D. and Cabri, L. J. (1976): Mineralogy and Concentration of Au and Pt Bearing Placers from the Tulameen River Area in British Columbia, Canadian Institute of Mining and Metallurgy Bulletin, June 1976, pages 111-119.
- Rice, H. M. A. (1947): Geology and Mineral Deposits of the Princeton Map-Area, British Columbia, Geological Survey of Canada, Memoir 243, 136 pages.

SELECTED BIBLIOGRAPHY

- Robertson, W. F. (1899): Platinum, Minister of Mines, B.C., Annual Report, pages 558-560.
- Roddick, J.C. (1970): The Geochronology of the Tulameen and Hedley Complexes, British Columbia, Unpublished M.Sc. Thesis, Queen's University, 128 pages.
- St. Louis, R. M. (1982): Platinoids in the Tulameen Ultramafic Complex, British Columbia Department of Energy, Mines and Petroleum Resources, Fieldwork 1981, pages 218-222.
- (1984): Geochemistry of the Platinum Group Elements in the Tulameen Ultramafic Complex, British Columbia, Unpublished M.Sc. Thesis, University of Edmonton, 127 pages.
- Uglow, W. L. (1920): Possibilities for Platinum in Western Canada, Canadian Mining Institute, Monthly Bulletin Number 95, pages 210-214.

ALPINE-TYPE AND DERIVED PLACERS

Alpine-type ultramafic bodies are dismembered basal portions of ophiolites that have been tectonically emplaced in an orogenic belt. Lenses and pods of dunites and peridotites serpentized to varying degrees are stretched out along major fault zones such as the Coquihalla, Hozameen, Fraser, Pinchi and Nahlin, all of which are associated with oceanic volcanic and sedimentary rocks of the Cache Creek and Slide Mountain Assemblages (Figure 1).

The concentrations of platinum-group elements in Alpine-type bodies, lower than those in Alaskan-type complexes, are closely associated with chromites and may be associated with magnetite and sulphides. The predominance of osmium and iridium and the absence of palladium in the chromitiferous dunites and peridotites reflects the low degree of differentiation within the magma as "an increasing Pt:Ir ratio is the characteristic trend expected in the normal crystallization in mafic magmas" (Crocket, 1982). The affinity of Os, Ir and Ru for chromites, which occur as lenses, pods and schlieren within the basal cumulus dunites has been attributed to their substitution for Cr^{3+} in chrome spinel (Crocket, 1982).

Within the cumulus gabbros, which overlie the tectonized harzburgites and lherzolites that compose the basal portion of the ophiolite, minor sulphide concentrations may provide a significant target for Pt and Pd. The process of fractional crystallization increases Pt and Pd concentrations, hence sulphide saturation of the magma "could potentially scavenge Pt and Pd when their concentration was at a maximum" (Crocket, 1982, page 57).

SCOTTIE CREEK

LOCATION

The chromite deposit on Scottie Creek is located approximately 6.4 kilometres from its confluence with the Bonaparte River, 32 kilometres north of Ashcroft.

REGIONAL GEOLOGY

The Scottie Creek property is situated within the Intermontane Belt near the eastern margin of the Cache Creek Assemblage. This eastern Cache Creek Belt of Pennsylvanian to Triassic radiolarian ribbon cherts, limestones, pillowed basalts, gabbro and serpentinized ultramafic rocks are bounded to the east by Eocene to Miocene Kamloops Group dacites and basalts and to the west by the Fraser River Fault. Small bodies of serpentinized dunites and peridotites are associated with the Cache Creek rocks, and occur in a northerly trending belt approximately 1/4 mile wide, that outcrops on "both sides of Bonaparte Valley from south of Cache Creek to Clinton" (Duffell and McTaggart, 1952, page 76). According to Wright et al. (1982), 85 ultramafic bodies have been mapped, ranging in size from a few to 300 metres in maximum dimension. The smaller bodies are extensively sheared, slickensided and completely serpentinized whereas the larger ones at Fergusson and Scottie Creeks are relatively unshattered. "Dunite forms more than 85% of the ultramafic rocks, and peridotite more than 10%. Dyke-like pyroxenite bodies appear to cut across dunite and peridotite," (Wright et al., 1982, page 1169) at Fergusson and Scottie Creeks.

LOCAL GEOLOGY

Chromite occurs as disseminated grains, massive lenses and tabular sheets within the irregularly spaced pods of faulted, serpentinized dunites. As described in the British Columbia Minister of Mines Annual Report for 1915 (page 285):

"The portion of Scottie Creek that the writer examined carefully is about three miles and a half above the mouth, where a deposit of chrome-iron ore occurs in the first range of hills back from the creek on the north side on right limit. These hills are made up chiefly of ridges and knolls of altered peridotite, carrying a large percentage of chrysolite, and decomposed magnesian rocks, greasy and very light colored, with tints of green, yellow and brown. The erosion has

LOCAL GEOLOGY (CONTINUED)

been so extensive as to cut a series of deep gulches through the hills, leaving many of the knolls isolated from the main range of hills and with sharp peaks and precipitous sides. This erosion has laid bare the iron ore on the sides of some of the knolls and the points remaining between the gulches, so that the outcroppings can be followed from point to point in an easterly direction for a considerable distance. Sometimes the outcroppings occur as solid masses or beds of ore; at other placers as nodules of varying sizes, some quite large, mixed with the decomposed rock."

As Wright et al. (1982, page 1169) observed: "At Scottie Creek, most chromite layers strike southwesterly and dip gently northwest, whereas at Ferguson Creek, 2.5 kilometres south of Scottie Creek, layers strike northerly and dip 50° east. This would be expected in separate blocks in a melange."

Southwest of the chromite bodies exposed in Chrome Creek are outcrops of Miocene basalts and augite andesites.

DISTRIBUTION AND OCCURRENCES OF PLATINUM-GROUP ELEMENTS

Samples of chromite ore obtained from deposits on Chrome Creek (north fork of Scottie Creek) returned values of 3.4 and 0.69 grams per tonne platinum (Thomlinson, 1920, page 179). Panned samples from bars and old diggings on Scottie Creek located 2.4 and 5.6 kilometres upstream from the confluence of Scottie Creek and Bonaparte River returned values of 4.8 and 1.37 grams per tonne platinum (Thomlinson, 1920, page 178). Two panned samples taken 0.4 and 0.8 kilometre up from the mouth of Chrome Creek produced 2.06 ounces per ton platinum each (Thomlinson, 1920, page 178).

SELECTED BIBLIOGRAPHY

- Duffell, S. and McTaggart, K. C. (1952): Ashcroft Map-Area, British Columbia, Geological Survey of Canada, Memoir 262, Pages 76-77, 96-99.
- Mertie, John B. Jr., (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, Page 29.
- Minister of Mines, British Columbia: Annual Reports, 1901, Page 1091; 1902, Page 198; 1915, Page 285; 1918, Page 227; 1920, Page 24; 1927, Page 211; 1928, Pages 220, 229; 1930, Page 198; 1931, Page 110.

SELECTED BIBLIOGRAPHY (CONTINUED)

- O'Neill, J. J. and Gunning, H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, page 102.
- Reinecke, L. (1920): Mineral Deposits between Lillooet and Prince George, British Columbia, Geological Survey of Canada, Memoir 118, page 86-91.
- Stevenson, J. S. (1941): Chromite Bulletin (Unpublished), British Columbia Department of Mines.
- Thomlinson, W. M. (1920): Platinum Mineral Investigations, Munition Resources Commission Canada, Final Report, page 177-180.
- White, G. E. (1980): Magnetometer Survey Report on the Flint, Scottie Creek Property, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 7859.
- Wright, R. L., Nagel, Joe and McTaggart, K. C. (1982): Alpine Ultramafic Rocks of South Western British Columbia, Canadian Journal of Earth Sciences, Volume 19, pages 1156-1173.
- Zbitnoff, George (1981): Diamond Drilling Report on the Barbara Claim, British Columbia Assessment Report 10208.

KEEFERS

LOCATION

The Keefers PGE occurrence is located on the north side of Nahatlach River, approximately 6.5 kilometres northwest of its confluence with the Fraser River, 27 kilometres due south of Lytton. The workings are located at 370 metres elevation.

REGIONAL GEOLOGY

In the vicinity of Keefers, a northwest trending serpentine belt occurs within a sequence of greenstones, quartzites, limestones and phyllites. It is bounded by the Cretaceous age granitic rocks of the Coast Plutonic complex on the west, and the Fraser River Fault on the east.

DISTRIBUTION AND OCCURRENCE OF PLATINUM-GROUP ELEMENTS

A description of the property geology and mineralization is unavailable, except for a brief account provided by the National Mineral Inventory (1975), as follows:

"The showings were discovered and staked by D. Pride in 1954; some open cutting was done at that time. Northwest Ventures Ltd. optioned the 23 claim property in 1955 and some surface work and diamond drilling was reported. A representative sample, sent by the company to Sudbury for assay, is reported to have returned values of 1.42 percent copper, 2.24 percent nickel, 0.15 percent cobalt, and 0.11 ounce platinum and 0.023 ounce palladium per ton."

SELECTED BIBLIOGRAPHY

Duffell, S. and McTaggart, K. C. (1952): Ashcroft Map-Area, British Columbia, Geological Survey of Canada, Memoir 262, Pages 24-25.

National Mineral Inventory (1975): Keefers, 92I/4 - Ni, Mineral Policy Sector, Department of Energy, Mines and Resources, Ottawa.

MASTODON

LOCATION

The Mastodon chromite property is located 4.8 kilometres southeast of Christina Lake and 19.2 kilometres southeast of Grand Forks at an elevation of 900 metres on the southwest slope of Cascade Mountain. The best showings are located adjacent to the Grand Forks - Rossland Highway, 10 kilometres southeast of Cascade (Stevenson, J. S. 1941).

REGIONAL GEOLOGY

Mastodon is located near the eastern margin of the Quesnel terrane within the Omineca Crystalline Belt. Paleozoic to late Triassic Anarchist assemblage and early Jurassic Rossland group volcanic arc and marine sedimentary rocks are complexly faulted and intruded by Jurassic Nelson granites and Eocene Coryell alkalic plutons in the vicinity of the Mastodon property. Tectonically emplaced serpentized dunites and pyroxenites are confined to the Anarchist assemblage (Peatfield, 1978)

LOCAL GEOLOGY

Small Lenses of chromite occur within a serpentized dunite which is associated with greenstone breccia, tuffs and flows. The Jurassic Nelson granites and associated diorite and porphyry cut the ultramafic body to the south-southeast of Cascade Mountain. Chromite occurs as irregular disseminations, long streaks (schlieren) and small lenses within the ultramafic body. The larger chromite lenses are aligned within the hanging wall of a northeast striking, southwest dipping fault zone, (Stevenson, 1941); the ultramafic body is associated with the Paleozoic arc and oceanic facies rocks; hence Peatfield (1978) designated these chromite-nickel bearing ultramafic bodies as 'alpine-type'.

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Samples of chromite ore were taken by Wm. Thomlinson (1920, Page 106) from several crown granted claims that are underlain by serpentinite. The results are as follows:

Cerargyrite Crown Grant

Fine green serpentine with some oxides of iron and altered rock, from the face of an open-cut near the bottom and from upper sides of a lens of chromite. Platinum - nil.

General sample of chromite on dump and in the lens. Platinum - trace.

Midnight Crown Grant

From open-cut toward north end of claim banded chromite enclosed in fine green serpentine. Platinum - .514 grams per tonne.

Blacktail Fraction

From the main workings, open-cuts on the north lens of ore. Soft reddish-brown oxidized material from streaks in the serpentine and from the lenses of chromite. Platinum - trace.

Blacktail Crown Grant

From the main workings, open cuts on the north lens of ore. General sample of chromite being taken out for shipment. Platinum - 0.69 grams per tonne.

Mastodon Crown Grant

Fine line of open-cuts where chromite was being mined for shipment. Soft altered rock with some firm serpentine containing grains of chromite. Platinum - trace.

From all the main workings, open-cuts. General sample of chromite being mined for shipment. Platinum - trace.

Blacktail

From open-cuts on north lens of ore. Special sample of granular chromite enclosed in amethyst coloured spar. Platinum - trace.

SELECTED BIBLIOGRAPHY

British Columbia Department of Mines and Petroleum Resources:

Exploration in British Columbia, 1977, Page E12.

Mertie, John B. Jr. (1969): Economic Geology of the Platinum Metals,
United States Geological Survey, Open File 630, Page 29.

O'Neill, J. J. (1918): The Platinum Situation in Canada, 1918,
Geological Survey of Canada, Summary Report 1918, Part G, Page 9G.

O'Neill J. J. and Gunning, H. C. (1934): Platinum and Allied Metal
Deposits of Canada, Geological Survey of Canada, Economic Geology
Series Number 13, page 106.

Peatfield, G. R. (1978): Geologic History and Metallogeny of the
'Boundary District', Southern British Columbia and Northern
Washington, Unpublished Ph.D. Thesis - Queens University.

Stevenson, J. S. (1941): Chromite Bulletin - unpublished, British
Columbia Department of Mines.

Thomlinson, Wm (1920): Platinum - Mineral Investigations, Munition
Resources Commission, Canada, Final Report, Page 166.

Cu-Ni SULPHIDE OCCURRENCES OF PLATINUM-GROUP ELEMENTS

In this class of deposits, the platinum group elements palladium and platinum occur as sulphides and sulfarsenides associated with pentlandite, pyrrhotite and chalcopyrite mineralization in gabbroic and related rocks. Pentlandite and pyrrhotite act as concentrators for palladium (Crocket, 1982). A positive correlation between Pd and Ni is evidenced by 'the mutual dissolution of Ni and Pd from pyrrhotite at sub-solidus temperatures' (Stumpfel and Tarkian, 1982). Chalcophyrite has been suggested (Crocket, 1982) to be a concentrator for platinum, however this is as yet unproven.

TOFINO NICKEL (RAINEY-PEARL)

LOCATION

The Tofino Nickel property is located opposite Similar Island on the west side of Deer Bay, in Tofino Inlet. The property is reached by boat 25 kilometres east-northeast from Tofino on the west coast of Vancouver Island.

REGIONAL GEOLOGY

The property, located on the western edge of the Insular Belt, is underlain by amphibolitic gneisses of the West Coast Crystalline complex (Muller, 1977). The gneisses are considered to have been derived from migmatization of Paleozoic eugeosynclinal rocks during a major Jurassic plutonic event. A U-Pb date obtained from well foliated quartz plagioclase biotite gneiss at Grice Bay, 15 kilometres southwest of the Tofino Nickel property gave a late Paleozoic age of 264 Ma (Muller, 1977).

LOCAL GEOLOGY

Amphibolite is exposed as lenses and dyke-like bodies (Mason, 1984) within a feldspar gneiss of the West Coast Complex. According to Stevenson (undated):

"The (gneissic) feldspar porphyry is the main rock in the stripping and also in the area extending for at least one-half mile easterly from it. The amphibolite occurs as remnant lenses that strike northwesterly and dip approximately 60 degrees southwestward, an attitude that is steeper but which roughly corresponds to the slope of the hillside, an important feature to be considered in prospecting the showing. The amphibolite appears to be a metamorphosed basic rock, probably originally basaltic to andesitic lava. The feldspar porphyry is a white rock, slightly gneissic in texture but otherwise lacking structure; the gneissosity strikes northwesterly. Replacement of the amphibolite by the feldspar porphyry is well shown at the lower end of the stripping."

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Pyrrhotite, chalcopyrite, pyrite and magnetite mineralization with nickel, platinum and palladium values is associated with the amphibolitic rocks. As described by Stevenson (undated):

"The pyrite occurs as scattered patches with ill defined borders to the amphibolite. The pyrrhotite occurs as patches and thin veinlets that form a lace-work in the amphibolite. The chalcopyrite occurs as small distinct lenses of nearly pure mineral, which range from one-half inch to 4 inches in width and from a few inches to 18 inches in length. The chalcopyrite lenses are well defined in the mineralized area and appear to be later than the other sulphides. The sulphides occur in two areas in the stripping, one, towards the upper end, measuring approximately 4 feet by 12 feet by 2 feet deep and the other towards the lower end, measuring 4 feet by 10 feet by 2 feet deep. Elsewhere the amphibolite is either relatively barren of minerals or may contain a little pyrite. The amphibolite, probably as lava, appears to have received the sulphide mineralization before the formation of the feldspar porphyry.

Samples of the sulphides and mineralized amphibolite taken by Stevenson returned values of up to 6.7 grams per tonne platinum and 9.8 grams per tonne palladium. Eastwood (1963), obtained 6.2 grams per tonne palladium and trace platinum from a grab sample of rock well mineralized with pyrite and chalcopyrite.

SELECTED BIBLIOGRAPHY

- Carson, D. J. T. (1968): Metallogenic Study of Vancouver Island with emphasis on the Relationships of Mineral Deposits to Plutonic Rocks. unpublished Ph.D. Thesis - Carleton University, page 156.
- Eastwood, G. E. P. (1963): Tofino Nickel Group, Minister of Mines, British Columbia, Annual Report, 1963, page 116.
- Mason, I. M. (1984): Assessment Report - Geology, Geochemistry and Geophysics of the Nickel I, 2, 3 and Loren Mineral Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 13121.
- Muller, J. E. (1977): Evolution of the Pacific Margin, Vancouver Island, and adjacent regions, Canadian Journal of Earth Sciences, Volume 14, Number 9, pages 2062-2065.

SELECTED BIBLIOGRAPHY (CONTINUED)

Muller, J. E. and Carson D. J. T. (1986): Geology and Mineral Deposits of Alberni Map-Area, Geological Survey of Canada, Paper 68-50, pages 36, 38, 40.

Stevenson, J. S. (Undated): Preliminary Notes on the Rainy-Pearl Copper Prospect, Tofino Inlet, British Columbia Open File, 92F 14E.

KITKAT

LOCATION

The Kitkat claim block is located on the west side of the Nitinat River approximately 5 kilometres southeast of Mt. McQuillan. The property can be reached from either Port Alberni (29 kilometres northwest of Kitkat) or Youbou (32 kilometres southeast) along Crown Forest Industries Ltd's Nitinat-Main Road.

REGIONAL GEOLOGY

The Kitkat property is situated on the north-northwestern edge of the Cowichan-Horne Lake Uplift which exposes Paleozoic age island arc volcanics and sedimentary rocks of the Sicker Group. The area is complexly block faulted; it is bounded on the southwest and northeast by parallel eastward-directed thrust faults. Isoclinal folds with north-northwest trending axis are associated with the episode of uplift which occurred at the end of the Paleozoic (Muller and Carson, 1968).

LOCAL GEOLOGY

Specific details of the property geology are lacking because the showings are a very recent discovery. According to Muller and Carson (1968), Pennsylvanian and older eugeosynclinal rocks of the Sicker group underly the area in the vicinity of the claims.

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Values of up to 1.65 grams per tonne platinum and 4.85 grams per tonne palladium as well as anomalous cobalt, chromium, copper and nickel values were obtained from irregular lenses, up to 4.5 centimetres in width, of semi-massive to massive stringer and disseminated pyrite and pyrrhotite within Sicker Group (George Cross Newsletter, 1985) basaltic volcanics. Malachite and azurite staining occurs on the gossanous weathering surface.

SELECTED BIBLIOGRAPHY

George Cross Newsletter, Nexus Resource Corporation, Number 222 (November 19, 1985).

Muller, J. E. and Carson, D. J. T. (1968): Geology and Mineral Deposits of Alberni Map-Area, British Columbia (92F), Geological Survey of Canada, Paper 68-50.

GIANT MASCOT (PACIFIC NICKEL, BRITISH COLUMBIA NICKEL)

LOCATION

The Giant Mascot, formerly the Pacific Nickel nickel-copper mine, is located approximately 10 kilometres north-northwest of Hope, at the head of the Stulkawhits (Texas) Creek which flows eastward into the Fraser River. Access is gained to the property along an 8.2 kilometer gravel road which intersects the Trans Canada Highway about 16 kilometres north of Hope.

REGIONAL GEOLOGY

Giant Mascot is situated on the eastern margin of the Coast Plutonic Complex, 3 kilometres west of the boundary with the Cascade Fold and Thrust Belt. The Hope fault, which follows this boundary, juxtaposes high grade metamorphic rocks of the Custer gneiss, on the eastern side, with upper Paleozoic Settler Schists and early to Mid-Cretaceous diorite and tonalites of the Spuzzum pluton, on the west (Monger, 1985). The metamorphic grade facies in the vicinity of Giant Mascot is sillimanite-almandine grade (Grove and James, 1974).

LOCAL GEOLOGY

Northerly trending, regionally metamorphosed Settler schists are intruded by a stock-like, crudely zoned ultramafic body enclosed by genetically related mid-Cretaceous Spuzzum diorites and tonalites (Aho, 1956). According to McLeod et al. (1976, page 115):

"K/Ar ages from the Giant Mascot Ultramafic body (119-95 m.y.) suggest that it is slightly older than adjacent diorite and tonalite (78-89), which are shown to be members of the mid-Cretaceous Spuzzum Intrusions."

However, the contact relationships between the two intrusive suites are ambiguous (McLeod et al., 1976). The ultramafic, which extends for 2.9 kilometres in an east-west direction and 2.25 kilometers north-south, is comprised of seventeen dunite to peridotite-cored, pipe-like bodies enclosed within a pyroxenite mass. Hornblende to pegmatitic hornblende reaction margins, up to 100 metres wide (Aho, 1956), rim the pyroxenite. Later hornblende and dioritic dykes cut the complex. The pipe-like ultrabasic structures vary in size from 15 metres by 15 metres to 60 metres by 100 metres; they broadly trend N75°W and plunge steeply to the north.

OCCURRENCE AND DISTRIBUTION OF P.G.E.

Nickeliferous pyrrhotite, chalcopyrite and pentlandite are exclusively associated with the ultrabasic rocks and occur as massive to disseminated sulphides. The average grade of the sulphides (Aho, 1956) is 1.4 percent nickel, 0.5 percent copper, 1 percent chromium, 0.1 percent cobalt, 0.7 grams per tonne gold and 0.3 grams per tonne platinum metals.

Aho (1956) defined three types of ore bodies at Giant Mascot; zoned bodies of replacement origin, massive sulphide of magmatic derivation, and vein-like later stage mineralization. The zoned ore bodies are elliptical in cross-section and form sulphide cores with shells or rings parallel to the long axes of the dunite and olivine peridotite pipes with which they are associated. These steeply plunging ore zones exhibit nickel bearing cores and copper bearing outer zones (Aho, 1956). Massive orebodies are irregular in outline and occur as lenses between or along the ultramafic units within the intrusive (Groves and James, 1974); occasionally they grade into disseminated ore.

Chalcopyrite-rich, vein-like mineralization up to 5 centimetres in width is present within and adjacent to the orebodies; rarely, it extends into the country rock.

A second type of mineralization consists of, or occurs in, heavily mineralized late stage hornblendite and pyroxenite dykes which cut the peridotite. As described by Aho:

"The pyroxenite dykes locally grade into the hornblendite dykes; sulphides within both are unevenly distributed and tend to be concentrated into massive blocks that extend irregularly like replacements into unaltered wall rocks. The sulphides are chiefly pyrrhotite, with smaller amounts of nickel and copper than sulfides in the ore bodies. In places the "Dykes" grade into massive sulphide "Veins" ."

A sample taken from a pegmatitic massive pyrrhotite vein from the 1500 orebody assayed 2 grams per tonne platinum and 7.2 grams per tonne palladium (Eastwood and Waterland, 1966). The 1500 orebody possesses maximum horizontal dimensions of 60 metres by 20 metres and a vertical dimension of 105 metres. Tonnage and grade quoted for this deposit are 607,273 tonnes at 1.35 percent nickel and 0.45 percent copper (Christopher and Robinson, 1974). Christopher and Robinson (1974 page 112) summarized the geological setting of the 1500 orebody as follows:

"Ore in both hornblende peridotite and hornblende pyroxenite occurs as both massive and lacy types with a concentration of massive near bottom; ore has been either dammed or cut off by a flat hornblendite dyke; the footwall contains a breccia with diorite fragments in mineralized hornblende pyroxenite".

SELECTED BIBLIOGRAPHY

- Aho, A. E. (1954): Geology of Ore Deposits of the Property of Pacific Nickel Mines near Hope, British Columbia, Unpublished Ph.D Thesis, University of California-Berkley.
-(1956). Geology and Genesis of Ultrabasic Nickel-Copper-Pyrrhotite Deposits at the Pacific Nickel Property, Southwestern British Columbia, Economic Geology, Volume 51, Number 5, pages 444-481.
-(1957). Pacific Nickel Property, Canadian Institute of Mining and Metallurgy, Volume 2, pages 27-36.
- British Columbia Ministry of Energy, Mines and Petroleum Resources, Geology and Exploration in British Columbia, 1971, pages 267-268; 1972, pages 117-118; 1973, pages 131-132.
- Cairnes, C. E. (1924): Nickeliferous Mineral Deposits, Emory Creek, Yale Mining Division, British Columbia, Geological Survey of Canada, Summary Report 1924, Part A, pages 100A-105A.
- Christopher, P. A. and Robinson, J. W. (1974): Pride of Emory Mine, British Columbia Ministry of Energy, Mines and Petroleum Resources, Geology and Exploration in British Columbia, 1974, pages 105-113.
- Clarke, W. E. (1969): Geology and Ore Controls, Giant Mascot Mines Limited, Western Miner, Volume 42, Number 6, pages 41-46.
-(1971). Geology of Giant Mascot, Western Miner, Volume 44, Number 9 pages 22-25.
- Cockfield, W. E. and Walker, J F. (1933): The Nickel Bearing Rocks Near Choate, British Columbia, Geological Survey of Canada, Summary Report 1933, Part A, pages 66A-68A.
- Eastwood, G. E. P. and Waterland, T. M. (1966): Pride of Emory, Giant Mascot Mines Limited, Minister of Mines, B.C., Annual Report 1966, pages 58-59.
- Grove, E. W. and James, A. R. C. (1974): Report on the Giant Mascot Mines, British Columbia Ministry of Energy, Mines and Petroleum Resources, Open File.
- Hill, H. L. and Starck, L. P. (1960): The Giant Nickel Mines; Western Miner and Oil Review, Volume 33, November, pages 39-42.

SELECTED BIBLIOGRAPHY

- Horwood, Ii. C. (1936): Geology and Mineral Deposits at the Mine of British Columbia Nickel Mines Limited, Yale District, British Columbia, pages I-15.
- McLeod, J. A. (1975): The Giant Mascot Ultramafite and its Related Ores, Unpublished M.Sc. Thesis, University of British Columbia, 123 pages.
- McLeod, J. A., Vining, M. and McTaggart, K.C. (1976): Note on the age of the Giant Mascot Ultramafic body, near Hope, British Columbia, Canadian Journal of Earth Sciences, Volume 13, Number 8, pages 1152-1158.
- Mertie, J. B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, page 28.
- Minister of Mines, B.C., Annual Reports; 1924, pages B137-138; 1928, page C227; 1929, pages C239-241; 1930, page A204; 1934, pages F17-F19; 1954, pages A161-163; 1958, pages 55-56; 1959, pages 124-127; 1960, pages 87-88; 1961, pages 86-88; 1963, pages 89-92; 1964, pages 137-142; 1965, pages 213-217; 1967, pages 63-64; 1968, pages 76-77.
- Monger, J. W. H. (1969): Hope Map-Area, West Half (92HW 1/2), British Columbia, Geological Survey of Canada, Paper 69-47, pages 63-64.
- Muir, J.E. (1971): A Study of the Petrology and Ore Genesis of the Giant Nickel 4600 Orebody, Hope, British Columbia, University of Toronto, Unpublished MSc Thesis, 113 pages.

SWEDE

LOCATION

The Swede Property is located on Moresby Island on Swede Peninsula on the east side of Anna Inlet in Klunkwoi Bay. The prospect is accessed by air, approximately 60 kilometres southwest from Sandspit.

REGIONAL AND LOCAL GEOLOGY

The Swede claims are located on the western boundary of the Insular Belt, within the Wrangellia Terrane. The oldest rocks in the area are tholeiitic basalts, andesites and coarser diabase dykes of the Triassic Karmutsen Formation. These are unconformably overlain by massive and thinly bedded limestones and argillites of the upper Triassic to lower Jurassic Kunga Formation. The property is underlain predominantly by andesites and basalts which have been intruded by three episodes of diabase dykes (Wilson, 1965). Erratic mineralization occurs as disseminations, blebs and fracture fillings of chalcopyrite and minor bornite in diabase dykes of the Karmutsen formation.

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Platinum, with local assays to 0.34 grams per tonne, and trace palladium were reported to occur within sporadic bornite mineralization within the diabase country rock (Minister of Mines, B.C., Annual Report 1922, page G39). This reported occurrence of the platinum group elements has not been substantiated.

SELECTED BIBLIOGRAPHY

McConnell, R. G. (1909): Moresby Island, Geological Survey of Canada, Summary Report 1909, pages 78-79.

Mertie, J. B. Jr. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, pages 28-29.

Minister of Mines, British Columbia: Annual Reports, 1907, pages 69-70; 1920, page 44; 1921, page 39; 1923, page 43; 1928, pages 65-66; 1929, page 57.

SELECTED BIBLIOGRAPHY (CONTINUED)

National Mineral Inventory (1975): Brandy (D),(Swede)(Last Chance),
103B/12 Cu₂, Mineral Resources Branch, Department of Energy, Mines
and Resources, Ottawa.

Sutherland Brown, A. (1968): Geology of the Queen Charlotte Islands,
British Columbia Department of Mines, Bulletin Number 54, pages
215-216.

Wilson, G. A. (1965): Report of D. Group Claims Situated on Swede
Peninsula, Lockeport, Moresby Island, B.C. Ministry of Energy, Mines
and Petroleum Resources, Assessment Report 662.

DYKES OPTION (MINERAL DYKE)

LOCATION

The Dykes Option copper, platinum, palladium and nickel occurrence is located 1 kilometre northwest of the junction of Riondel Road with the Kootenay Bay - Crawford Bay Road (Hoy, 1980).

REGIONAL GEOLOGY

The property is situated on the western limb of the overturned Crawford antiform within the Kootenay Arc, a north trending regional structure of folded and thrust faulted Hadrynian to early Mesozoic age strata (Hoy, et al., 1985). The Dykes Option is contained within a north-south trending, 20 kilometres wide belt, of sillimanite-garnet-biotite grade metamorphism (Hoy, 1980).

LOCAL GEOLOGY

Calc-silicate gneiss with amphibolite, schist and impure marble layers of the Middle Cambrian Index formation of the Lardeau Group underlie the property. A flat lying diorite sill or dyke (Rice, 1941) has intruded the metasedimentary rocks. Pegmatite dykes cut the diorite intrusive (O'Grady, 1930).

OCCURRENCE AND DISRIBUTION OF PLATINUM GROUP ELEMENTS

Details of the mineralization and geology of the Dykes Option mineral showing are limited to brief descriptions by O'Grady (1930), Richmond (1931) and Rice (1941). Platinum, palladium, nickel and copper mineralization is reported to be present as disseminations within the diorite sill or dyke. The best showing, exposed in an 80 metre adit, occurs near the margin of the intrusive diorite (Rice, 1941); no assays are available.

SELECTED BIBLIOGRAPHY

- Hoy, T (1980): Geology of the Riondel Area, Central Kootenay Arc, Southeastern British Columbia, British Columbia Ministry of Energy, Mines and Petroleum Resources, Bulletin 73, pages 38, 49, 88.
- Hoy T., Berg, N., Fyles, J. T., Delaney, G. D., McMurdo, D. and Ransom, P. W. (1985): Stratabound Base Metal Deposits in Southeastern British Columbia, in, Geological Society of America, Field Guide to Geology and Mineral Deposits in the Southern Canadian Cordillera, Geological Society of America Cordilleran Section Meeting, Vancouver, British Columbia, May 1986, pages 11-1 - 11-32.

SELECTED BIBLIOGRAPHY

- O'Grady, B. T. (1930): Mineral dyke, Minister of Mines, B.C., Annual Report 1929, page C325.
- Rice, H. M. A. (1938): Preliminary Report, East Half Nelson Map-Area, British Columbia, Geological Survey of Canada, Paper 38-17, page 9.
- . . (1941). Nelson Map Area, East Half, British Columbia, Geological Survey of Canada, Memoir 228, page 60.
- Richmond, A. M. (1931): Mineral Dyke, Minister of Mines, B.C., Annual Report 1930, page A254.

MISCELLANEOUS PGE OCCURRENCES

Platinum group-elements have been detected in trace to appreciable amounts in hydrothermal, alkalic-hosted, skarn, porphyry copper, and stratabound base metal environments in British Columbia.

The mobility of platinum-group minerals at low temperatures in aqueous solutions, and the extraction from silicates, oxides and sulphides during the process of serpentinization, has been studied by Stumpfl and Tarkian (1976). The mobility is such that in extreme situations PGE may be concentrated in "locations determined by geological structure, without any spatial relation to igneous rocks" (Stumpfl and Tarkian, 1982).

i) Hydrothermal Platinum-group Element Occurrences

Assays obtained from quartz veins and shear zones from the Cable and Nome claims, Mt. Ida property, and former Mother Lode Mine, range from 0.6 to 3.4 grams per tonne platinum. All of these hydrothermally related showings, prospects and deposits occur in areas where there has been regional extensional tectonic events within the Omineca Crystalline Complex.

CABLE CLAIM

The Cable Claim, is situated between Woodbury and Sunrise Mountains, at an elevation of 2300 metres on a northern tributary of the Silver Spray Creek. This region now lies within the boundaries of Kokanee Glacier Park.

The area is entirely underlain by granites of the Jurassic Nelson Batholith. An assay of 2.40 grams per tonne platinum (O'Neill and Gunning, 1934), was obtained from an auriferous quartz vein within the granite. Further details of the geology, mineralization and workings are unavailable.

SELECTED BIBLIOGRAPHY

Cairnes, C. E. (1934): Sardon (Slocan and Ainsworth Mining Divisions), Kootenay District, British Columbia, Geological Survey of Canada, Map 273A.

... ..(1935). Description of Properties, Slocan Mining Camp, British Columbia, Geological Survey of Canada, Memoir 184, page 199.

O'Neill, J. J. (1919): The Platinum Situation in Canada, 1918, Geological Survey of Canada, Summary Report 1918, Part G, page 8G.

O'Neill, J. J. and Gunning H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, page 107.

Mertie, John B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, page 29.

i) Hydrothermally Related Occurrences

NOME CLAIM

LOCATION

The Nome claim is located on the northeast side of Klowala (Cariboo) Creek which flows northeastward into Keen Creek, the south fork of the Kaslo River. The claim lies outside the northern boundary of Kokanee Glacier Park.

REGIONAL AND LOCAL GEOLOGY

The Nome claim lies on the eastern boundary of the Omineca Crystalline Belt, within the middle Jurassic Nelson Batholith. The property is underlain by a coarse grained granitic phase of the batholith which is cut by narrow, basic dykes (Cairnes, 1934).

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Assays of 2.74 grams per tonne and 1.71 grams per tonne platinum were obtained from grab-samples from a rock slide of granodiorite containing "specks and small masses of a hard, dark-colored mineral which appeared to be almost free from weathering effects, more so than the rock itself", (Thomlinson, 1920, Page 169). According to Cairnes (1935):

"Workings include two adits about 200 feet apart (vertically), the lower follows a shear zone about 4 feet wide. The zone strikes north 35 degrees east, dips steeply to the northwest, and shows little mineralization other than pyrite which is partly oxidized near the surface. The upper adit lies 1300 feet to the southeast of the other and was inaccessible at the time visited (June, 1927), but appears, from the material on the dump, to have encountered somewhat similar conditions to those in the lower adit and is probably driven on a parallel shear".

SELECTED BIBLIOGRAPHY

Cairnes, C. E. (1935): Description of Properties, Slocan Mining Camp, British Columbia, Geological Survey of Canada, Memoir 184 pages 238-239.

Mertie, John B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional page 630, page 29.

SELECTED BIBLIOGRAPHY (CONTINUED)

- O'Neill, J. J. (1919): The Platinum Situation in Canada, 1918,
Geological Survey of Canada, Summary Report 1918, page 8G.
- O'Neill, J. J. and Gunning, H. C. (1934): Platinum and Allied Metal
Deposits of Canada, Geological Survey of Canada, Economic Geology
Series Number 13, page 107.
- Thomlinson, Wm. (1920): Mineral Investigations-Platinum, Munitions
Resources Canada, Final Report, page 169.

i) Hydrothermally Related Occurrences

MT. IDA

LOCATION

Three platinum group element occurrences - Bonnie Brae, Mount Ida and Sunset - have been documented on the north and northeast slopes of Mount Ida, 0.8 kilometres southeast of Salmon Arm.

REGIONAL AND LOCAL GEOLOGY

Mount Ida, situated within the Kootenay Terrane of the Shuswap complex, is tectonically bounded on the east by the Monashee complex and on the west by the Harper Ranch assemblage of the Quesnel Terrane.

Argillites and limestones of the Precambrian to Mississippian Sicamous Formation, which unconformably overly Cambro-Ordovician schists of the Silver Creek Formation, comprise the middle to upper stratigraphic portion of the Mount Ida group. This sequence is intruded by Cretaceous age granites of the Coast Intrusions which in turn are capped by basaltic lava flows of the Eocene Kamloops Group.

DISTRIBUTION AND OCCURRENCE OF PLATINUM GROUP ELEMENTS

Platinum values have been obtained from quartz filled shear zones along with copper, lead and zinc sulphides, within impure quartzites and schists, adjacent to dykes of feldspar porphyry. The mineralized zones occur as "a system of sub-parallel ore bodies 18 inches to 7 feet wide that generally strike north-east and dip from an angle of 45 degree to almost vertical towards the southeast" (Brewer, 1913).

Three mineralized localities exposed in workings on the north and northeast sides of Mt. Ida were sampled by W.F. Ferrier of the Munition Resource Commission in 1918 as a follow up to previously reported assays of 2.74 and 7.54 grams per tonne platinum. As reported by Ferrier in the Final Report of the Commission:

"The first claims sampled was the White Cliff on the northeast slope of Mt. Ida, at an elevation of about 3,150 feet. Two tunnels have been driven. The upper one is a cross-cut 122 feet in length which has just reach the foot-wall of a vein at the face. About 100 feet below this another tunnel, 24 feet in length, follows a vein the dimensions of which were not satisfactorily determined. The strike is apparently northeast with a dip to the southeast. A cut was taken along the western wall of the tunnel to determine if platinum is present.

Sulphides, chiefly sphalerite, with some pyrite and galena, occur somewhat sparingly in the quartz. A second sample, consisting principally of these sulphides, was chipped at random from both walls of the tunnel to ascertain if the platinum was closely associated with the sulphides".

The assay results of these samples returned values of 13.03 grams per tonne gold, 1.03 grams per tonne platinum, and 3.42 grams per tonne gold, and no values in platinum respectively (Ferrier, 1920).

"The Mountain View claim was next visited and a sample taken in an open-cut near the eastern boundary of the claim, across the full width of the vein. The strike here is about east and west, with a dip of 60° to the south, but Mr. Thornton stated that the vein had been traced through the White Cliff, running across four claims. The general character of the vein material resembles that seen at the White Cliff. As before, a second sample was taken here by cutting across the centre of the vein, which appeared to be the more highly mineralized portion".

Samples from the Mountain View claim assayed 12.0 grams per tonne gold, 6.86 grams per tonne platinum and 2.79 grams per tonne gold 0.69 grams per tonne platinum respectively (Ferrier, 1920).

The Miller tunnel, on the north side of Mt. Ida, was the location for two samples taken along the face of an exposed shear zone full of quartz stringers in a granitic rock. Ferrier (1920) reports:

"The face of the drift, which does not expose any walls, was sampled across its full width. Another sample was taken from a small highly mineralized streak in the face.

Sulphides, consisting of sphalerite, galena, chalcopryrite, and pyrite, appear to be more abundant in the quartz here than in that at the claims visited on the previous day".

Values of 8.23 grams per tonne gold and 0.69 grams per tonne platinum were recovered from the 4.75 foot cut across the full width of the face, and 4.80 grams per tonne gold and 1.03 grams per tonne platinum are reported from across the mineralized streak.

SELECTED BIBLIOGRAPHY

- Brewer, W. M. (1914): Minister of Mines, B.C. Annual Report 1913, pages 198-199.
- Ferrier, W. F. (1920): Platinum at Mt. Ida, near Salmon Arm, Kamloops, Munition Resources Commission, Canada, Final Report, pages 183-185.
- Jones, A. J. (1959): Vernon Map Area, British Columbia, Geological Survey of Canada, Memoir 296, page 148.
- Minister of Mines, B.C.: Annual Reports, 1899, pages 733-734; 1926, page 188; 1930, pages 183-184.
- Mertie, John B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, page 29.
- O'Neill, J. J. and Gunning, H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, pages 183-185.
- Peto, Peter (1983): Geochemistry Report on the Bonnie Brae Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12055.
- Tully, Donald W. (1980): Geochemical Soil Survey on the Arm #1, 2 - Ida #1, 2, 3, 4, Claim Group, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 8995.
- Tully, Donald W. (1982): Detailed Geochemical Soil Surveys at Selected Locations on the Ida 2 and 4 claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 10244.

i) Hydrothermally Related Occurrences

MOTHER LODE (BURNT BASIN)

LOCATION

The Mother Lode property located in the Burnt Basin area, is situated at the head of Josh Creek, a tributary of McRae Creek, about 1.6 kilometres southwest of Paulson Bridge on the Christina Lake - Kinnaird Highway.

REGIONAL GEOLOGY

Oceanic sedimentary and intercalated volcanics of the upper Paleozoic Mount Robert's formation (and possibly Upper Triassic Rossland Group) which underly the Mother Lode crown grant, are intruded to the northwest by Jurassic Nelson granites and to the southeast by the Tertiary Coryell batholith. This area is in a tectonic depression within the Omineca crystalline Belt, approximately 10 kilometres east of the Grand Forks Complex, which is bounded by Tertiary normal faults (Parrish et al. 1985).

LOCAL GEOLOGY

Limey argillites, limestones and greenstones of the Paleozoic Mount Roberts Formation are cut by granodiorite and basic gabbroic dykes related to the Jurassic Nelson and Eocene Coryell plutonic events. A quartz vein, ranging in size from two to six feet in width, contains pyrite, galena, sphalerite and some molybdenite; it is hosted by the greenstone, between two wide mafic syenitic dykes.

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Assays of the quartz vein performed by Baker and Company of Newark, New Jersey returned values from trace to 8.57 grams platinum per tonne (O'Neill and Gunning 1934, page 104). Subsequent resampling of the vein by the Geological Survey of Canada (O'Neill and Gunning, 1934) indicated a platinum content of 3.43 grams per tonne.

SELECTED BIBLIOGRAPHY

Mertie, John B. (1969): Economic Geology of the Platinum Metals, United Stated Geological Survey, Professional Paper 630, page 29.

Minister of Mines, British Columbia: Annual Reports 1901, page 1066; 1920, page 304; 1917, page 201; 1932, pages 123-124; 1968, page 236.

O'Neill J. J. (1919): The Platinum Situation in Canada, 1918, Geological Survey of Canada, Summary Report, **1918**, Part G, page 9G.

O'Neill J. J. and Gunning, H. C. **(1934)**: Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series, Number 13, page 104.

Parrish, R., Carr, S. and Parkinson, D. (1985): Metamorphic Complexes and Extensional Tectonics, Southern Shuswap Complex, Southeastern British Columbia in Geological Society of America, Field Guides to Geology and Mineral Deposits in the Southern Canadian Cordillera, May 1985.

Uglow, w. L. (1920): Possibilities for Platinum in Western Canada, Canadian Mining Institute Bulletin, March 20, page 212.

ii) Alkalic Hosted Platinum-Group Occurrences

Platinum values at the Franklin Camp and Sappho property near Greenwood are associated with copper mineralization occurring within marginal shonkinitic phases of Tertiary Coryell alkalic stocks in southeastern British Columbia. Both properties are underlain by oceanic volcanic and sedimentary rocks of Paleozoic to Jurassic age (and associated basic and ultrabasic bodies) of the Quesnel Terrane, that were intruded during the Jurassic time by Nelson granites and in Tertiary time by Coryell alkalic plutons. The Tertiary episode of igneous activity occurred during the waning stages of a period of Eocene extensional tectonics, manifested by the horst and graben structures observed from the Okanagan Valley eastward to Christina Lake. The Franklin Camp and Sappho Property are both located on boundaries of such structures.

Both occurrences are associated with greenstones and ultramafics and are proximal to regional normal faults; if the Paleozoic to Jurassic ultramafic rocks are the platinum group element source, perhaps the PGE were remobilized and leached during deformation and concentrated in contemporaneous alkalic Coryell plutons in the vicinity of the source rocks. Therefore, alkalic bodies in this tectonic setting may be targets for platinum-group element exploration.

ii) Alkalic Hosted Occurrences

FRANKLIN MINING DISTRICT

LOCATION

The Franklin Mining Camp is situated between the Franklin and Glouster tributaries of Burrel Creek, the eastern branch of the North Fork of the Kettle River. The camp is approximately 72 kilometres from Grand Forks. The area is accessed from Grand Forks via the paved Granby Creeks logging road, which follows the north fork of the Kettle River for approximately 40 kilometres to its junction with the Burrel Creek road at kilometre 44 (Cukor, 1982). A secondary dirt road leads into the Franklin Mining Camp.

The first claims in the area, the Banner and McKinley, were staked in 1896. Considerable development was carried out in the area through to 1906 by which time practically all the ground in the mineral belt was staked (Drysdale, 1915).

REGIONAL GEOLOGY

The Franklin Mining Camp is situated within the Omineca Crystalline Belt, near the tectonic boundary between the Quesnel and North America Terranes. The property lies within a tectonically depressed area west of the Granby River fault. This fault forms the eastern boundary of a Tertiary north-northeast trending regional structure, the Republic Graben, which extends southward across the International Boundary (Lisle and Seraphim, 1980).

According to Drysdale (1915), the basal formation of the Franklin Camp is comprised of Carboniferous Franklin Group greenstones, tuffs argillites and cherty quartzites and Glouster Formation crystalline limestone. The "Paleozoic" rocks occur as roof pendants through the Jurassic age Nelson plutonic complex. Church (pers. comm) correlates Drysdale's Franklin Group greenstones with the Knob Hill Group and argillites with the Permo-Carboniferous Attwood Group. The limestone, previously referred to as Paleozoic Glouster Formation by Drysdale has been redesignated as the Triassic Brooklyn Group (Church, 1985).

Sandstones, conglomerates and tuffs of the Eocene Kettle River Formation are overlain by augite syenites with shonkinite-pyroxenite of the Coryell Formation; these comprise the Penticton Group, which lithologically dominates the Franklin Camp.

LOCAL GEOLOGY

Mineral claims within the Franklin Camp have been staked along the intrusive contacts between the Paleozoic greenstones and limestone with the Nelson granites and Coryell augite syenites.

A pyroxenite unit, referred to by Drysdale (1915) as a skonkinite-pyroxenite, is interpreted to be a basic cumulate or marginal phase of the augite syenite intrusion. Locally known as the "Black Lead", this pyroxenite hosts the mineralization, which occurs as pods, veinlets and disseminations of chalcopyrite, pyrite and copper carbonates at the outer margins of the body.


According to Chilcott and Lisle (1980), the sulphides, chalcopyrite, pyrite, sphalerite and platinum with malachite and azurite present as oxides in a gangue of quartzite and greenstone on the Mountain Lion crown grant, and in a pyroxenite-syenite gangue on the Maple Leaf, Buffalo and Averill crown grants.

OCCURRENCE AND DISTRIBUTION OF PLATINUM GROUP ELEMENTS

Platinum values of up to 15.4 grams per tonne were recorded in 1915 from assays of 'Black Lead' ore obtained from workings on the Maple Leaf Claim (B.C. Annual Reports 1920, page 153; 1929, page C254).

The Maple Leaf claim as well as the Columbia, Ottawa, Evening Star, Iron Hill, Buffalo, Blue Jay, Averill, Mountain Lion, and Lucky Jack Claims all of which are located on the 'Black Lead' (figure 4) were sampled by Wm. Thomlison of the Munition Resource Commission in 1918; his assays substantiated the earlier platinum results. Thomlinson's assays of this study and later data obtained are listed in Table 2. Thomlinson (1918) concluded that the platinum values observed in the ore zone (Black Lead) are proportional to the primary copper sulphides mainly chalcopyrite; the richest copper ore yielded the highest platinum concentrations. Sperrylite (Pt,As₂) is closely associated with sulphides; it is the major platinum mineral.

Mertie (1969) noted the occurrence of platinum within a quartz vein in the Union Mine Camp, located west of the Maple Leaf Claim.



300 0 300 600 900 1200

Scale (m.)





-  Tertiary sediments and volcanics
-  Eocene Coryell augite syenite
-  Jurassic Nelson Batholith
-  Paleozoic to Triassic Anarchist group
Knob Hill, Attwood and Brooklyn Formations

Figure 4. Generalized geology and claims with platinum-group element occurrences in the Franklin Camp (after Drysdale, 1915).

TABLE 2

ASSAYS OF PLATINUM OBTAINED FROM 'BLACK LEAD' IN THE FRANKLIN CAMP, AND DRAINAGE CHANNELS

CLAIM	DESCRIPTION	PLATINUM GRAMS PER TONNE	SOURCE
Maple Leaf	From carlode of picked copper ore shipped from upper workings.	15.4	B.C. Annual Reports; 1920, page 153 1929, page C254
	Samples taken from almost pure mass of chalcopyrite occurring in the pyroxenite near the margin with the augite syenite hanging wall side of the lode.	13.0	Thomlinson, 1920, page 162
	Sample taken from main open cut at upper workings.	5.14, 5.83	Thomlinson, 1920, page 165
Lucky Jack	Dump of pyroxenite at mouth of short drift near west end-line. Selected pieces of dark-coloured close-grained rock with chalcopyrite and small crystals of whitish metallic mineral.	2.74	Thomlinson, 1920, page 163
	From small shaft 200 feet easterly from above sample location. Chalcopyrite and pyrite within lens of dark, close-grained rock.	1.37	Thomlinson, 1920, page 163
	From opencut near east end-line, medium grained pyroxenite with chalcopyrite, pyrite and stained by copper carbonates.	2.06	Thomlinson, 1920, page 163
Mountain Lion	Oxidized pyroxenite and iron sulphide from small shaft and open cut on ridge 1000 feet west of shaft on Gloucester claim.	3.09	Thomlinson, 1920, page 163
	Samples from small shaft and large open cut. Decomposed pyroxenite with reddish-brown iron oxides.	0.69	

TABLE CONTINUED

CLAIM	DESCRIPTION	PLATINUM GRAMS PER TONNE	SOURCE
Golden Age	Oxidized pyroxenite with fine particles of chalcopyrite and stains of copper carbonates from small shaft near southwest of claim.	2.06	Thomlinson, 1920, page 164
Averill Group	From dump at lower drift near the cabin. Coarse grained pyroxenite associated with fine grained syenite, with chalcopyrite and bornite.	3.06	Thomlinson, 1920, page 164
	Dump at upper shaft. As above.	3.06	Thomlinson, 1920, page 164
Buffalo	Dump at old shaft below and near trail at southwest end of claim. Pyroxenite, with small masses of chalcopyrite, some pyrite and copper carbonate.	6.51	Thomlinson, 1920, page 165
	From open cuts in dry ravine below trail near southeast corner, selected pieces with pyrite and chalcopyrite.	2.74	Thomlinson, 1920, page 165
Ottawa	Pyroxenite with magnetite, pyrite and chalcopyrite from large open-cuts in dry ravine below trail near southeast corner.	2.06	Thomlinson, 1920, page 165
Columbia	Pyroxenite with pyrite and chalcopyrite from dumps of two drifts on side-hill north of cabin on north bank of Franklin Creek.	1.37	Thomlinson, 1920, page 165

TABLE CONTINUED

CLAIM	DESCRIPTION	PLATINUM		SOURCE
		GRAMS	PER TONNE	
Davenport	Highly mineralized pyroxenite with minute specks of bornite and masses of chalcopyrite from east end of 75 foot long tunnel.	3.77		B.C. Annual Report 1921 , page G180
	Panned sample near mouth of Franklin Creek.	1.03		O'Neill and Gunning 1934, page 99
	Panned sample from McDonald Creek.	3.09		O'Neill and Gunning 1934, page 99
	Panned sample from west branch of north fork of Kettle River.	2.06		O'Neill and Gunning 1934, page 99

SELECTED BIBLIOGRAPHY

- Allen, Guy (1982): Prospecting report on the Glouster and G. H. Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 10953.
- Chilcott, R. P. and Lisle, T. E. **(1964)**: Report on the Franklin Mining Camp, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 637.
- Church, B. N. (1985): Geology of the Mount Attwood-Phoenix area, Greenwood (82E/2). British Columbia Ministry of Energy, Mines and Petroleum Resources, Geological Fieldwork, 1984, Paper 1985-1, pages 17-21.
- Coryell, John (1900): Franklin Camp, Minister of Mines, B.C., Annual Report 1901, page 1066.
- Cox, P. M. and Friesen, P. J. **(1972)**: Geochemical Soil Survey in the Franklin Mining Camp, Assessment Report on the Little Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 10604.
- Drysdale, Charles W. (1915): Geology of the Franklin Mining Camp, British Columbia, Geological Survey of Canada, Memoir 56, 246 pages.
- Freeland, P. B. (1920): Franklin Camp, Minister of Mines, B.C., Annual Report 1919, pages K206-K207.
- (1921). Franklin Camp, Minister of Mines, B.C., Annual Report 1920, pages 153-154.
- (1922). Franklin Camp, Minister of Mines, B.C., Annual Report 1921, page G181.
- (1930). Franklin Camp, Minister of Mines, B.C., Annual Report 1929, pages C253-254.
- **(1933)**. Franklin Camp, Minister of Mines, B.C., Annual Report 1932, page A121.
- Galloway, J. D. **(1914)**: Franklin and Glouster Camps, Minister of Mines, B.C., Annual Report **1913**, pages K167-169.
- Keating, John (1983): Assessment Report on the Geological and Geochemical Survey on the Pl 1-3 Mineral Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12254.

SELECTED BIBLIOGRAPHY

- Larson, A. G. **(1915)**: Mineral and Other Resources of the North Fork of the Kettle River, Minister of Mines, B.C. Annual Report 1914, page K340, 354.
- Lisle, T. E. (1976): Assessment Report on the Glouster Claim, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 6228.
-(1980). Preliminary Geological Report on the Spring Group, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Reports 8126, 8149.
- Lisle, T. E. and Seraphim, R. H. **(1983)**: Drilling Report on the Spring Group, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report **9115**.
- Mertie, John B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, page 29.
- O'Neill, J. J. and Gunning H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada - Economic Geological Series, Number 13, pages 79,99,104,106.
- "Sourdough" **(1955)**: The Discovery Post, Western Miner and Oil Review, Volume 28, Number 9, page 44.
- Stephen J. C. and Shearer, J. T. **(1980)**: Geological and Geochemical Report on White Bear Group, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 7918.
- Thomlinson, Wm. (1920): Platinum-Mineral Investigations, Munitions Resource Commission, Canada, Final Report, pages 161-166.
- Walls, J. R. (1984): Geological and Geochemical Report on the White Bear Crown Grant and Tenderloin (A) Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 12508.

ii) Alkalic Hosted Occurrences

SAPPHO

LOCATION

The Sappho property is located 8 kilometers south of Greenwood and 5 kilometres east of Midway on the International Boundary. Access is from Highway 3, about 3 kilometres east of Midway to the Norwegian Creek road which is followed southeasterly for about 3 kilometres until it crosses Norwegian Creek. From here, a 0.2 kilometre hike along an old wagon trail leads into the old workings.

REGIONAL AND LOCAL GEOLOGY

The Sappho Crown Grant is situated on the eastern fault boundary of the north-northeast-trending Tertiary Toroda Creek graben, within the Omineca belt of British Columbia. Interpretation of the geology of the area has been hampered by extensive Eocene extensional tectonics. Late Paleozoic Knob Hill greenstones occur as roof pendants through a Triassic microdiorite (Church, 1985). Tertiary Coryell alkalic intrusives (shonkinites, feldspar porphyry dykes) occur within the central and northeastern portions of the Sappho claim. A serpentine schist body of unknown age occurs several hundred metres to the north of the claim and sheared serpentine fragments have been found in the dumps of some old workings in the area of the claims (Church, 1983).

DISTRIBUTION AND OCCURRENCE OF P.G.E.

Platinum values of 1.0 gram per tonne were obtained from chalcopyrite-rich ore within a pyroxenite, in 1927 (O'Neill and Gunning, 1934). Subsequent rock sampling and trenching of the copper mineralization in the pyroxenite, done in 1975, 1978 and **1981**, confirmed the presence of platinum; by assays of up to 1.75 grams per tonne platinum were found (Gilmour, **1981**).

The host to the copper-platinum mineralization is variably identified as Coryell shonkinite, a pyroxene-rich marginal phase of an alkalic intrusive, by Church (1985) and a serpentinized and foliated syenodiorite to gabbro by Gilmour (**1981**). Copper, silver and platinum mineralization occur as blebs and disseminations in shear zones or are associated with a skarn-like zone of strong biotite, garnet and chlorite alteration (Gilmour, **1981**) near the margin between the mafic rocks and the intrusive microdiorite stock. The highest platinum assay (1.75 grams per tonne - Gilmour, 1981) was obtained from a 50 cm. long pod of massive chalcopyrite within sheared and extensively (biotite, garnet and calcite) altered upper Paleozoic greenstones (Gilmour, 1981); these would correspond to the Tertiary biotite shonkinite of Church (1983). According to Fyles and Keating (1984), the Sappho showings are situated at or close to the point where a north-northeast trending zone of Eocene normal faulting associated with the regional Toroda Creek graben crosses the altered mafic volcanic rocks. Evidently, no agreement exists about either the nature of the host rocks or their ages.

SELECTED BIBLIOGRAPHY

- B.C. Ministry of Energy, Mines and Petroleum Resources (1975): Sappho (Cabin), Exploration in British Columbia, 1975, pages E12-E13.
- Brock, R. N. (1903): Preliminary Report on the Boundary Creek District, British Columbia. Geological Survey of Canada, Summary Report, 1902, Volume XV, pages 92A-138A.
- Church, B. N. (1985): Geology of the Mount Attwood-Phoenix Area, B.C. Ministry of Energy, Mines and Petroleum Resources, Preliminary Map 59.
- Church, B. N. and Robertson S. (1983): Geology and Magnetometer Survey of the Sappho Gold-Silver-Platinum-Copper Prospect, B.C. Ministry of Energy, Mines and Petroleum Resources, Geological Fieldwork, 1982, Paper 1893-1, pages 27-31.
- Freeland, P. B. (1928): Sappho Greenwood Section of the Southern Mineral Survey District (Number 4), Minister of Mines, B.C., Annual Report 1927, pages
- Fyles, J. T. and Keating John (1984): Geological and Geochemical Survey on the Sappho 83 Group of Claims, B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment report 12924.
- Gilmour, W. R. (1981): Assessment Report on the Sappho Property for Kettle River Resources, Ltd., B.C. Ministry of Energy, Mines and Petroleum Resources, Assessment Report 9364.
- Little, H. W. (1983): Geology of the Greenwood Map Area, British Columbia, Geological Survey of Canada, Paper 79-29, 37 pages.
- McKechnie N. D. and Olson, P. E. (1964): Cabin Group, Minister of Mines, B.C., Annual Report 1964, page 110.
- Mertie, John B. (1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, page 29.
- O'Neill, J.J. (1919): The Platinum Situation in Canada, 1918, Geological Survey of Canada, Summary Report, 1918, Part G, page 8G.
- O'Neill, J. J. and Gunning, H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series, Number 13, pages 103-104.

MISCELLANEOUS (CONTINUED)

iii) Platinum-Group Element Occurrences in Skarns, Porphyry Copper and Stratabound Lead-Zinc Deposits

Smelters at the Nickel Plate and Copper Mountain-Ingerbelle Mines recorded trace platinum. The Sullivan Mine, is presumed to be the source of palladium and platinum recovered as by-products of refining at the Trail Smelter during the period **1928-1930**.

SULLIVAN MINE

The Sullivan Mine is a stratabound Pb-Zn-Ag deposit located near Kimberley, British Columbia. The ore is confined and conformable to Aldridge Formation metasedimentary rocks of the mid-Proterozoic Purcell Supergroup. According to O'Neill and Gunning (1934, Page 107) palladium and platinum, worth \$25,365.00 and \$3,177.00 (**1928-1930** prices) respectively, were recovered "as a by-product of refining at the Trail plant of the Consolidated Mining and Smelting Company of Canada".

No platinum or palladium has been reported, however, in recent and routine analyses of lead or zinc concentrates and of pyrrhotite which have failed to indicate platinum or palladium above detection limits of 50 ppb (Hamilton, J., 1986, pers. comm.).

O'Neill (1919), remarks that palladium was noted at the Trail Smelter of the B.C. Consolidated property, in association with copper-gold ore.

iii) Platinum-Group Element Occurrences in Skarns, Porphyry Copper, and
and Stratabound Lead-Zinc Deposits (Continued)

NICKEL PLATE MINE

At the Nickel Plate Mine, auriferous arsenopyrite and bismuth telluride ore occurs at the outer margins of a pyroxene skarn zone between limy clastic rocks and calcareous sediments of the Nicola group, and diorite porphyry sills of the Hedley Intrusions. Assays of 0.5 percent platinum, occurring as sperrylite ($PtAs_2$), were reported from a residue on the plates at the stamp mill at the mine (Camsell, 1910, page 137).

COPPER MOUNTAIN

Chalcopyrite and bornite as disseminated blebs and fracture fillings mineralization occur within highly fractured Upper Triassic Nicola Group bedded tuffs and andesites which have been intruded by the alkalic Copper Mountain intrusions. According to O'Neill and Gunning (1934, Page 103); "Jules Catharinet (1905) mentions reports of platinum assays from Copper Mountain and claims to have identified sperrylite in a pegmatite on the Copper Cliff claim. Minute crystals and grains are said to have been observed in biotite in a thin section and in bornite and chalcopyrite microscopically".

Recent analyses of sulphide concentrates obtained from the Copper Mountain - Ingerbelle Mine substantiate the presences of palladium and platinum in the ore (Mutschler, et al., in press).

SELECTED BIBLIOGRAPHY

Camsell, Charles (1910): Geology and Ore Deposits of Hedley Mining District, Geological Survey of Canada, Memoir 2, Page 137.

Catharinet, Jules (1905): Copper Mountain, British Columbia, Engineering and Mining Journal, Volume 79, Pages 125-127.

Mertie, J. B. Jr. **(1969)**: Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, Pages 28-29.

Mutschler, F. E., Griffen, M. E., Stevens, Scott D and Shannon, S. S. (In press): Previous Metal Deposits Related to Alkaline Rocks in the North American Cordillera - Transactions of the Institute of Mining and Metallurgy - South Africa.

O'Neill, J. J. (1919): The Platinum Situation Canada, 1918, Geological Survey of Canada, Summary Report, 1918, Part G, pages IG-19G.

O'Neill J. J. and Gunning, H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, Pages 103, 107.

TABLE 3. PLATINUM GROUP ELEMENT PLACERS IN BRITISH COLUMBIA

River	Tributary	Description	Assay	Source
Similkameen		On Pedersons Flat, 4 miles above Princeton platinum occurs in gravels	\$0.60/yd (\$115/oz)	MMAR 1924, p.B176
		Combined black sand from two pans of sandstone bedrock on Tulameen River, 2 miles above Princeton on south bank of river	200.9 g/t	MacKenzie, 1920, p.150
		One pan of river gravel 2 miles above Princeton on south bank of river	239.3 g/t	MacKenzie, 1920, p. 150
Tulameen (92H/7)		Platinum was recovered along the Tulameen from its mouth near Princeton, to the mouth of Granite Creek (approximately 12 miles) and between the mouths of Olivine and Champion Creeks. Champion Creek, Brakeburn Creek, Newton Creek which all have their sources in the ultramafic complexes, bear platinum and iridium in their gravels. Other platinum-bearing tributaries to the Tulameen are Cedar Creek, Slate (Olivine), Bear, Pine, Eagle, Otter and Boulder		
Tulameen	Bear Creek (92H/5)	Combined black sand and gravel from two pans of medium-sized gravel from Tulameen River at mouth of Bear Creek	1733.5 g/t	MacKenzie, 1920, p.150
Tulameen	Slate (Olivine) Creek (92H/10)	Combined black sand from five pans of medium-sized gravel from Tulameen River at mouth of Slate (Olivine) Creek	600.3 g/t	MacKenzie, 1920, p.150

River	Tributary	Description	Assay	Source
Tulameen	Slate (Olivine) Creek (92H/7)	Combined black sand from three pans of medium-sized gravel from Tulameen at mouth of Slate (Olivine) Creek	1316.6 g/t	Mun. Res. Com., 1920, p.150
		Washings of 90 cubic yards of gravel on Tulameen River opposite Slate Creek	8.46 g/t	Mun. Res. Com., 1920, p.152
Tulameen	Granite Creek (92H/7)	Analyses of 17.89 grams of native platinum from Granite Creek (a) non-magnetic and (b) magnetic separations. Granite Creek has been mined from the Tulameen River to Newton Creek where the gold to platinum ratio was 4:1	Pt (a) 68.19% (b) 78.43% Pd (a) 0.29% (b) 0.09% Rh (a) 3.10% (b) 1.70% Ir (a) 1.21% (b) 1.40% OsIr (a) 14.62% (b) 3.77%	GSCAR 1899, p.5-9
Tulameen	Siwash Creek (92H/11)	Platinum is reported to occur with gold in black sands		MMAR 1911, p.K183
Coquihalla	Sowoqua (92H/6)	Drains Coquihalla River 1.5 miles below Jessica. "Platinum values are claimed by owners" in areas underlain by serpentines of the Coquihalla Belt		MMAR 1922, p.N143
Okanagan	Shuttleworth (82E/5)	Evidence of presence of platinum but not in commercial quantities		Mun. Res. Com., 1920, p.168-169; Camsell, 1919, p.28B
Kettle	Rock Creek (82E/5)	"Platinum Occurs in exceedingly minute to moderate sized, coarse irregularly shaped grains, the largest of which measures 4 mm in diameter; a little chromite was detected in one of the pellets of platinum"		GSCAR 1892-93, p.14

River	Tributary	Description	Assay	Source
Kettle	Rock Creek (82E/5)	Sample of heavy black sand taken from riffle of sluice boxes at Camp McKinney	44% native platinum	GSCAR 1892-93, p.14
Kettle	Franklin (82E/9)	Analyses of black sands and pannings taken near mouth of Franklin Creek	1.03 g/t	Mun. Res. Com., 1920, p.167
		Black sand pannings from McDonald Creek, branch of Franklin Creek	3.09 g/t	Mun. Res. Com., 1920, p. 167
Lardeau (82K/2)		"Pannings from bar below small canyon at bridge on road to Truine Mine"	1.37 g/t	Mun. Res. 1920, p.171
		"Pannings from Ninemile point, near old cabin above falls"	0.34 g/t 1920, p.171	Mun. Res.Com.,
		"Pannings from bar at Eightmile point just above bridge on road to Silver Cup Mine"	5.49 g/t 1920, p.171	Mun. Res. Com.
		"Pannings from bars at mouth of Finkle Creek which enters river from north at Sevenmile point, near Livingstone's Camp"	4.11 g/t	Mun. Res. Com. 1920, p. 171
		"Pannings from bars 300 to 500 yds above old power house of Silver Cup mill, about 6 miles from Ferguson"	4.11 g/t	Mun. Res. Com. 1920, p.171.
		"Discarded 'black sand' found lying near cabin at placer workings one half mile from Ferguson, and mouth of North Fork"	0.69 g/t	Mun. Res. Com. 1920, p. 171
		"Pannings from North fork of Lardeau River. Mainly from bars below small canyon about 2 1/2 miles from Ferguson"	3.77 g/t	Mun. Res. Com, 1920, p. 171

River	Tributary	Description	Assay	Source
Lardeau (82K/2)		"Pannings from North fork of Lardeau River. Small bars in canyon below wagon-road bridge, about 1/2 mile from Ferguson"	1.03 g/t	Mun. Res. Com., 1920, p. 171
		"Lardeau River, Bench placer on main river near Ferguson townsite"	0.34 g/t	Mun. Res. Com., 1920, p. 171
		"Black sand said to be from small bars in centre of main canyon of the Lardeau River, halfway between Trout Lake and Ferguson"	7.54 g/t	Mun. Res. Com., 1920, p.172
		"Pannings from bars near mouth of canyon, Lardeau River about one half mile north of town of Trout Lake-"	5.83 g/t	Mun. Res. Com., 1920, p. 172
Lardeau	Five-mile Creek (82K/2)	Material from panning	3.09 g/t	Mun. Res. Com., 1920, p.172
Lardeau	Canon Creek, near Gerrard	Material from panning	0.69 g/t	Mun. Res. Com., 1920, p.172
		Pannings from Lardeau River near mouth of Cascade Creek	1.37 g/t	Mun. Res. Com., 1920, p.173
		"Pannings from bars near old dredge at Gold Hill, Lardeau River; also from point on South side of channel for about 1 mile upstream from dredge"	1.37 g/t	Mun. Res. Com., 1920, p.173
		"Pannings from bars near old dredge and along south side of channel, for about 1/2 mile downstream from dredge"	0.69 g/t	Mun. Res. Com., 1920, p.173
Columbia	Toby Creek (82K/8, 9)	Platinum reported mixed with coarse gold		O'Neill, 1919, p.10G

River	Tributary	Description	Assay	Source
Columbia	Isaac Creek	"Pannings from bars along south bank of Isaac Creek, for 3/4 of a mile upstream from a Point 1/2 mile above falls"	1.37 g/t	Mun. Res. Com., 1920, p.175
		"Bars along Isaac Creek for 3/4 of a mile up stream from a point 1 1/4 miles above falls"	0.69 g/t	Mun. Res. Com., 1920 p . 175
		"Bars along Isaac Creek for one-half mile up stream from a point 2 miles above falls"	0 g/t	Mun. Res. Com., 1920, p.175
Thompson	Clearwater (92P/9)	Near mouth of Clearwater, traces of platinum recorded. Samples of black sand sent to Baker and Co. - Newark, New Jersey (platinum refiners)	4251 g/t	MMAR 1900, p.894; 1901, p.1082
Thompson	Tranquille (92I/10)	Pannings from points along Tranquille River, near schoolhouse, up stream to Sharps cabin, a distance of 1/2 mile	0 g/t	Mun. Res. Com., 1920, p. 176
		Pannings from points along Tranquille River from Sharps cabin to main canyon, a distance of 1 mile	0.69 g/t	Mun. Res. Com., 1920, p. 176
Thompson	Criss Creek (tributary of Deadman River) (92I/15)	Sample concentrated from the gravels in a pit at the junction of Criss Creek and Deadman River	4.11 g/t	O'Neill and Gunning, 1934, P. 88
Bonaparte	Scottie Creek (92I/14)	Pannings from bars and old workings between Hunter ranch and dam at Chinese workings	0.34 g/t	Mun. Res. Com., 1920, p.177
		Pannings from bars along creek from dam at old diggings to a point below canyon. Gold may contain iridium, osmium, ruthenium, rhodium,	0 g/t	Mun. Res. Com., 1920, p.177

River	Tributary	Description	Assay	Source
Bonaparte	Scottie Creek (921/14)	Pannings from cabin on Gagne's claims to old shaft on low bench near creek, a distance of 1/3 mile	4.80 g/t	Mun. Res. Com., 1920, p.178
		Pannings from bars along Scottie Creek near old shaft on Gagne's claims to mouth of Chrome Creek, a distance of about 2/3 mile	1.37 g/t	Mun. Res. Com., 1920, p. 178
Bonaparte	Chrome Creek (921/14)	Pannings from mouth upstream for about 1/4 mile	2.06 g/t	Mun. Res. Com., 1920, p.178
		Pannings from 1/4 mile from mouth to a point opposite the Chromite claims, a distance of about 1/2 mile	2.06 g/t	Mun. Res. Com., 1920, p.178
Fraser	(92H/6)	Platinum has been found in placers along the Fraser River from a few miles below the mouth of Ouesnel River to a few miles north of Hope		
	(92H/11)	Black sands near Yale - average of 6 samples of 5 assay tons each	13.7 g/t	O'Neill and Gunning, 1934, p.101
		Black sands opposite Saddle Rock Station, near Yale	4.80 g/t	O'Neill and Gunning, 1934, p.101
	(92H/14)	Boston Bar - a little platinum recovered in concentrates from dredge samples		O'Neill and Gunning, 1934, p.101
	(92H/14)	North Bend - platinum reported in black sands		O'Neill, 1919, p.107

River	Tributary	Description	Assay	Source
Fraser	(92I/12)	4-5 miles south of Lillooet on east side of river, platinum obtained from cleaning up concentrating tables of old dredge at Lillooet. "Platinum was in fair sized flakes and Dawson reports the occurrence of the- metal in very fine scales, with gold, about 10 miles below Lillooet"	17.1 g/t	GSCAR 1887-88, p.156; O'Neill and Gunning, 1934, p.101
	(92I/5)	Cobledick Dredge - at the Van Winkle bar near Lytton, platinum and iridium occur in black sands	5681.1 g/t	O'Neill and Gunning, 1934, p.101; B.C.O.F. 92I-4E
	(92I/12)	Fosters Bar - 23 miles north of Lytton platinum occurs with gold in black sands. Assay results were obtained from four 1-ton concentrates	174.2 g/t	B.C.O.F. 92I/NW-89
			199.9 g/t	
			141.3 g/t	
			291.4 g/t	
	(92N/1)	Glasgow - 18 miles southeast of Clinton; platinum occurs in gravel benches, terraces and beaches. Emission spectrography of a 1.6 oz of 'platinum' sample indicated 0.5 oz platinum, 1.0 oz palladium and 0.1 oz	Pt	Geiger, 1970
			21.6 g/t	
			Ir	
		Sample of black sand taken from Fraser River at mouth of Quesnel River	Pt 71% Os 3.1%	O'Neill and Gunning, 1934, P. 101
			82.3 g/t	
		Sample of black sand taken 2 miles above Quesnel River mouth		O'Neill and Gunning, 1934, p.101
		Black sand, 3 miles above Quesnel River mouth	Pt 8.6 g/t OsIr 109.7 g/t	O'Neill and Gunning, 1934, P. 101

River	Tributary	Description	Assay	Source
Fraser	(92N/1)	Black sand, 15 miles above Quesnel River mouth	133.7 g/t	O'Neill and Gunning, 1934, p.101
	Lillooet River (92G/16W)	Platinum, palladium and iridium occur in sands with gold and silver. (a) PML 811, (b) PML 813	(a) Pt 1.44 g/t Pd 1.06 g/t Ir 0.07 g/t (b) Pt 1.58 g/t Pd 0.69 g/t Ir 0.07 g/t	B.C.O.F. 92GNE013 B.C.O.F. 9213NE019
	Quesnel River (93A/14)	Head of Harvey Creek	0 g/t	O'Neill and Gunning, 1934, p.84
	(93A/14)	Upper Cunningham Creek	0 g/t	"
	(93A/7)	Eureka Creek	0 g/t	"
	(93A/11)	Keithly Creek Black sands, 82 miles above mouth	0 g/t 34.3 g/t	"
	(93B/16)	Black sands, 40 miles above mouth	96.0 g/t 4.8 g/t	"
	93B/16)	Black sands, 25 miles above mouth	13.7 g/t	"
	(93B/16)	Black sands, 13 miles above Quesnel	267.4 g/t	"
		Black sands, 30 miles above Quesnel	219.4 g/t	"
	(93A/12)	Black sands, 25 miles below forks	17.1 g/t 85.7 g/t	"
	(93G/1)	Three miles above Quesnel	8.6 g/t	"
	(93A/6)	Horsefly River - platinum reported in small quantities with gold	0 g/t	O'Neill and Gunning, 1934, P.83

River	Tributary	Description	Assay	Source
Fraser	Quesnel River (93A/5)	Bullion pit - platinum, osmiridium and palladium are present in the heavy concentrates that remain in the sluices after cleaning up and also in gravels as fine grains	0 g/t	O'Neill and Gunning, 1934, p.85; MMAR 1928, p.C201
	Cottonwood (93G/2)	Platinum and iridium Occur in gravels in river bars, benches of gravel and deposits in rock benches where the river crosses ultrabasic rocks	0 g/t	MMAR 1936, p.C19
Quesnel	Morehead Creek (93A/12)	On the south bank of tile Quesnel River at mouth of Morehead CreeK, crude platinum was reported by Water tight Dipper Dredge and Mining Co. Ltd. of Calgary		MMAR 1913, p.62
Quesnel	Twenty Mile Creek (93A/12)	Platinum, palladium and osmiridium are found in minute metallic grains and enclosed in small fragments and nuggets of magnetite and chromite. Analysis of sample of a pan of concentrates taken from tile sluices after cleanup	Pt 2194.3 g/t Pd 2208.0 g/t Os 1440.0 g/t	Uglow, 1920, p.215
Fraser	Government Creek (tributary of Hixon Creel.-:') (93G/9)	Platinum in black sands was recorded in creek beds and shallow covered benches	96.0 g/t	O'Neill and Gunning, 1934, p.83; MMAR 1919, p.K128
Parsnip	McLeod (93J/14)	Iridium, and platinum occur with gold in shallow gravels on rock benches and also in the cracks and crevices of the rock under tile gravel. Pyroxene intrusions nearby suggest a source for the platinum group elements		MMAR 1932, p.A88

River	Tributary	Description	Assay	Source
Parsnip	McLeod (93J/14)	C. Nelson's claim, located at the lower end of a low-lying rock bench, 1/2 mile south of the mouth of the McDougall River	8.6 g/t	MMAR 1932, p.A88
		Sample of black sands from which gold had been removed by amalgamation - from Cluckhoo claim, 1 1/2 miles below mouth of McDougall River on McLeod River on the right bank	\$4500/t Pt \$780/t Ir (1931 price)	MMAR 1934, p.A8
Prsnip	(93O/13E)	Platinum was reported to occur with gold in sands at Bill Cuts Bar which is now flooded by Williston Lake		MMAR 1894
Nat i on	Rainbow Creek (93O/4W)	Platinum and iridium or-cur in black sands i n the canyon. Ratio of platinum to gold averages 1:2	\$96/t Pt, Ir (1931 price)	O'Neill and Gunning, 1934, p.82
Stuart	Dog Creek (93K/8)	Platinum occurs with gold in gravels overlying false bedrock on extensive bench near creek	\$0.01/yd3	MMAR 1931,
Ingenika Creek	McConnell (94D/16)	Trace platinum and palladium occur with gold in terraced gravel floor along McConnell Creek and within 1 to 8-foot-deep pockets of sand, gravels and boulders which extend from the mouth of McConnell Creek 10 miles down the Ingenika River		O'Neill and Gunning, 1934 p. 83
		Highest platinum value from north end of a pit, 3 feet deep on Lase No. 457	\$0.8/yd3 (1932 price)	O'Neill and Gunning, 1934 p.38
		Sample of black sand after gold removed by amalgamation	\$45.2 Pt/t (1932 price)	O'Neill and Gunning, 1934 p.83

River	Tributary	Description	Assay	Source
Manson Creek	(93N/9)	Platinum reported in placers along creek		O'Neill, 1919, p.10G
Germanson	Ah Hoo (93N/10)	At Ah Hoo Creek a belt of nickeliferous (0.18%) serpentine is cut by the creek. Below this point placer deposits contain small amounts of platinum		MMAR 1936, p.C6
Peace	(94B/4)	Sample of black sand concentrate (971 lbs) obtained from top 8 inches of bar gravels on Pete Toy Bar, 3 miles above Finlay Forks. This area is now flooded by Williston Lake	33.3 g/t	O'Neill and Gunning, 1934, p.82
Peace	(94B/2)	Placer platinum has been reported to occur with gold at the upper end of Rocky Mountain Canyon avoid the junctions of the Finlay and Parsnip Rivers with the Peace River, at Branhams Flats. This area is flooded by the Portage Mountain Dam		Mun. Res. Com., 1920, p.156-160
	94B/3)	"Platinum has been found in small amounts in the bars and benches of the upper Peace River, chiefly between Hudson's Hope and the Ne Parle pas rapids"	\$0.80/yd (1918 price)	Uglow, 1920, p.218; O'Neill and Gunning, 1934, p.83
Peace	McLean Creek (93P/4) MacDonald Creek	Platinum occurrence is by report only		O'Neill, 1919, p.10G
Dease	Thibert Creek (104J/15)	Platinum and osmium occur in black sands in gravels on Thibert Creek below mouth of Berry Creek. The richest gravels are only a few feet thick and rest on bedrock. Serpentinities occur along Thibert Fault		O'Neill and Gunning, 1934, p.76-77; MMAR 1905, p.77--78

River	Tributary	Description	Assay	Source
Dease	Thibert Creek (104J/15)	Analyses of black sand remaining in the sluice boxes and undercurrents Assays of concentrates from top gravel obtained from (a) B.C. Government Assay office and (b) San Francisco	44.1% Pt 11.9% Os, Ir (a) Pt 2057.1 g/t (b) Pt 514.3 g/t	MMAR 1902, p. 44
Atlin Lake	Ruby Creek (104N/11)	Crystals and grains of iridosmine were obtained from black sands on Ruby Creek		O'Neill and Gunning, 1934, p.80-81
Atlin Lake	Boulder Creek (104N/11)	Assay taken from black sand concentrate	2.7 g/t	B.C.O.F. 104N/5, 6,11,12
Graham Island (103J/4)		Platinum has been reported in the beach sands on the northeast coast of Graham Island from Toe Hill to Rose Point. However this was not substantiated by Munitions Resource Commission		O'Neill and Gunning, 1934, p.81
Vancouver Island	Wreck Bay (Florenca Bay) (92C/13)	Platinum was detected in two samples of auriferous black beach sands	1.7 g/t 1.0 g/t	O'Neill and Gunning, 1934, p.100
Jervis Inlet	Potato Creek (92J/4)	Platinum occurs in gravel bars, pits and benches in the main channel of Potato Creek	0.69 g/t 5.8 g/t	Mun. Res. Com., 1920, p.182

GSCAR - Geological Survey of Canada Annual Report
MMAR Minister of Mines, B.C. Annual Report
Mun. Res. Com. - Munition Resources Commission
B.C.O.F. - British Columbia Open File
g/t - grams per tonne

ACKNOWLEDGMENTS

Open File 1986-7 was funded by, and a contribution to the Canada-British Columbia Mineral Agreement. Preparation and compilation of this publication was supervised by Mr. A. F. Wilcox and Mr. J. G. McArthur, Manager of Resource Data and Analysis Section of the British Columbia Geological Survey Branch.

Valuable assistance by staff geologists with the British Columbia Geological Survey Branch, especially Mr. T. E. Kalnins, Mr. D. J. Alldrick, Dr. B. N. Church, Dr. G. E. Ray and Dr. W. J. McMillan, contract geologists Mr. B. Grant, Mr. G. Dawson and Dr. J. Fyles of Kettle River Resources was invaluable.

Aid in gathering vital resource material was given by Ms. S. Ferris and Ms L. Peacock, librarians at the library of the British Columbia Ministry of Energy, Mines and Petroleum Resources; their help is acknowledged with respect and thanks.

Dr. W. J. McMillan provided key support and manuscript revision. Technical assistance by Mrs. R. Moir and Mr. J. Armitage and manuscript preparation by Ms. B. Gordon and Mrs. D. Bulinckx is acknowledged with appreciation.

BIBLIOGRAPHY

- Agiorgitis, G. and Wolfe, R. (1978): Aspects of Osmium, Ruthenium and Iridium. Contents in some Greek Chromitites, Chemical Geology, Volume 23, pages 267-272.
- Aho, A. E. (1956): Geology and Genesis of Ultrabasic Nickel-Copper-Pyrrhotite Deposits at the Pacific Nickel Property British Columbia: Economic Geology, Volume 51, pages 444-481.
- Allen C. C. (1960): The Platinum Metals: Canada Department Mines and Technical Surveys, Minerals Resources Division, Mineral Report 3, 68 pages.
- Angeli, N. and Choudhuri: Ultramafic Complexes and Associated Mineral Deposits in the Pre-Cambrian of Eastern Minas Gerais, Brazil, Economic Geology, Volume 20, Number 4, page 809.
- Anslett, T. F., Bleiwos, D. I., and Sheng-Fogg, C. (1982): Platinum Availability - Market Economy Countries; A Minerals Availability Systems Appraisal. United States Department of the Interior, Bureau of Mines, Information Circular 8897.
- Baglin, E. G. (1985): Recovery of Platinum, Palladium, and Gold from Stillwater Complex Flotation Concentrate by a Roasting-Leaching Procedure, United States Geological Survey, Report of Investigations 8970, 12 pages.
- Bancroft, H. (1910): Platinum in Southeastern Nevada (Clark County), United States Geological Survey, Bulletin 430, pages 192-199.
- Barker, J.C. (1982): Critical and Strategic Minerals in Alaska: Cobalt, the Platinum group metals, and chromite, United States Geological Survey, Information Circular 8869, 8 pages.
- Barker, J.C., Still, J.C., Mowalt, T. C. and Mulligan, J. J. (1981): Critical and Strategic Minerals in Alaska, United States Bureau of Mines, Information Circular 8869.
- Barker, J.C. Thomas, D. L. and Hawkins, D. B. (1985): Analyses of Sampling Variance From Curtian Platinum and Palladium Deposits in Alaska, - United States Geological Survey, Report of Investigations, 8948, 26 pages.

BIBLIOGRAPHY (CONTINUED)

- Beath, C. B., Cousins, C.A. and Westwood, R. J. (1961): The Exploration of the Platiniferous Ores of the Bushveld Igneous Complex with Particular Reference to the Rustenburg Platinum Mines, Commonwealth Mining and Metallurgical Congress, 7th, South Africa, Transaction, Volume 1, pages 217-243.
- Belincourt, L. E., Hummel, and H. H. and Skinner, B. J. (1982): Phases and Phase Relations of the Platinum-Group Elements, in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L.J. Cabri, C.I.M. Special Volume 23, Chapter 3, pages 19-46.
- Betekhtin, A. G. (1930): Character, Classification, Occurrence and Genesis of Platinum Mineral Deposits of Russia, Academy of Science USSR, 148 pages.
- Bonatti, E. (1978): The Origin of Metal Deposits in the Oceanic Lithosphere, Scientific American, Volume 238, Number 2, pages 54-61.
- Boudreau, Alan E. (1985): The Composition and Role of High-Temperature Hydrothermal Fluids in the Genesis of PGE Deposits in Layered Intrusions (abstract).
- Boyle, R. W. (1982): Gold, Silver and Platinum Metal Deposits in The Canadian Cordillera - Their geological and geochemical setting in Precious Metals in the Northern Cordillera, ed. A. A. Levinson, The Association of Exploration Geochemists, pages 1-18.
- Brownlee, Donald E., Bates, B. A. and Wheelock, M. M. (1984): Platinum Group Nuggets in Deep Sea Sediments, (abstract) in Abstracts of Papers Presented at the 47th Annual Meeting of Meteoritical Society, Meteoritics, Volume 19, Number 4, pages 199-200.
- Buchanan, D. L. and Jones, M. J. (Ed) (1984): Sulphide Deposits in Mafic and Ultramafic Rocks. Proceedings of I.G.C.P. Projects 161 and 91, Third Nickel Sulphide Field Conference, Perth, Western Australia, 23-25, May, 1982, The Institute of Mining and Metallurgy, London, England 164 pages.
- Buddhas, J. D. (1951): Minerals of the Platinum Group: American Mineralogist, Volume 19, pages 350-352.
- Bureau of mines Minerals Yearbook (1984): Platinum Group Metals, United States Department of the interior, 10 pages.

BIBLIOGRAPHY (CONTINUED)

- Cabri L. J. (1972): The Mineralogy of the Platinum-Group Elements, Mining Science and Engineering, Volume 4, Number 3.
-(1976): Glossary of Platinum-Group Minerals, Economic Geology, Volume 71, Number 7, pages 1476-1480.
-(1981 a): Nature and Distribution of the Platinum-Group Element Deposits, Episodes, 1981, Number 2, pages 31-35.
-(1982): Classification of Platinum-Group Element Deposits with Reference to the Canadian Cordillera in Precious Metals in the Northern Cordillera, Ed. Levinson, The Association of Exploration Geochemists, pages 21-31.
-(1982): The Platinum-Group Minerals in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M., Special Volume 23, Chapter 7, pages 83-150.
-(1982): Relationship of Mineralogy to the Recovery of PGE from Ores in Platinum Group Elements; Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M., Special Volume 23, Chapter II, pages 234-250.
-(1982): Unnamed Platinum-Group Minerals. in Platinum-Group Elements; Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 9, pages 175-196.
- Cabri, L. J. and Harris, D. C. (1975): Zoning in Is-Ir-Alloys and the Relation of the Geological and Tectonic Environment of the Source Rocks to the Bulk Pt:Pt+Ir+Os Ratio for Placers, Canadian mineralogist, Volume 13, pages 266-274.
- Cabri, L. J. and LaFlamme, J. H. G. (1974): Rhodium, Platinum and Gold Alloys from the Stillwater Complex, Canadian Mineralogist, Volume 12, pages 399-403.
-(1976): The Mineralogy of the Platinum-Group Elements from some Copper-Nickel Deposits of the Sudbury Area, Ontario, Economic Geology, Volume 71, Number 7, pages 1159-1195.
-(1982): Analyses of Minerals Containing Platinum-Group Elements. in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 8, pages 151-174.

BIBLIOGRAPHY (CONTINUED)

- Cabri, L. J. and LaFlamme J. H. G. (1982): Sample Preparation techniques for PGE-Bearing Materials. in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 5, pages 65-70.
- Cabri, L. J., LaFlamme, J. H. G., Stewart, J. M., Rowland, J. F. and Chen, T. T. (1975): New Data on some Palladium Arsenid and Antimonides, Canadian Mineralogist, Volume 13, pages 321-335.
- Cabri, Louis J, Owens, Dalton R. and Laflamme, J. H. Gilles: Tulameenite, A New Platinum-Iron-Copper Mineral from Placers in the Tulameen River Area, British Columbia, Canadian Mineralogis., Volume 12, pages 21-25 (1973).
- Cabri, L. J. and Trail, R. J. (1966): New Palladium Minerals from Norilisk, Western Siberia, Canadian Mineralogist, Volume 8, pages 541-550.
- California Mining Journal (1985): Prospecting for Platinum metals, California Mining Journal, Volume 54, Number 6, pages 36-50.
- Campbell, Ian H., Barnes, Stephen, J. (1984): A Model for the Geochemistry of PGE in Magmatic Sulphide Deposits in Ore Deposits and Related Petrology of Mafic and Ultramafic Suites' Canadian Mineralogist, Volume 22, Pages 151-160.
- Carlson, C. A. (1985): Analyses for Platinum-Group Elements in Samples from Podiform Chromite Deposits, California and Oregon, United States Geological Survey, Open File 85-442, 14 Pages.
- Carlson, R. R., Carlson C. A. and Grimes, D. J. (1982): Geochemical Analysis for Platinum Group Elements in Rock Samples from the Kalmiopsis Wilderness, Southwest Oregon, United States Geological Survey, Open File 85-442, 14 pages.
- Chapin, Theodore (1919): Platinum-Bearing Auriferous Gravels of Chistochina River, Alaska, United States Geological Survey, Bulletin 692, pages 137-141.
- Christian, Jeffrey M. (1986): Platinum Demand Strengths, Supply Tightens, Prices Rise, Engineering and Mining Journal, March 1986, pages 43-45.

BIBLIOGRAPHY (CONTINUED)

- Chyi, L. L. and Crocket, J. H. (1976): Partition of Platinum, Palladium, Iridium and Gold Among Co-Existing Minerals from the Deep Ore Zone, Strathcona Mine, Sudbury, Ontario, Economic Geology, Volume 71, Number 7, Pages 1196-1205.
- Clark, A. L. and Greenwood, W. R. (1972): Geochemistry and Distribution of Platinum-Group Metals in Mafic to Ultramafic Complexes of Southern and Southeastern Alaska, United States Geological Survey, Professional Paper 800-C, Pages 157-160.
- Coles, B. R. (1964): The Magnetic Properties of Platinum metals and Alloys, A Survey of Recent Research: Platinum Metals Review, Volume 8, Number 1, Pages 9-11.
- Conn, H. K. (1979): The Johns - Manville Platinum-Palladium Prospect, Stillwater Complex, Montana, USA, Canadian Mineralogist, Volume 17, Pages 463-468.
- Cooley, E. F., Curry, K. J. and Carlson R. R. (1976): Analyses for the Platinum Group Minerals and Gold by Fire Assay Emission Spectrography, Applied Spectroscopy, Volume 50, Number 1, Pages 52-56.
- Coonrad, Warren L. (1978): Geochemical Map Showing Distribution and Abundance of Gold and Platinum in the Vicinity of Platinum, Goodnews and Hagemester Island Quadrangles Region, Southwestern Alaska, United States Geological Survey, Open File 78-95.
- Coulter, C. C. (1936): The Platinum Group Discovery at Centennial Wyoming, Mining Journal (Phoenix, Arizona), Volume 20, Number 6, Page 5.
- Cousins, C. A. (1973): Notes on the Geochemistry of the Platinum Transactions of the Geological Society of South Africa, Volume 76, Pages 77-81.
- Crampton, F. A. (1916): Platinum at the Boss Mine, Goodsprings, Nevada, Mining and Scientific Press, Volume 112, Pages 479-482.
- Crocket, J. H. (1969): Platinum Metals. in Wedepohl, K. H. (editor), Handbook of Geochemistry 11/5, Springer-Verlag, Berlin.

BIBLIOGRAPHY (CONTINUED)

- Crocket, J. H. (1972): Some Aspects of the Geochemistry of Ru, Os, Ir and Pt in Iron Meteorites, Geochimica and Cosmochimica Acta, Volume 36, Pages 517-535.
-(1979): Platinum-Group Elements in Mafic and Ultramafic Rocks - A Survey, Canadian Mineralogist, Volume 17, Pages 391-402 and 889-890.
-(1982): Geochemistry of the Platinum-Group Elements. in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 4, pages 47-64
- Crocket, J. H. and Cabri, L. J. (1982): Analytical Methods for the Platinum-Group Elements. In Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 6, Pages 71-82.
- Crocket, J. H. and Chyi, L. L. (1972): Abundance of Pd, Ir, Os and Au in an Alpine Ultramafic Pluton, 24th, I.G.C., Section 10, Pages 202-209.
- Crocket, J. H., MacDougall, J. D. and Harriss, R. C. (1973): Gold, Palladium and Iridium in Marine Sediments, Geochimica and Cosmochimica Acta, Volume 37, Pages 2547-2556.
- Crocket, J. H. and Teruta, Y. (1977): Palladium, Iridium and Gold Contents of Mafic and Ultramafic Rocks Drilled from the Mid-Atlantic Ridge, Leg 37, Deep Sea Drilling Project, Canadian Journal of Earth Sciences, Volume 14, Pages 777-784.
- Czamanske, G. K., Hatthy, Joseph and Nabbs, S. W. (1984): Pt, Pd and Rh Analyses and Beneficiation of Mineralized Mafic Rocks from the La Perouse Layered Gabbro, Alaska, Economic Geology, Volume 76, Pages 2004-2011.
- Darling, A. S. (1960): Iridium-Platinum Alloys; A Critical Review of their Constitution and Properties: Platinum Metals Review, Volume 4, Number 1, Pages 18-21.

BIBLIOGRAPHY (CONTINUED)

- Darling, A. S. (1961): Rhodium-Platinum Alloys; A Critical Review of their Constitution and Properties: Platinum Metals Review, Part 1, Volume 5, Number 2, Pages 58-65 and Part 2, Volume 5, Number 3, Pages 97-100.
-(1962): Gold-Platinum Alloys; A Critical Review of their Constitution and Properties: Platinum Metals Review, Part 1, Volume 6, Number 2, Pages 60-77 and Part 2, Volume 6, Number 3, Pages 106-111.
-(1963): Cobalt-Platinum Alloys; A Critical Review of Their Constitution and Properties: Platinum Metals Review, Volume 7, Number 3, Pages 96-104.
- Darling, A. S. and Yorke, J. M. (1960): The Ruthenium-Palladium System; Platinum Metals Review, Volume 4, Number 3, Pages 104-110.
- Das Sarma, B., Sen, B. N., and Chowdhury, A. N. (1966): Studies on the Geochemistry of Platinum, Economic Geology, Volume 61, Pages 592-597.
- Day, D. T. (1905-10): Platinum, United States Geological Survey, Mineral Resources, United States, 1904, Pages 359-360; 1906, Pages 551-552; 1907, Part 1, Pages 731-732; 1905-09, Pages 781-791.
-(1913): Platinum and Allied Metals: United States Geological Survey, Mineral Resources United States, 1912, Part 1, Pages 1055-1059; 1913, Part 1, Pages 445-457.
- De Hautpick, E. (1913): Gold and Platinum in Mongolia, Mining Journal, Volume 100, Pages 107.
- Dickson, C. N. (1905): The Distribution of Platinum Metals in Other Sources than Placers: Canadian Mining Institute Journal, Volume 8, Pages 192-214.
- Dudley, W. L. (1885): Iridium, with a bibliography of 149 References from 1803-1885, United States Geological Survey, Resources, United States, 1883-1884, Pages 102-104.

BIBLIOGRAPHY (CONTINUED)

- Dunstan, B. (1920): Platinum, Queensland Geological Survey Department of Mines, Publication 268, Pages 127-172.
- Economic Geology, (1976): An Issue Devoted to Platinum-Group Elements, Economic Geology, Volume 71, Number 7.
- Economic Geology, (1976): An Issue Devoted to Mineral Deposits in Southern Africa, Economic Geology, Volume 71, Number 1.
- Edwards, N. B., Anderson, J. S. and Hart, J. G. (1942): On the Occurrence of Platinum and Palladium at the Thomson River Copper Mine, Victoria, With a Note on the Optical Properties of Braggite, Australasian Institute Mining and Metallurgy Proceedings, Number 125, Pages 61-69.
- Elford, H. S. (1932): Osmiridium. in Tasmania, Engineering and Mining Journal, Volume 133, Pages 580-581.
- Emmons, S. F. (1903): Platinum in Copper Ores in Wyoming, United States Geological Survey, Bulletin 213, Pages 94-97.
- Engelhard, C. W. (1963): Uses of Platinum Metals, Engineering and Mining Journal, Volume 164, Pages 119.
- Finch, J. W. (1925): Platinum Discoveries in Wyoming (Encampment District), Engineering and Mining Journal, Volume 120, Pages 95-96.
- Francis, A. V. (1962): Platinum Group Metals and Rhenium, Metallurgie, Volume 94, Number 3, Pages 233-245.
- Gladhill, T. L. (1921): Iridosmine Crystals from Ruby Creek, Atlin District, British Columbia, Toronto University Studies, Geological Series, Number 12, Page 40.
- Griffith, Laurence (1940): The Determination of Platinum and Palladium in Low Grade Materials, Canadian Institute Mining and Metallurgy, Transactions, Volume 43, Pages 153-158.
- Griffith, W. P. (1967): The Chemistry of the Rarer Platinum Metals (Ir,Os, Ru, Rh), Interscience Publishers Inc., New York, 49 Pages.

BIBLIOGRAPHY (CONTINUED)

Grimaldi, F. S. and Schnepfe, M. M. (1969): Mode of Occurrence of Platinum, Palladium and Rhodium in Chromite, United States Geological Survey, Professional Paper, 650-C, Pages 149-151.

Groeneveld, Meijer, W. O. J. (1955): The Geochemistry of the Platinum Metals with Respect to their occurrence in Nickeliferous Sulphide Deposits (Abstract), Canadian Mining Journal, Volume 76, Page 80.

.....(1955): Synthesis, Structures, and Properties of Platinum Metal Tellurides, American Mineralogist Volume 40, Pages 646-657.

Gruetter, T. W. (1916): Platinum in Compton Mines, Grant County, Oregon, Engineering and Mining Journal, Volume 101, Pages 1135.

.....(1916): Platinum on the Pacific Coast, Mining Science Press, Volume 113, Page 20.

Haffly, J., Riley, L. B. and Goss, W. D. (1977): A manual on the Fire Assaying and Determination of the Nobel Metals in Geological Materials, United States Geological Survey, Bulletin 1445, 58 Pages.

Haffly, J., Haubert, A. W. and Page, N. J. (1980): Determination of Iridium and Ruthenium in Geological Samples By Fire Assay and Emission Spectrograph, United States Geological Survey, Professional Paper 1129-C, Pages G1-G4

Hale, F. A. Jr. (1914): Platinum Ore in Southern Nevada, Boss Mine, Engineering and Mining Journal, Volume 98, Page 641.

Harrington, G. L. (1919 a): The Gold and Platinum Placers of the Tolstoi District, Alaska, United States Geological Survey, Bulletin 692, Pages 339-351.

.....(1919 b): The Gold and Platinum Placers of the Kiwalik-Koyuk Region, Alaska, United States Geological Survey, Bulletin 692, Pages 369-400.

Hausel, D. W., Graff, P. J. and Albert, K. G. (1985): Economic Geology of the Copper Mountain Supracrustal Belt, Owl Creek Mountains, Fremont County, Wyoming, Geological Survey of Wyoming, Report of Investigations.

BIBLIOGRAPHY (CONTINUED)

- Heyding, R. D. and Calvert, L. D. (1961): Arsenides of the Transition Metals, IV - A note on the Platinum Metal Arsenides, Canadian Journal of Chemistry, 39, Pages 955-957.
- Hawley, J. E., and Rimsaite, Y. (1953): Platinum in Some Canadian Uranium and Sulphide Ores, American Mineralogist, Volume 38, Pages 463-475.
- Hiemstra, S. A. (1979): The Role of Collector in the Formation of the Platinum Deposits in the Bushveld Complex, Canadian Mineralogist, Volume 17, Pages 437-452.
- Himmelberg, G. R. , Loney, R. A. and Craig, J. T. (1986): Petrogenesis of the Ultramafic Complex of the Blashke, Southeastern Alaska, United States Geological Survey, Bulletin 1662.
- Hite, T. H. (1933): Fine Gold and Platinum of Snake River, Economic Geology, Volume 28, Page 256-265.
- Hoffman, G. C. (1888): On a specimen of Canadian Native Platinum from British Columbia, Royal Society Canada Proceedings and Transactions 5, Part 3, Pages 17-22.
- Holt, S. P., Shepard, J. G., Thorne, R. L., Tolonen, A. W. and Foose, E. L. (1948): Investigation of the Salt Chuck Copper Mine, Kasaan Peninsula, Prince of Wales Island Southeastern Alaska, United States Bureau of Mines, Report of Investigations, 4358, 16 pages.
- Houston, Robert, S., Schuster, Eric J., Ebbelt, Ballard E. (1975): Preliminary Report on the distribution of Copper and Platinum-Group Metals in the Mafic Igneous Rocks of the Sierra Madre, Wyoming, United States Geological Survey, Open File Report 75-85.
- Howe, J. L. and Holtz, H. C. (1919): Bibliography of the Metals of the Platinum Group; Platinum, Palladium, Iridium, Rhodium, Osmium, Ruthenium, 1748-1917, United States Geological Survey, Bulletin 694, 5882.
- Howe, J. L. and Staff of Baker and Co., Inc. (1947): Bibliography of the Platinum metals, 1918-1930, N. J. Baker and Co. Ltd., New York, 138 Pages.

BIBLIOGRAPHY (CONTINUED)

Howe, J. L. and Staff of Baker and Co - , Inc, (1949): Bibliography of the Platinum Metals, 1931-40, N. J. Baker and Co. Inc., New York, 248 Pages.

.....(1956): Bibliography of the Platinum Metals, 1941-1950, N. J. Baker and Co. Inc., Baker, New York, 93 Pages.

Imperial Institute (1936): Platinum and Allied Metals Mineral Resources Department, 2nd Edition, London, 137 pages.

Irvine, T. N. (1963): Origin of the Ultramafic Complex at Duke Island, Southwestern Alaska, Mineralogical Society of America, Special Paper 1, 1963, Pages 36-45.

.....(1975): Crystallization Sequences in the Muscox Intrusion and other Layered intrusions II: Origin of Chromitite Layers and Similar Deposits of other Magmatic Ores, Geochimica and Cosmochimica Acta, Volume 39, Pages 991-1020.

Johnston, C. W. (1962): Platinum Mining in Alaska; Dredge and Dragline Operations at Goodnews Bay, Platinum Metals Review, Volume 6, Number 2, Pages 68-74.

Jolly, J. H. (1978): Platinum-Group Metals, United States Department of The Interior, Mineral Commodity Profile 22, 23 pages.

.....(1980): Platinum-Group Metals, in Mineral Facts and Problems, 1980 Edition, United States Bureau of Mines, Bulletin 671, Pages 683-7-6.

Kellogg, A. E. (1919): Placer Mining in Oregon (Waldo District, Josephine County), Engineering and Mining Journal, Volume 108, Pages 90-91.

.....(1922): Platinum in Quartz Veins of Southwest Oregon; Engineering and Mining Journal, Volume 113, Page 1000.

.....(1929): Platinum in Southwestern Oregon, Mining Journal (Phoenix, Arizona), Volume 12, Pages 5-6.

BIBLIOGRAPHY (CONTINUED)

- Kemp, J. F. (1902): The Geological Relations and Distribution of Platinum and Associated Metals, United States Geological Survey, Bulletin 193, 95 Pages.
- (1904): Platinum in the Rambler Mine, Wyoming, United States Mineral Resources, United States, 1902, Pages 244-250.
-(1926): General Geological Relationships of Platinum, Engineering and Mining Journal, Volume 121, Pages 717-719.
- Khrostopa, V. P., Golocnya, S. V., Chernysheva, N. V. and Bukhanova, A. I. (1976): Distribution of the Platinum Group Metals in Chromite Ores and Ultramafic rocks of the Ray-Iz Massif (Polar Urals), Geochem International, Volume 13, Pages 35-39.
- Knopf, A. (1915 a): A Gold-Platinum Lode in Southern Nevada, United States Geological Survey, Bulletin 620, Pages 1-18.
- (1915 b): Platinum at the Boss Mine, Mining and Science Press, Volume 110, Pages 867-879.
- Koraloff, Paul (1926): Platinum and Platinoid Metals, A Memorandum Based on Part 3 of "The Geology and World Distribution of Platinum", 1923, -South Africa Mining and Engineering Journal, Volume 37, Pages 519-521, 516-562 and 585-587.
- Lennon, P. O. (1927): Osmiridium Mining in Tasmania, Mining Magazine, Volume 37, Pages 343-349.
- Lipovsky, M. (1962): The Transvaal Platinum Deposits, South Africa Mining and Engineering Journal, Volume 37, Part 2, Pages 273-276 and 405.
- Lundy, David E. (1923): The Platinum Group Metals-An in Depth View of the Industry, 1983, International Precious Metals Institute.
- MacKay, R. A. (1953): The World's Sources of Platinum, Mining Magazine, Volume 89, Pages 216-218.

BIBLIOGRAPHY (CONTINUED)

- MacKenzie, G. C. (1920): An Investigation of Certain Canadian Platinum and Manganese Resources, Canadian Mining Institute Transactions, Volume 22, Pages 305-319.
- Makovicky, Emil, Makovicky, Milota and Rose-Hansen, John (1985): Experimental Studies on the Solubility and Distribution of Platinum Group Elements in Base Metal Sulphides in P.G.M. Deposits (Abstract) in Program and Abstracts, Fourth International Platinum Symposium, Canadian Mineralogis, Volume 23, Part 2, Pages 317-318.
- Macaulay, R. M. (1919): The Source of Placer Platinum in the Tulameen District of British Columbia, Engineering and Mining Journal, Volume 107, Pages 303-336.
- McCallum, M. E. (1968): The Centennial Ridge Gold-Platinum District, Albany County, Wyoming, Geological Survey of Wyoming - Preliminary Report Number 87.
- McCallum, M. E., Loucks, R. R., Carlson, R. R., Cooley, E. F. and Dverge, T. A. (1976): Platinum Metals Associated with Hydrothermal Copper Ores of the New Rambler Mine, Medicine Bow mountains, Wyoming, Economic Geology, Volume 71, Number 7, Paes 1429-1450.
- McCallum, M. E. and Orback, C. J. (1968): The New Rambler Copper-Gold-Platinum. District, Albany and Carbon Counties Wyoming, Geological Survey of Wyoming, Preliminary Report Number 8.
- McCallum, M. E. and Menzer, F. J. (1982): Metal Distribution in the Battle Lake Area, Grand Encampment Mining District, Carbon County, Wyoming: with comparisons to deposits in the New Rambler Mine District, Albany and Carbon Counties Wyoming, United States Geological Survey, Open File 82-179, 20 Pages.
- McCaskey, H. D. (1902): Platinum and Associated Rare Metals in Placer Formations, Bureau of Public Printing - United States, Pages 3-8.
- McDonald, Donald and Hunt, Leslie B. (1982): A History of Platinum and its allied metals - Johnson Mathey, Europa Publications, 450 Pages.
- McTaggart, K. C. (1971): On The Origin of Ultramafic Rocks, Geological Society of America, Bulletin 82, Pages 23-42.

BIBLIOGRAPHY (CONTINUED)

- Mertie, J. B. Jr. (1919): Platinum Bearing Gold Placers of the Kahi Hna Valley, Alaska, United States Geological Survey, Bulletin 692, Pages 233-264.
-(1920): The Salt Chuck Palladium-Copper Mine, Prince of Wales Island, Alaska, Engineering and Mining Journal, Volume 110, Pages 17-20.
-(1939): Platinum Deposits of the Goodnews Bay District, Alaska, United States Geological Survey, Bulletin 910, Pages 17-20.
-(1940b): The Goodnews Platinum Deposits, Alaska, United States Geological Survey, Bulletin 918, 37 Pages.
-(1969): Economic Geology of the Platinum Metals, United States Geological Survey, Professional Paper 630, 120 Pages.
- Mihalik, P., Jacobson, J. B. E. and Hiemstra, S. A. (1974): Platinum-Group Minerals from a Hydrothermal Environment, Economic Geology, Volume 69, Page 257-262.
- Mining Journal (1986): Positive Outlook for Platinum, Mining Journal, Volume 306, Number 7848.
- Ministry of Natural Resources (1979): Platinum Group Metals - Ontario and the World, Ontario Mineral Policy Background, Paper Number 7, Mineral Resources Branch, 162 pages.
- Mitchell, R. H., Keays, R. R. (1981): Abundance and distribution of gold, palladium and iridium in some spinel and garnet lherzolites; implications for the nature and origin of precious metal-rich intergranular components of the upper mantle, Geochimica and Cosmochimica Acta, Volume 45, Pages 2425-2442.
- Molly, E. N. (1959): Platinum Deposits of Ethiopia, Economic Geology, Volume 54, Pages 467-477.
- Mowatt, Thomas C. and Jansons, U. (1985): Platinum and Palladium in some Mafic and Ultramafic Rocks from the Rabbit Creek Area in the Noatak Quadrangle, Alaska, United States Geological Survey, open File Report 85-45.

BIBLIOGRAPHY (CONTINUED)

Munition Resources Commission (1920): Platinum Mineral Investigations, in Final Report of the Work of the Commission, Industrial and Technical Press, Page 134-194.

Naldrett, A. J. (1982): Platinum-Group Element Deposits, in Platinum-Group Elements, Mineralogy, Geology and Recovery, Ed. by L. J. Cabri, C.I.M., Special Volume 23, Chapter 10, Page 197-233.

Naldrett, A. J. and Duke, J. M. (1980): Platinum Metals in Magmatic Sulphide ores, Science, Volume 208, Page 1417-1425.

Naldrett, A. J., Hoffmanm E. L., Green, A. H., Chou, C. L., Naldrett, S. R. and Alcock, R. A. (1979): The Composition of Ni-Sulphide Ores, with particular Reference to their Content of PGE and Au, Canadian Mineralogist, Volume 17, Page 403-416.

Newman, S. C. 1973, Platinum: Institute Mining Metallurgy Transactions, Volume 82, Pages A52-A56

Nicholl, J. J. (1925): Platinum in Mexico, Mining Journal, Volume 149, Page 352.

O'Neill, J. J., The Platinum Situation in Canada, 1918, Geological Survey of Canada, Summary Report, 1918, Part G, Page 1G-19G.

O'Neill, J. J. and Gunning, H. C. (1934): Platinum and Allied Metal Deposits of Canada, Geological Survey of Canada, Economic Geology Series Number 13, 165 Pages.

O'Neill, P. N. (1956): Platinum Mining in Columbia, South America. Mining Engineering, Volume 8, Number 5. Pages 496-500.

Oshin, I. O. and Crocket, J. H. (1982): Noble Metals in Thetford Mines Ophiolites, Quebec, Canada. Part 1: Distribution of Gold, Iridium, Platinum and Palladium in the Ultramafic and Gabbroic Rocks. Economic Geology, Volume 77, Pages 1556-1570.

BIBLIOGRAPHY (CONTINUED)

Page, N. J. (1969): Platinum Content of Ultramafic Rocks, United States Geological Survey, Circular 622, 5 pages.

..... (1985): Platinum in Soils and Rocks from the Lower Coon Mountain Pluton, De Norte County, California, United States Geological Survey, Open File Report 85-149.

Page, N. J. and Carlson, R. R. (1980): Review of Platinum-Group Metal Geochemistry and the Major Occurrences in the World, United States Geological Survey, Open File 80-90, 21 pages.

Page, N. J. and Jackson, E. D. (1967): Preliminary Report on sulphide and Platinum-Group Minerals in the chromite of the stillwater Complex, Montana, in Geological Survey Research, 1967, United States Geological Survey, Professional Paper 575-D. pages D123-126.

Page, N. J., Meyers, J. S., Haffly, J., Simon, F. O. and Aruseqvage, P. J. (1980): Platinum, Palladium and Rhodium in the Fiskingesset Complex, Southwest Greenland, Economic Geology, Volume 75, page 907-915.

Page, N. J., Pallister, J. S., Brown, M. A., Smewing, J. D. and Haffly, J. (1979): Platinum-Group Metals in Chromite Rich Rocks from two Traverses Through the Semial Ophiolite, Oman, EOS, Volume 60, page 963.

Page, N. J., Rowe, L. J. and Haffly, J. (1976): Platinum Metals in the Stillwater Complex, Montana, Economic Geology, Volume 71, pages 1352-1363.

Page, N. J., Riley, L. B. and Haffly, J. (1972): Vertical and Lateral Variations of Platinum, Palladium and Rhodium in the Stillwater Complex, Montana, Economic Geology, Volume 67, pages 915-923.

Page, N. J., Theodore, T. G. and Bradley, L. A. (1986): Discussion of Ultramafic and Mafic Rocks and Platinum-Group Element Analyses from the Lost Basin Mining District, Northwestern Arizona, United States Geological Survey, Open File 86-33, 13 pages.

Patterson, R. M. (1979): Concerning the Origin and Occurrence of Gold, Platinum and Diamond in the United States, United States Geological Survey, Open File 78-9-5.

BIBLIOGRAPHY (CONTINUED)

- Pelzer, E. E. (1950): The Rankin Inlet Nickel-Copper Deposit; a Microscopic Study: Canadian Mining Journal, Volume 71, Number 9, pages 79-83.
- Plafker, G. and MacKerett, E. M. Jr. (1970): Mafic and Ultramafic Rocks from a Layered Pluton at Mount Fairweather, Alaska, in Geological Survey Research Survey 1970, United States Geological Survey, Professional Paper 700-B, pages B21-B26.
- Poiterin, E. (1924): Platiniferous Rocks from Tulameen Map Area, Yale District, British Columbia and Ural Mountains Russia, Geological Survey of Canada, Summary Report 1923, Part A, pages A84-A101.
- Pollet, J. D. (1931): Platinum Mining in Sierra Leone: Engineering and Mining World, Volume 2, pages 747-748.
- Pretorius, D. A. (1981): Abstracts of Papers: Third International Platinum Symposium, Pretoria, South Africa, 28 June - 9 July, 1981, Society of Economic Geologists, Geological Society of South Africa, 47 pages.
- Purrington, C. W. (1899): The Platinum Deposits of the Tura River System, Ural Mountains, Russia, American Institute of Mining Engineers Transactions, Volume 29, pages 3-16.
- Raicevic, D. and Cabri, L. J. (1976): Mineralogy and Concentration of Au and Pt from alluvial gravel from the Tulameen River Area in British Columbia, Canadian Institute Mining Metallurgy, Volume 69.
- Razin, L. V. (1976): Geologic and Genetic Features of Forsterite Dunites and their Platinum-Group Mineralization, Economic Geology, Volume 71, Number 7, pages 1372-1376.
- Razin, L. V. and Kohmenlo, G. P. (1969): Accumulation of Osmium Ruthenium, and other Platinum-Group Metals in Chrome Spinel in Platinum Bearing Dunites, Geochemical International, Volume 6, pages 547-557.
- Read, T. J. (1905): Platinum and Palladium in Certain Copper Ores, Engineering and Mining Journal, Volume 79, pages 985-988.

BIBLIOGRAPHY CONTINUED

- Rose, A. W. (1966): Geology of Chromite-Bearing Ultramafic Rocks near Eklutna, Anchorage Quadrangle, Alaska, Geological Society of America, Bulletin, Volume 70.
- Schmidt, E. R. (1953): The structure and composition of the Merensky Reef and Associated Rocks in the Rustenburg Platinum Mine, Geological Society of South Africa, Transactions, Volume 55, 47 pages.
- Society of Economic Geologists (1975): International Platinum Symposium, Denver, October 16, 17, 18, 1975, Program and abstracts, Society of Economic Geologists.
- Still, Jan C. (1974): Copper, Gold, Platinum and Palladium Sample Results from the Klukwan Mafic Ultramafic Complex, Southeast Alaska, United States Geological Survey, Open File 84-21.
- (1984): Copper, Gold, Platinum and Palladium Sample Results from the Klukwan Mafic/Ultramafic Complex, Southeast Alaska, United States Geological Survey, Open File 84-21, 52 pages.
- Stumpfel, E. F. (1961): Some New Platinoid-Rich Minerals, Identified with the Election Microanalyser, Mining Magazine, Volume 32 pages.
- (1962): Some Aspects of the Genesis of Platinum Deposits; Economic Geology, Volume 57, pages 619-623.
- Stumpfl, E. F. and Tarkian, M. (1976): Platinum Genesis; New Mineralogical Evidence, Economic Geology, Volume 71, Number 7, pages 1451-1460.
- Taft, H. H. (1918): A Wyoming Platinum Mine, Engineering and Mining Journal, Volume 106, page 900.
- Tanton, T. L. (1922): Palladium-Bearing Nickel Deposit at Shebandowan Lake, Thunder Bay District, Ontario, Canadian Geological Survey Report for 1922, Part D, page 1-8.
- Taylor, H. P. Jr (1976): The Zoned Ultramafic Complexes of Southeastern Alaska in Ultramafic and related Rocks, ed. P. J. Wylie, John Wiley and Sons, page 96-118.

BIBLIOGRAPHY (CONTINUED)

Thomson, A. G. (1951): South African Platinum Metals, Mining Journal, Volume 237, page 160-161.

Till, Alison B. and Page, N. J. (1979): Reported Occurrences of Platinum in the USSR, United States Geological Survey, Miscellaneous Field Studies; Map MF-1071.

Tredoux, Marian, Sellschop, J. P. Friedrich; Davies, Gladstone and DeWit Maarton, J. (1985): The Behavior of the PGE and Au during Partial Melting and Fractional Crystallization; Evidence from Magmas on the Kappraal Craton, (abstract) in Program and abstracts; Fourth International Platinum Symposium, Canadian Mineralogist 23, Part 2, pages 317-318.

Turner, H. W. (1922): Platinum in Quartz Veins, Engineering and Mining Journal, Volume 113, pages 488-489.

Tyler, Paul and Santmyers, R. M. (1931): Platinum, United States Bureau of Mines, Information Circular 6389, 69 pages.

Uglow, W. L. (1919): Geology of Platinum Deposits, Parts 1 and II, Engineering and Mining Journal, Volume 108, pages 352-355 and 390-393.

.....(1921): Possibilities for Platinum in Western Canada: Canadian Mining Inst., Monthly Bulletin, No. 95, pages 207-220.

United States Geological Survey: Platinum mineral Resources Reports 1904, pages 359-360; 1906, pages 551-502; 1907, part 1, pages 731-732, 1905-1909, pages 781-791.

Van Wyk, Elwyn and Dixon, Kenneth (1983): The Recovery of Platinum-Group Metals and Gold by the Lead-Collection Step of the Fire Assay Procedure, Council for Mineral Technology (South Africa), Report No. 188, 18 pages.

Vines, R. F. and Wise, E. M. (1941): The Platinum Metals and Their Alloys, The International Nickel Company Inc., New York, 141 pages.

BIBLIOGRAPHY (CONTINUED)


- Vogt, J. H. L. (1927): Geology of the Platinum Metals, Economic Geology, Volume 22, pages 740-741.
- Wagner, P. A. (1925): Notes on the Platinum Deposits of the Bushveld Complex, Geological Society of South Africa Transactions, Volume 28, pages 83-133.
-(1925): The Platinum Deposits on the Western Part of the Lydenburg District, Transvaal, South Africa Journal of Industries, Volume 8, pages 80-113.
-(1926): Preliminary Report on the Platinum Deposits in the Southeastern Part of the Rustenburg District, South Africa Geological Survey, Memoir 24, 39 pages.
- (1926): Occurrence of the Platinum Metals in South Africa, Economic Geology, Volume 231, Number 2, pages 109-134, Number 3, pages 243-270.
-(1926): Recent Discussions on the origin of the Bushveld Platinum Deposits, Geological Society of South Africa, Transactions, Volume 29, pages 145-147.
- Watson, R. J. (1928): Platinum-Bearing Nickel-Copper Deposit on Lower Shebandowan Lake, District of Thunder Bay, Ontario Department of Mines, Annual Report, Volume 37, Part 4, pages 128-149.
- Westland, A. D. (1982): Inorganic Chemistry of the Platinum-Group Elements, in Platinum-Group Elements: Mineralogy, Geology, Recovery, Ed. by L. J. Cabri, C.I.M. Special Volume 23, Chapter 2, pages 5-18.
- Wilson, N. D. B. and Anderson, D. T. (1959): The Composition of Canadian Sulphide Ore Deposits: Canadian Institute Mining and Metallurgy Transactions, Volume 62, pages 324-337.
- Wise, E. M. (1953): The Platinum Metals, Alloys, and Compounds (Wise); and Coordination Compounds (Gilchrist): New York Interscience Encyclopedia Inc., reprinted (1954) by Development and Research Division, The International Nickel Co. Ltd., New York, pages A1-A24.

BIBLIOGRAPHY(CONTINUED)

Wright, R. L. and Fleischer, M. (1965): Geochemistry of the Platinum Metals, United States Geological Survey, Bulletin 1214-A, pages A1-A24.

Wright, R. L., Nagel, J. and McTaggart, K. C. (1982): Alpine Ultramafic Rocks of Southwestern British Columbia, Canadian Journal of the Earth Sciences, Volume 19, pages 1156-1173.

Wright, Thomas Loewelyn and Fleischer, Michael (1985): Geochemistry of the Platinum Metals, United States Geological Survey, Bulletin 1214-A, 24 pages.



Geological Survey Branch
Ministry of Energy, Mines and Petroleum Resources
Province of British Columbia

OPEN FILE 1986-7 ①

**PLATINUM GROUP ELEMENT OCCURRENCES
IN BRITISH COLUMBIA**

SCALE 1:2 000 000

KILOMETRES 0 50 100 150 200

LEGEND

ROCKS	ASSEMBLAGE	GROUP	LITHOLOGY
MISSISSIPPIAN-TRIASSIC	SLIDE MOUNTAIN-STYLVESTER	Antler, Milford, Kalo, Tachina, Lay Range, Geyser, Slide Mountain, Sylvester, Fennel	upper: marine diabase, local gabbro, alpine ultramafics lower: chert, argillite, sandstone, conglomerate
THOMPSON		Kobau, Thompson, Churchill, Corvado, Mount Roberts, Knob Hill, Brooklyn, Attwood	marine volcanoclastics, chert, argillite, limestone, local basalt
CACHE CREEK-ANVIL RANGE		Bridge River, Hozomeen, Darrington, Shukun, Marble Canyon, Chapperton, French Range, Nakina, Cache Creek, Redonda, Pavilion, Anvil Range	marine and non-marine basalt, limestone, argillite, chert, melange
STICKER-SKOLAI		Slicker, West Coast, Skolai, Battle Lake	marine and non-marine (locally) basalt, andesite, rhyolite, tuff, carbonate, argillite, chert
IGNEOUS INTRUSIONS			
MID-CRETACEOUS	ALASKAN-TYPE ULTRAMAFICS	Alaskan Panhandle	dunite, peridotite, pyroxenite, hornblende, gabbro, diorite
UPPER TRIASSIC-LOWER JURASSIC	ALASKAN-TYPE ULTRAMAFICS	Tulameen, Tarnagait River, Pularis, Luan Creek, Meslinka, Wrede Creek	dunite, peridotite, pyroxenite, hornblende, gabbro, diorite
LATE PALEOZOIC-TRIASSIC	ALPINE-TYPE ULTRAMAFICS	Brulonne, Twin Sisters, Shulaps, Cassiar, Nahlin, Trembleur	dunites, peridotites

SYMBOLS

TRANSFORM FAULT
THrust FAULT, TEETH ON UPPER PLATE
NORMAL FAULT, TEETH ON UPPER PLATE
UNDEFINED FAULT
PLATINUM-GROUP ELEMENT LODE OCCURRENCE	●
PLATINUM-GROUP ELEMENT PLACER OCCURRENCE	○
CHROMITE OCCURRENCE	□
IRIDIUM OCCURRENCE	△
DRAINAGE WITH REPORTED PLATINUM VALUES, POTENTIAL VALUES	~~~~~

Legend and geology modified from Geological Survey of Canada Map 1050A, 1981 and Parrish et al., 1985 in Field Guides to Geology and Mineral Deposits in the Southern Canadian Cordillera-GSA Section Meeting, May, 1985.
Compiled by V.J. Rublee, June 1986.

- | LODE | PLACER |
|---------------------|---------------------------|
| 1. Tulameen | (19) Tulameen River |
| 2. Scottie Creek | (20) Shuttleworth Creek |
| 3. Keefe's | (21) Rock Creek |
| 4. Mastodon | (22) Franklin Creek |
| 5. Swede | (23) Kaslo River |
| 6. Tofino Nickel | (24) Toby Creek |
| 7. Kitkat | (25) Lardeau River |
| 8. Mineral Dyke | (26) Isaac Creek |
| 9. Giant Mascot | (27) Clearwater River |
| 10. Mt. Ida | (28) Tranquille River |
| 11. Mother Lode | (29) Criss Creek |
| 12. Nome | (30) Scottie Creek |
| 13. Cable | (31) Vale |
| 14. Franklin | (32) Van Winkle Bar |
| 15. Sappho | (33) Foster's Bar |
| 16. Sullivan | (34) Glasgow |
| 17. Nickel Plate | (35) Quesnel River |
| 18. Copper Mountain | (36) Bullion Pit |
| | (37) Twenty-Mile Creek |
| | (38) Horsefly River |
| | (39) Cottonwood River |
| | (40) Hixon Creek |
| | (41) Dog Creek |
| | (42) McLeod River |
| | (43) Rainbow Creek |
| | (44) Bill Cus's Bar |
| | (45) Manson Creek |
| | (46) Ah Hoo Creek |
| | (47) Branham Flats |
| | (48) Pete's Toy Bar |
| | (49) McConnel Creek |
| | (50) Thibert Creek |
| | (51) Ruby Creek |
| | (52) Boulder Creek |
| | (53) Toe Hill |
| | (54) Wreck (Florence) Bay |
| | (55) Potato Creek |
| | (56) Lilloet River |

THIS PROJECT IS A CONTRIBUTION TO THE CANADA-BRITISH COLUMBIA MINERAL DEVELOPMENT AGREEMENT, 1985-1990