

**LEGEND**

**(1) TERRAIN UNIT SYMBOLS**

**Simple Terrain Units:**

e.g. Ft (SURFICIAL MATERIAL)(surface expression)

Note: Two letters (in order of decreasing importance based on their areal extent) may be used to describe surface expression, or letters may be omitted if information is lacking.

**Composite Units:** Two or three groups of letters are used to indicate that two or three kinds of terrain are present within a map unit.

e.g. Mv+R indicates that both components are approximately equal in proportion.  
e.g. Mv/R indicates that "Mv" is more extensive than "R".  
e.g. Mv/R indicates that "Mv" is considerably more extensive than "R".

**Stratigraphic Units:** Groups of letters are arranged one above the other where one or more kinds of surficial material overlie a different material. Venues are assumed to overlie bedrock, unless otherwise noted in the stratigraphic unit label.

e.g. Cv FG indicates that "Cv" overlies "FG".

**(2) MATERIALS**

A	Anthropogenic	Culturally-made or culturally-modified geological materials such that their initial physical properties (e.g. structure, cohesion, compaction) have been drastically altered. Includes excavations and tailings from mining operations.
C	Colluvial	Materials that have reached their present positions as a result of direct, gravity-induced movement involving no agent of transportation such as water or ice, although the moving material may have contained water and/or ice. Includes talus and debris flow deposits, as well as reworked till on steep slopes.
F	Fluvial	Materials transported and deposited by streams and rivers, synonymous with alluvial. Includes well-sorted sand, gravel, and overbank silt in post-Fraser glaciation floodplains, terraces, and fans.
FG	Glaciofluvial	Materials that exhibit clear evidence of having been deposited by glacial meltwater streams either directly in front of, or in contact with, glacier ice. Includes sand and gravel, often stratified, which may show evidence of ice melting (slumped structures). Features include deltas, kame terraces, river terraces, and meltwater channels.
LG	Glaciolacustrine	Lacustrine materials deposited in or along the margins of glacial (ice-dammed) lakes, including sediments that were released by the melting of floating ice. Includes laminated or bedded silt and clay, and may contain ice-crafted stones.
M	(Moraine) till	Material deposited directly by glacier ice without modification by any other agent of transportation. Includes three subclasses.
M <sup>1</sup>		Light to dark brown, moderately to highly consolidated lodgement till derived from phyllite, chlorite schist and greenstone, limestone and dolostone, quartzite and slate (Eagle Bay Assemblage). Texture of matrix is generally a mix of silt and sand. Average clast size is 1-2 cm; largest clasts are large boulders (over 2 m).
M <sup>2</sup>		Light to medium grey, loose, sandy, clay-supported ablation till derived from granite and granodiorite of the Baldy Batholith. Generally found in boulder fields at moderate to high elevations. In the southern half of the Adams Plateau map area, M <sup>2</sup> is derived from Eagle Bay Assemblage rocks (described for M <sup>1</sup> ).
M <sup>3</sup>		Light to medium grey, very highly consolidated lodgement till derived from granite and granodiorite of the Baldy Batholith. Texture of matrix is generally a mix of coarse sand and clay. Average clast size is 0.25 cm, approximately the size of mineral grains within the granite. Maximum clast size is only 5-10 cm.
O	Organic	Sediments composed largely of organic materials resulting from the accumulation of vegetative matter. They contain at least 30% organic matter by weight (17% or more organic carbon). Includes swampy areas on high plateaus and on floodplains.
R	Bedrock	Bedrock outcrops and rock covered by a thin mantle (up to 10 cm thick) of unconsolidated or organic materials. Includes granite and granodiorite of the Baldy Batholith, metasedimentary and metavolcanic rocks of the Fennel Formation and Eagle Bay Assemblage.

**(3) SURFACE EXPRESSION**



b	Blanket	A mantle of unconsolidated materials thick enough to mask minor irregularities of the surface of the underlying unit, but still conforms to the general underlying topography. Greater than 1 m thick, and possesses no construction forms typical of the materials genesis; outcrops of the underlying unit are rare.
c	Cone	A cone or segment of a cone with a relatively smooth slope gradient from apex to toe greater than 15°, and a longitudinal profile that is either straight, concave or convex.
d	Depression	Circular or irregular area of lower elevation (hollow) than the surrounding terrain and marked by an abrupt break in slope; side slopes within the depression are steeper than the surrounding terrain; depressions are two or more metres in depth.
f	Fan	A relatively smooth segment of a cone with a slope gradient from apex to toe, up to, and including 15°, and a longitudinal profile that is either straight, concave, or convex.
h	Hummock	Sleep sided hillocks and hollows of unconsolidated material with multidirectional slopes dominantly between 15 and 35°. Local relief is greater than 1 metre. In plan, an assemblage of non-linear, generally chaotic forms that are rounded or irregular in cross-profile.
p	Plain	A level or very gently sloping, unidirectional surface with gradients up to and including 3°, local surface irregularities generally have a relief of less than 1 metre.
r	Ridged	Elongate hillock with slopes dominantly between 15° and 35° if composed of unconsolidated materials; bedrock slopes may be steeper. Local relief is greater than 1 m. In plan, an assemblage of parallel or sub-parallel linear forms.
t	Terraced	A single or assemblage of step-like forms where each step-like form consists of a scarp face and a horizontal or gently inclined surface above.
v	Veneer	A mantle of unconsolidated materials too thin to mask the minor irregularities of the surface of the underlying material. It ranges in thickness from 10 cm to 1 metre, and possesses no form typical of the material genesis.

**CRITERIA FOR DETERMINING LINES/BOUNDARIES:**

—	(Solid) Well-defined, sharp boundaries that can be precisely delimited at the scale of mapping.
- - -	(Dashed) Boundaries that are gradational over a short distance or that can be only approximately located, or where precise boundary locations are masked by forest.
.....	(Dotted) Assumed boundaries, and boundaries that are gradational over considerable distances.

**SYMBOLS**

	Drumlin, drumlinoid ridge		Meltwater channel (minor)
	Lineations		Till sample site
	Striae, glacial grooves		Fieldcheck site
	Meltwater channel (major)		Stratigraphic section



**BRITISH COLUMBIA**  
Ministry of Employment and Investment  
Energy and Minerals Division  
Geological Survey Branch

Library  
Energy, Mines and  
Petroleum Resources  
Victoria, B.C.

Geological Survey Branch  
**OPEN FILE 1997-6**  
**TERRAIN GEOLOGY OF THE  
NORTH BARRIERE LAKE AREA**  
NTS 82 M/5  
By E.R. Leboe (G.I.T.), P.T. Bobrowsky (P.Geo.),  
A. Dixon-Warren, and A. Ledwon  
1:50 000

For an overview of the Quaternary geology of the North Barriere Lake area please refer to the following reports: "Eagle Bay Project: Till geochemistry of the Adams Plateau (82 M/4) and North Barriere Lake (82 M/5) map areas" by P.T. Bobrowsky et al. and "Eagle Bay Project: Surficial geology of the Adams Plateau (82 M/4) and North Barriere Lake (82 M/5) map areas" by A. Dixon-Warren et al. both in Geological Fieldwork 1996, Lefebvre, D.V., McMillan, W.J., and McArthur, G., Editors, B.C. Ministry of Employment and Investment, Paper 1997-1. Geology based on air photo interpretation by E.R. Leboe, followed by ground truthing in areas indicated in 1997-1. Geology based on air photo interpretation by E.R. Leboe, followed by ground truthing in areas indicated in 1997-1.

**QUATERNARY GEOLOGY NOTES**

**QUATERNARY STRATIGRAPHY OF SOUTH-CENTRAL BC**

The present-day landscape of the North Barriere Lake map area is the result of two cycles of glaciation, one interglacial, and early-Holocene erosion and sedimentation (Fulton and Smith, 1978; Ryder et al., 1991). Although not necessarily present in this map area, the following lithologic units and their correlative glacial climate units have been identified in south-central British Columbia.

The oldest Quaternary deposits, identified only at two locations some 60 and 100 km to the south, are the interglacial Westwood Sediments. These deposits consist of cross-stratified gravelly sand capped by marl, sand, silt, and clay, all of which are equivalent to the Highbury non-glacial interval in the Fraser Lowland (Sangamonian). Next in age are Okanagan Centre Drift deposits, consisting of coarse, poorly stratified gravel, till, and laminated silt, identified at Heffley Creek (20 km south of the map area), and elsewhere farther south. These sediments were deposited during the Okanagan Centre Glaciation, equivalent to the Semahmoo Glaciation in the Fraser Lowland (early Wisconsinan). Middle Wisconsinan, Olympic Non-Glacial Besette Sediments overlie the Okanagan Centre Drift. They consist of nonglacial silt, sand and gravel with some organic material and up to two tephras. The Kamloops Lake Drift overlies the Besette sediments, and underlies the present-day surface cover of postglacial deposits. This unit consists of silt, sand, gravel, and till deposited during the Fraser Glaciation (Late Wisconsinan).

The surface and near-surface sediments mapped in the North Barriere Lake map area directly result from the last cycle of glaciation and deglaciation (Fraser Glaciation), and ensuing postglacial activity.

**FRASER GLACIATION**

The onset of Fraser Glaciation in the map areas began in the Coast, Cariboo, and Monashee Mountains. Valley glaciers descended to form piedmont lobes on the Interior Plateau, and eventually coalesced to form a mountain ice sheet (Ryder et al., 1991). Ice sheet margins reached a maximum elevation between 2200 and 2400 m along rimming mountains. The entire Shuswap Highland, except perhaps Dinn Peak (2630 m) and higher peaks to the north, was completely buried beneath an ice cap by approximately 19 ka. At Fraser Glaciation maximum, regional ice flow was to the south-southeast, with deviations up to 45° (Fulton et al., 1986). Flow was locally diverted down valleys. Basal till deposits, which range widely in texture with the underlying bedrock, now blanketed the land surface.

Deglaciation of the Interior Plateau was rapid, the equilibrium line rose considerably, reducing the area of accumulation for the Cordilleran ice sheet, and the ice mass decayed by downwasting. Ablation till was deposited by stagnating ice in several high-elevation portions of the map areas. As uplands were deglaciated prior to low benches and valleys, meltwater was channelled to valley sides, resulting in kame terraces and ice-contact sediments. Valleys clear of ice above the stagnating glaciers in their lower reaches became the confinement for blocking meltwater drainage, resulting in local mantles of glaciolacustrine sediments. Radiocarbon dates of 11.3 ka immediately north of the North Barriere Lake map area indicates that deglaciation began shortly before this time (Dyck et al., 1965). Minor, local ice readvances occurred in some areas of the Shuswap Highland (Duford and Osborn, 1973).

**HOLOCENE POST-GLACIAL**

Once ice-dammed lakes were released, meltwaters carrying heavy sediment loads deposited thick units of stratified sand and gravel in valleys. As sediment loads decreased, deposition was replaced by erosion, and water courses cut down through valley fills, leaving glaciofluvial terraces abandoned on valley sides. Immediately following the complete deglaciation of the region, unstable and unvegetated slopes were highly susceptible to erosion and sedimentation. Intense mass wasting of surface deposits on oversteepened valley slopes resulted in the deposition of colluvial fans and aprons along valley bottoms. Most post-glacial deposition occurred within the first few hundred years of deglaciation, and certainly before the eruption of Mt. Mazama, 6.6 ka, which deposited tephra near the present-day ground surface.

**REFERENCES**

Duford, J.M. and Osborn, G.D. (1973). Holocene and latest Pleistocene cirque glaciations in the Shuswap Highland, British Columbia. *Canadian Journal of Earth Sciences*, 15: 865-873.

Dyck, W., Fyles, J.G. and Blake, W., Jr. (1965). Geological Survey of Canada Radiocarbon Dates IV; *Geological Survey of Canada*, Paper 65-4, 23 p.

Fulton, R.J. and Smith, G.W. (1978). Late Pleistocene stratigraphy of south-central British Columbia. *Canadian Journal of Earth Sciences*, 15: 971-980.

Fulton, R.J., Alley, N.F., and Achard, R.A. (1986). Surficial geology, Seymour arm, British Columbia. *Geological Survey of Canada Map 1609A*, 1:250 000.

Ryder, J.M., Fulton, R.J., and Clague, J.J. (1991). The Cordilleran ice sheet and the glacial geomorphology of southern and central British Columbia. *Géographie Physique et Quaternaire*, 45: 365-377.

