
BRITISH Energy, Mines and Petroleum Resources

British Columbia Geological Survey Open File 2017-05

Basal till potential of the Christensen Creek map area (NTS 093C/11), British Columbia

Geoscience BC Map 2017-02-04

D. Sacco, H. Arnold, T. Ferbey and W. Jackaman



Scale 1:50,000

NOTE: Where map units are composed of multiple surficial materials, a compound map unit designator is used, separating more extensive materials from less extensive (e.g., for Tb.Tv, Tb is more extensive than Tv).

QUATERNARY

HOLOCENE

ORGANIC DEPOSITS: Mostly saturated organic materials, consisting mainly of mosses, sedges, or

other hydrophytic vegetation. Undifferentiated organic deposits: Bogs, fens, and swamps; generally occur where

shallow lakes are infilled and in depressions along floodplains and abandoned meltwater

COLLUVIAL DEPOSITS: Materials deposited by direct gravity-induced movement; lithologic composition dependent on source material; typically poorly sorted, massive to crudely stratified

Cv Colluvial veneer: Deposits less <2 m thick that conform to underlying topography; typically on steep slopes.

LATE WISCONSINAN

PROGLACIAL AND GLACIAL ENVIRONMENT GLACIOLACUSTRINE DEPOSITS: Well sorted, stratified sand, silt, and clay deposited by suspension settling and interflow and underflow currents; diamictons released from floating ice or colluviated from valley sides into glacial lakes; grain size may increase, and sorting may decrease in ice proximal environments.

GLv Glaciolacustrine veneer: Deposits <2 m thick that conform to underlying topography; predominantly fine-grained material or silt-rich diamictons.

GLb Glaciolacustrine blanket: Deposits >2 m thick that mask subtle variations in substrate

surface but generally conform to underlying topography; predominantly in areas of low relief.

GLACIOFLUVIAL DEPOSITS: Poorly sorted to well sorted sand and gravel transported and deposited directly by glacial meltwater.

GFv Glaciofluvial veneer: Deposits <2 m thick that conform to underlying topography.

ablation till (e.g., Tb.Cb; Tv.R).

Surficial material other than till.

GFb Glaciofluvial blanket: Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography.

Only thick basal till (Tb or Ts); may contain lesser amounts of thin basal

Only thin basal till (Tv); may contain lesser amounts of thick basal till

Basal till with lesser amounts of another surficial material, excluding

Another surficial material with lesser amounts of basal till (e.g., Cb.Tb).

Only ablation or ridged till at surface (Th,Tu, or Tr), or ablation or ridged till and another surficial material, excluding basal till (e.g., Th.O; GFb.Tu;

Basal till with lesser amounts of ablation till (e.g., Tb.Tu).

GFr Eskers: Sinuous ridges of stratified sand and gravel deposited in subglacial, englacial, or supraglacial channels.

BASAL TILL DEPOSITS: Diamictons eroded, transported and deposited at the base of an active glacier. They are dense, massive, and matrix supported and can be fissile and jointed. Matrix is typically composed of silt, sand, and clay. Clasts are often sub-angular to sub-rounded and can be faceted and striated.

- **Till veneer:** Deposits <2 m thick that conform to underlying topography; predominantly in
- upland regions with isolated bedrock exposures.
- Till blanket: Deposits >2 m thick that mask subtle variations in substrate surface but generally conform to underlying topography; predominantly in areas of low relief.
- Ts Streamlined till: Flutings, drumlins, and the sediment (down-ice) part of crag-and-tails.

ABLATION TILL DEPOSITS: Diamictons deposited by melt out, commonly from stagnant ice, of far-

travelled supraglacial and englacial material. These deposits typically lack the density of basal till and

have a high percentage of matrix sand. May be stratified and include sorted sands and gravels. **Undulating till:** Loose, sandy diamicton commonly representing thinner deposits near the

margins of widespread ice stagnation, or in depressions where localized ice stagnation occurred; consist of hillocks and hollows with slopes ≤ 15°.

widespread ice stagnation; consist of hillocks and hollows with slopes ≥ 15°. OTHER TILL DEPOSITS: Diamictons composed of subglacial, englacial, and (or) supraglacial

Hummocky till: Loose sandy diamicton commonly representing thicker deposits and

debris produced by glacial thrust, push, or meltout processes. Tr Ridged till: Elongate ridges oriented perpendicular to ice-flow direction, formed subglacially

PRE-QUATERNARY BEDROCK: Lithology varies greatly across the map area and includes sedimentary, metamorphic, volcanic, and intrusive rocks of Permian to Quaternary age; outcrop is generally limited to areas of

Undifferentiated bedrock: High-angle slopes in upland areas or in incised meltwater channels; may be susceptible to rock fall; hummocky, or undulating expressions are the result of glacial or meltwater erosion, or preferential erosion due to structural weaknesses; streamlined bedrock is the result of glacial erosion.

TILL SAMPLES (Labeled with sample number; Jackaman et al., 2015) ICE-FLOW INDICATORS (Arnold et al., 2016)

Drumlin, Drumlinoid or fluting (flow direction known, unknown).

National and Provincial parks and protected areas

DEPOSIT TYPE*

Striation (flow direction known, unknown).

No MINFILE occurrences in map area, as of November 2017. Ablation till with lesser amounts of basal till (e.g., Tu.Tb). *If deposit type is available, see Lefebure and Ray (1995), Lefebure and Höy (1996), and Simandl et al. (1999) for mineral deposit profile codes and definitions.

MINERAL OCCURRENCES

MINFILE NUMBER NAME

North American Datum 1983 Universal Transverse Mercator Zone 10 North Shuttle Radar Topography (SRTM) DEM, 1 arcsecond (30 m) resolution

DESCRIPTIVE NOTES

The Christensen Creek map area is in the Fraser Plateau, a physiographic subdivision of the Interior Plateau defined by a flat to gently rolling topography. Glacial sediments cover much of the region, and bedrock outcrops are rare (Holland, 1976). Previous work in the area includes soils and terrain mapping by Baender (1980) and glacial features mapping by Tipper (1971). To the east, Kerr and Giles (1993a, b) and Proudfoot and Allison (1993a, b) completed surficial geology mapping. Bedrock geology was originally mapped by Tipper (1969) and has been updated since by Souther and Souther (1994), Bordet 2014), and Angen et al. (2017). The present basal till potential map continues the series published by Sacco et al. (2014a to j) for Geoscience BC's Targeting Resources for Exploration and Knowledge (TREK) project area (Clifford and Hart, 2014; Sacco et al., 2014k; Sacco and Jackaman, 2015).


Surficial sediment geochemical and mineralogical anomalies can be used to locate buried bedrock mineralization (Saarnisto, 1990; Klassen, 2001). Basal till is ideal for assessing bedrock hosted mineral potential in areas covered by Quaternary sediments because it is commonly a first derivative of bedrock (Shilts, 1993), has a relatively simple and predictable transport history, and produces a geochemical and mineralogical signature that is more extensive than its bedrock source (Levson, 2001). Glacial transport and deposition of basal till produces a dispersal train elongated down ice from its bedrock source (Fig. 1) To date, all till orientation surveys conducted in British Columbia have identified known mineralized sources (Plouffe et al., 2016).

The purpose of the basal till potential map series is to assist in the design of surface sediment exploration programs by identifying areas where basal till is most likely to occur. Ice flow indicators compiled by Arnold et al. (2016) are included in the maps to illustrate the general transport directions of basal till. These data should be supplemented with additional field measurements to assess for local variability.

The basal till potential map series builds on earlier drift exploration potential maps developed by Proudfoot et al. (1995). Existing surficial geology, terrain, or soils and landform mapping data were reviewed and updated to produce the maps. Map unit definitions are based on conventions outlined by Cocking et al. (2016) and Deblonde et al. (2012) and unit colours are related to basal till potential classes. Each unit includes a label that describes the surficial material within it (mainly unconsolidated sediments) and its surface expression (individual plan-view forms and patterns of forms; Howes and Kenk, 1997).

New mapping focused on distinguishing basal till (Fig. 2) from ablation till (Fig. 3) which, because of a more complex transport and depositional history, is ill-suited for mineral exploration. The relationship between surface expression and till facies is predictable (Maynard, 1989; Aario and Peuraniemi, 1992; Proudfoot et al., 1995; Spirito et al., 2011; McClenaghan et al., 2013). For example, blanket, veneer, LONGITUDINAL SECTION and streamlined units typically contain basal till facies, whereas undulating and hummocky units typically contain ablation till facies. Based on these relationships we used air photographs supplemented by sparse field data to construct -4

Basal till consists of sediment eroded, transported, and deposited at the base of an active glacier (Dreimanis, 1989). It typically has a relatively subdued surface expression that either follows underlying topography (Tb, Tv) or is streamlined



in the direction of ice flow (Ts). It is a dense, massive, matrixsupported diamicton, with a matrix mainly of silt with lesser (dark blue) define the head of a dispersal train at surface, and decrease exponentially in the References cited amounts of sand and clay (Fig. 2). Vertical joints and down-ice direction (light purple). Note how the head of a dispersal train is offset, in the Aario, R., and Peuraniemi, V., 1992. Glacial dispersal of till down-ice direction, from its subcrop source.

subhorizontal fissility intersections can give basal till a blocky appearance in section. Clasts are mostly subangular to subrounded and are commonly striated. The transport path of basal till is relatively simple and short and can be established by measuring the azimuth of ice-flow indicators produced by subglacial flow. However, multiple ice-flow events can create a more complex transport path, highlighting the importance of ice-flow history reconstructions (Ferbey and Levson, 2009; Plouffe et al., 2016).

Compared to basal till, the transport distance of ablation till is longer and the depositional history more complex. Ablation till consists of material transported in the englacial and supraglacial environments and commonly deposited by passive melt out processes. Melting of remnant ice-blocks mantled or surrounded by glacial debris produces irregular, undulating to hummocky topography (Tu, Th). Ablation till is less consolidated, has a higher percentage of gravel-sized material and a sandier matrix (Fig. 3). It can be massive to crudely stratified and may contain lenses of sorted sand and gravel. Deposited during deglaciation, ablation tills are typically the youngest Late Wisconsinan till facies exposed at surface, and can overlie basal tills. Windows through an ablation till, into an underlying basal till, can exist but may be

In a basal till potential map, each unit with till, as a primary or secondary surface material is assigned a basal till potential rating. High potential is assigned to units containing mainly basal till. The highest potential category (1) includes till blankets (>2 m thick) and streamlined till with some till veneer (<2 m thick). In these units, samples can be collected from most exposures. In the second category of high potential (2), till veneers are predominant and likely include some bedrock exposures. In these areas. sample collection may be most productive down-ice from bedrock outcrop, where till might be sufficiently thick to avoid post-depositional surface processes such as pedogenesis. In the third category of high

potential (3), map units are mostly basal till (Tb, Ts, Tv) with lesser amounts of another surface material (excluding Tu or Th). Knowledge of the surface expression of this secondary material, which is provided in the map unit label, will assist in targeting basal till.

Moderate potential is assigned to units containing varying amounts of basal till and ablation till. These map units typically represent (4) thick basal till deposits in depressions or small valleys where ablation till has been deposited, or (5) near the margins of extensive ablation till map units where basal till may be found within a few metres of surface or in areas of higher elevation where ablation till thins.

Low potential (6) is assigned to units consisting mainly of surface material other than till. These areas may include basal till deposits that are too small to resolve at the current map scale, or are discontinuous. Poor potential (7) is assigned to areas of thick ablation till. These areas typically consist of hummocky ablation till and may include lesser amounts of another surficial material (e.g., ice-contact glaciofluvial deposits). These areas are still mapped as having potential because underlying basal till deposits may be present at

We thank L.B. Aspler and A.S. Hickin for reviewing this map. The British Columbia Ministry of Forests, Lands and Natural Resource Operations provided access to stereo models and

constituents in morainic landforms of different types. Geomorphology, 6, 9-25. Angen, J.J., Rahimi, M., Hart, C.J.R., Westberg, E., Logan, J.M. and Kim, R., 2017. Bedrock geology TREK proiect area, northern Interior Plateau, central British Columbia; Geoscience BC Map 2017-06-01 and MDRU Map 12-2017, 1:250,000 scale. Arnold, H., Ferbey, T., and Hickin, A.S., 2016. Ice-flow indicator compilation, British Columbia and Yukon. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2016-04, Geological Survey of Canada Open File 8083, 1:1,750,000 scale Baender, H., 1980. Upper Dean River resource folio, terrain. British Columbia Ministry of Environment.

unpublished, 1:50,000 scale. Bordet, E., 2014. Eocene volcanic response to the tectonic evolution of the Canadian Cordillera. Unpublished Ph.D. thesis, University of British Columbia, 488 p. Clifford, A., and Hart, C.J.R., 2014. Targeting Resources through Exploration and Knowledge (TREK). Geoscience BC's newest minerals project, Interior Plateau region, central British Columbia (NTS 093B, C, F, G). In: Geoscience BC Summary of Activities 2013, Geoscience BC, Report 2014-1,

Cocking, R.B., Deblonde, C., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Laviolette, A., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2016. Surficial data model, version 2.2.0: Revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps. Geological Survey of Canada, Open File 8041, 45 p. Deblonde, C., Plouffe, A., É. Boisvert, Buller, G., Davenport, P., Everett, D., Huntley, D., Inglis, E., Kerr, D.

Moore, A., Paradis, S.J., Parent, M., Smith, I.R., St. Onge, D., and Weatherston, A., 2012.



Fig. 2. Basal till in vertical exposure. Note blocky appearance. Granule and coarser-sized clasts float in a claysilt matrix. Measuring tape in centimetres.

Science language for an integrated Geological Survey of Canada data model for surficial geology maps, version 1.2. Geological Survey of Canada, Open File 7003, 224 p. Dreimanis, A., 1989, Tills: their genetic terminology and classification, In: Goldthwait, R.P., and Matsch. C.L. (Eds.), Genetic Classification of Glacigenic Deposits, Balkema, Rotterdam, 17-83. Ferbey, T., and Levson, V.M., 2009. The influence of ice-flow reversals on the vertical and horizontal distribution of trace elements in tills, Huckleberry mine area, west-central British Columbia. In: Paulen, R.C. and McMartin, I. (Eds.), Application of Till and Stream Sediment Heavy Mineral and Geochemical Methods to Mineral Exploration in Western and Northern Canada, Geological Association of Canada, Short Course Notes 18, pp. 145-151. Holland, S.S., 1976. Landforms of British Columbia: a physiographic outline. British Columbia Ministry of Energy and Mines, and Petroleum Resources, British Columbia Geological Survey Bulletin 48, Howes, D. E., and Kenk, E., 1997. Terrain classification system for British Columbia (revised Edition).

British Columbia Ministry of Environment, Manual 10 (Version 2), 102 p. Jackaman, W., Sacco, D.A., and Lett, R.E., 2015. Regional geochemical and mineralogical data, TREK project – year 2, Interior Plateau, British Columbia. Geoscience BC, Report 2015-12, 13 p. Kerr, D.E., and Giles, T.R., 1993a. Surficial geology of the Chilanko Forks area (NTS 93C/1). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-16, 1:50,000 scale. Kerr, D.E., and Giles, T.R., 1993b. Surficial geology of the Chezacut area (NTS 93C/8). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open

Fig. 3. Ablation till exposed in road cut with higher percentage of sand and gravel, and lower density, compared to typical basal tills (see Fig. 2). Pick for scale (65 cm).

Klassen, R.A., 2001. A Quaternary geological perspective on geochemical exploration in glaciated terrain. In: McClenaghan, M.B., Bobrowsky, P.T., Hall, G.E.M., and Cook, S.J. (Eds.), Drift Exploration in Glaciated Terrain, Geological Society, Special Publication 185, p. 1-17. Lefebure, D.V., and Ray, G.E. (Eds.), 1995. Selected British Columbia mineral deposit profiles, volume 1 metallics and coal. British Columbia Ministry of Energy, Mines, and Petroleum Resources,

British Columbia Geological Survey Open File 1995-20, digital. efebure. D.V., and Höy, T. (Eds.), 1996. Selected British Columbia mineral deposit profiles, volume 2 – more metallic deposits. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1996-13, digital. Levson, V.M., 2001. Regional till geochemical surveys in the Canadian Cordillera: sample media, methods, and anomaly evaluation. In: McClenaghan, M.B., Bobrowsky, P.T., Hall, G.E.M. and Cook, S.J. (Eds.), Drift Exploration in Glaciated Terrain, Geological Society, Special Publication

Maynard, D.E., 1989. Terrain analysis in exploration geochemistry. Explore, 65, 14-16. McClenaghan, M.B., Plouffe, A., McMartin, I., Campbell, J.E., Spirito, W.A., Paulen, R.C., Garrett, R.G., Hall, G.E.M., 2013. Till sampling and geochemical analytical protocols used by the Geological Survey of Canada. Geochemistry: Exploration, Environment, Analysis, 13, 285-301. Miller, J.K., 1984. Model for clastic indicator trains in till. In: Prospecting in areas of glaciated terrain. Institution of Mining and Metallurgy, London. pp. 69-77.

Plouffe, A., Ferbey, T., Hashmi, S., and Ward, B.C., 2016. Till geochemistry and mineralogy: vectoring towards Cu porphyry deposits in British Columbia, Canada. Geochemistry: Exploration, Environment, Analysis, http://doi.org/10.1144/geochem2015-398. Proudfoot, D.N., and Allison, R.F., 1993a. Surficial geology of the Clusko River area (NTS 93C/9). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-18, 1:50,000 scale. roudfoot, D.N., and Allison, R.F., 1993b. Surficial geology of the Toil Mountain area (NTS 93C/16). British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 1993-19, 1:50,000 scale.

Proudfoot, D.N., Bobrowsky, P.T., and Meldrum, D.G., 1995. Drift exploration potential maps derived from terrain geology maps. In: Bobrowsky, P.T., Sibbick, S.J., Newell, J.M., and Matysek, P.F. (Eds.), Drift Exploration in the Canadian Cordillera, British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Paper 1995-2, p. 23-31. Saarnisto, M., 1990. An outline of glacial indicator tracing. In: Kujansuu, R., and Saarnisto, M. (Eds.), Glacial Indicator Tracing, Balkema, Rotterdam, 1-13. Sacco, D.A., and Jackaman, W., 2015. Targeted geochemical and mineralogical surveys in the TREK project area, central British Columbia (parts of NTS 093B, C, F, G): year two. In: Geoscience BC

Summary of Activities 2014, Geoscience BC, Report 2015-1, pp. 1–12. Sacco, D., Ferbey, T., and Jackaman, W., 2014a. Basal till potential of the Clusko River map area (NTS 093C/09), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-06; Geoscience BC Map 2014-06-01, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014b. Basal till potential of the Clisbako River map area (NTS 093B/12), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-07; Geoscience BC Map 2014-06-02, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014c. Basal till potential of the Marmot Lake map area (NTS 093B/13), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-08; Geoscience BC Map 2014-06-03, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014d. Basal till potential of the Toil Mountain map area (NTS

093C/16), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia

Geological Survey Open File 2014-09; Geoscience BC Map 2014-06-04, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014e. Basal till potential of the Suscha Creek map area (NTS 093F/01), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-10; Geoscience BC Map 2014-06-05, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014f. Basal till potential of the Coglistiko River map area (NTS 093G/04), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-11; Geoscience BC Map 2014-06-06, 1:50,000 scale.

Sacco, D., Ferbey, T., and Jackaman, W., 2014g. Basal till potential of the Pelican Lake map area (NTS

093G/05), British Columbia. British Columbia Ministry of Energy and Mines, British Colombia Geological Survey Open File 2014-12; Geoscience BC Map 2014-06-07, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014h. Basal till potential of the Euchiniko River map area (NTS 093F/08), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-13; Geoscience BC Map 2014-06-08, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014i. Basal till potential of the Chilako River map area (NTS 093G/12), British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-14; Geoscience BC Map 2014-06-09, 1:50,000 scale. Sacco, D., Ferbey, T., and Jackaman, W., 2014j. Basal till potential of the Hulatt map area (NTS 093G/13) British Columbia. British Columbia Ministry of Energy and Mines, British Columbia Geological Survey Open File 2014-15; Geoscience BC Map 2014-06-10, 1:50,000 scale. Sacco, D.A., Jackaman, W., and Ferbey, T., 2014k. Targeted geochemical and mineralogical surveys in


Summary of Activities 2013, Geoscience BC, Report 2014-1, pp. 19–34 Shilts, W., 1993. Geological Survey of Canada's contributions to understanding the composition of glacial sediments. Canadian Journal of Earth Sciences, 30, p. 333–353. Simandl, G.J., Hora, Z.D., and Lefebure, D.V. (Eds.), 1999. Selected British Columbia mineral deposit profiles, volume 3 – industrial minerals and gemstones. British Columbia Ministry of Energy, Mines, and Petroleum Resources, British Columbia Geological Survey Open File 1999-10,

the TREK Project area, central British Columbia (parts of NTS 093B, C, F, G). In: Geoscience BC

Souther, J.G., and Souther, M.E.K., 1994. Geology, Ilgachuz Range and adjacent parts of the Interior Plateau, British Columbia, Geological Survey of Canada, Map 1845A, 1:50.000 scale. Spirito, W., McClenaghan, M.B., Plouffe, A., McMartin, I., Campbell, J.E., Paulen, R.C., Garrett, R.G., and Hall, G.E.M., 2011. Till sampling and analytical protocols for GEM projects: from field to archive. Geological Survey of Canada, Open File 6850, 83 p. Tipper, H.W., 1969. Geology, Anahim Lake, British Columbia. Geological Survey of Canada, Map 1202A, 1:253,440 scale.

Tipper, H.W., 1971. Surficial geology, Anahim Lake. Geological Survey of Canada, Map 1289A,





Sacco, D., Arnold, H., Ferbey, T., and Jackaman, W., 2017. Basal till potential of the Christensen Creek map area (NTS 093C/11), British Columbia. British Columbia Ministry of Energy, Mines and Petroleum Resources, British Columbia Geological Survey Open File 2017-05, Geoscience BC Map 2017-02-04, 1:50,000 scale.