



**BRITISH
COLUMBIA**

Ministry of Energy and Mines
Resource Development and
Geoscience Branch

Coalbed Gas Potential in British Columbia

***BRITISH COLUMBIA,
CANADA***

Petroleum Geology Paper 2004-1



**BRITISH
COLUMBIA**

Ministry of Energy and Mines
Oil and Gas Division
Resource Development and Geoscience Branch

Parts of this publication may be quoted or reproduced if source is acknowledged.

This publication is also available as digital files in Adobe Acrobat PDF format from the BC Ministry of Energy and Mines internet website at:

<http://www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/studies.htm>

PETROLEUM GEOLOGY PAPER 2004 - 1

Coalbed Gas Potential in British Columbia

This publication was researched and compiled by Total Earth Science Services with expert guidance and in coordination with Barry Ryan and Mary Coward of the British Columbia Ministry of Energy and Mines.

VICTORIA
BRITISH COLUMBIA
CANADA

JUNE 2004

TABLE OF CONTENTS

| | |
|---|----|
| INTRODUCTION..... | 1 |
| LOCATION MAP..... | 2 |
| ROCKY MOUNTAIN BELT COALFIELDS OR UNITS | |
| Peace River..... | 3 |
| Gates..... | 6 |
| Gething..... | 7 |
| Minnes..... | 7 |
| East Kootenay..... | 11 |
| Elk Valley..... | 12 |
| Crowsnest..... | 14 |
| Flathead..... | 16 |
| INSULAR BELT COALFIELDS | |
| Vancouver Island..... | 19 |
| Nanaimo..... | 20 |
| Comox..... | 23 |
| Suquash..... | 27 |
| Graham Island..... | 27 |
| INTERMONTANE BELT COALFIELDS | |
| Pre-Tertiary | |
| Klappan and Groundhog..... | 31 |
| Telkwa..... | 35 |
| Tertiary | |
| Hat Creek..... | 39 |
| Princeton..... | 43 |
| Tuya River..... | 47 |
| Tulameen..... | 51 |
| Merritt..... | 55 |
| Bowron River..... | 59 |
| Coal River..... | 63 |
| APPENDIX I | |
| References..... | 67 |

INTRODUCTION

This publication provides a brief overview of the major coalfields in British Columbia and their coalbed gas potential (coalbed gas (CBG) is often referred to as coalbed methane (CBM)). Each coalfield is summarized and illustrated with simple figures, images and tables.

At present there are no producing coalbed gas wells in British Columbia but within western Canada and especially British Columbia there is tremendous incentive and opportunity to develop coalbed gas resources.

The major coalfields of British Columbia are grouped into Rocky Mountain, Insular and Intermontane belts. The Insular Belt includes the coalfields and deposits on Vancouver Island and small deposits on the Queen Charlotte Islands to the north. The Intermontane Belt includes a number of coalfields and deposits through the centre of the province and the Rocky Mountain Belt includes the important coalfields in the southeast and northeast of the province, where most of the mines are located. About 80 per cent of the coal resource in the province is contained in the Upper Jurassic to Lower Cretaceous coalfields in the Rocky Mountain Belt.

British Columbia has an estimated coal resource in the range of 250 billion tonnes to a depth of 2000 metres that is available for coalbed gas exploration. The potential coalbed gas resource in this coal is about 90 Tcf but the percentage of this resource that may eventually be recoverable will be much less.

Published coalbed gas resource assessments for the East Kootenay coalfields in the southeast corner of the province indicate potential for more than 14 Tcf and for northwest British Columbia 8 Tcf (Ryan and Dawson, 1993). Ryan (2003) estimates a resource of about 60 Tcf for the Peace River Coalfield in the northeast and 0.3 to 1.6 Tcf. for Vancouver Island coalfields. If gas contents are 1 to 3 cm³/g at Hat Creek, there could be a sizable coalbed gas resource of over 0.5 Tcf concentrated in a small area.

With a well developed network of gas pipelines, British Columbia affords exceptional opportunities to those interested in the development of coalbed gas.

If you would like to know more about opportunities for coalbed gas in British Columbia, visit the Ministry of Energy and Mines website at:

www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/cbg.htm


We encourage you to contact us for additional data required to initiate your exploration program in British Columbia.

Coalbed Gas Team


Phone: (250) 952-0408

E-mail: CBGInfo@gems9.gov.bc.ca

Coalfields and Coalbed Gas Potential in British Columbia




Peace River Coalfield



The northern Inner Foothills Belt, called the Peace River Coalfield, extends 400 kilometres from the Kakwa to the Sikanni rivers in northeastern British Columbia. The coalfield contains a number of coal-bearing units that dip to the east into the Western Canadian Sedimentary Basin. In order of decreasing age these are the Jura-Cretaceous Minnes Group and Lower Cretaceous Gething and Gates formations.

The coal resource in the Peace River Coalfield to a depth of 2000 metres is estimated to be more than 160 billion tonnes of medium and low-volatile bituminous coal. This is mainly divided between the Gates (10 billion tonnes plus) and Gething formations (120 billion tonnes plus). To date, two major mines (Quintette and Bullmoose) were developed in the Gates but have recently closed.

Coalbed gas potential in the area is largely restricted to the Gething and the Gates formations but potential also exists in the Minnes Group and Upper Cretaceous strata. A conservative estimate of gas content in the coalfield is 60 Tcf. Pipeline infrastructure is already well-developed in this area, the heartland of province's oil and gas industry.



Map of Peace River Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Coal


- The lower Gething Formation contains coal over an extensive area, though the best development is in the area between the Sukunka and Pine rivers. The formation is enclosed by the underlying Cadomin conglomerate and the overlying Bluesky conglomerate, above which is the marine Moosebar Formation. This formation is overlain by the coal-bearing Gates Formation, which contains coal from the Sukunka River southeast to the Alberta border. The deformed belt of the coalfield (Inner Foothills), which trends northwest, is defined by the outcrop of the Gething Formation on the west and a number of major thrusts on the east; the main one being the Gwillam Lake Thrust. East of the thrusts, Cretaceous beds dip into the trough of the Western Canadian Sedimentary Basin and are in places too deep to be of interest for coalbed gas development. Fold style is generally chevron with well-developed flat limbs and shorter steep dipping limbs. Regional thrusts are west dipping, though at least at Willow Creek reverse faults and axial planes dip steeply to the east.

| PERIOD | Rocky Mountains and Foothills North | | | |
|------------------|-------------------------------------|--|-------------------|-------------------|
| | NORTHEASTERN BC | | | |
| TERTIARY | | | | |
| UPPER CRETACEOUS | | | WAPITI GP. | |
| | | | SMOKY GP. | |
| LOWER CRETACEOUS | | | DUNVEGAN FM. | |
| | | | FORT ST. JOHN GP. | SHAFTESBURY FM. |
| | | | | BOULDER CREEK FM. |
| | | | | HULCROSS FM. |
| | | | | GATES FM. |
| | | | | Torrens Fm. |
| | | | | MOOSEBAR FM. |
| BULL-HEAD GP. | GETHING FM. | | | |
| | CADOMIN FM. | | | |
| JURASSIC | MINNES GP. | | | |
| | FERNIE GP. | | | |

(Smith, 1989)

Gates

- The Gates Formation contains coal from the Sukunka River south to the Saxon property near the Alberta border, although thicknesses appear to thin to the east at depth in the equivalent formation (Falher Formation).
- The formation is mainly medium-volatile bituminous but ranks on the western margin of the coalfield are lower.
- The formation is up to 280 metres thick and contains four or five laterally extensive seams ranging in thickness from 5 to 10 metres that have an aggregate thickness of up to 46 metres.
- Two major mines (Quintette and Bullmoose) in the formation were established by the early 1980s and several other properties have reached advanced stages of exploration. Quintette closed in 2000 and Bullmoose in 2003. Western Canadian Coal Corporation is applying for a mining lease on a property near the Quintette Mine called Perry Creek.



| Shipped Coal Gates Formation (Bullmoose Mine) | |
|--|---------|
| Moisture (shipped) (%) | 8.0 |
| Ash (db) (%) | 8.5 |
| Volatile matter (db) (%)..... | 26.6 |
| Fixed carbon (db) (%)..... | 56.9 |
| Sulphur (db) (%)..... | 0.4 |
| Btu/lb | 13,800 |
| Mj/kg (adb net) | 30.18 |
| FSI..... | 5.5 - 7 |
| Hardgrove Index..... | 70 - 80 |
| Rmax (%) (average) | 1.1 |
| (Ryan, 2002) | |

Gething



- The Gething Formation is up to 1036 metres thick and contains more than 100 coal beds ranging in thickness from a few centimetres to 4.3 metres. The formation underlies a large area of the Peace River Coalfield.
- The best coal development is between Williston Lake and Sukunka River to the south though the formation also contains coal at depth to the east towards Dawson Creek. South of Sukunka River the formation is thin and contains only a few seams.
- Coal rank is generally medium-volatile bituminous but decreases to the north and northeast toward Williston Lake. West and north of Sukunka River, rank increases to low-volatile bituminous.
- There has been limited mining in the Gething Formation despite the fact that it hosts the first coal to be discovered in British Columbia in 1793. More recently the Pine Valley Mining Corporation has obtained permits for a mine in the Willow Creek area.

Minnes

- Strata of the Minnes Group occupy a large area of surface bedrock of the Foothills Belt and is equivalent to the Nikanassin and Mist Mountain formations to the south.


Coalbed Gas

- The coalbed gas potential for the Peace River Coalfield is enormous and has variously been calculated to be between 60 and 200 Tcf. This is an estimate based on coal quantities and preliminary data and should be treated with caution.
- Existing coalbed gas exploration in the Peace River Coalfield falls into a number of phases:
 - ▶ Estimates were made of desorbed gas contents of samples collected from shallow holes during coal exploration from 1975 to 1985. These data were



collected as part of the appraisal of the underground coal mining potential of a number of properties. The data generally indicate low gas contents, but the data are old and were collected at shallow depths and may have little relevance to coalbed gas resource evaluation.

- ▶ In 1996, Phillips Petroleum drilled four holes in the Monkman area (Gates Formation) and the data are summarized in Dawson et al. (2000). The holes intersected cumulative coal thicknesses averaging 20 metres at depths ranging from 1200 to 1550 metres. Gas contents ranged from 6 to 26 cm³/g on an as received basis. Comparisons of desorption and adsorption data indicate that the seams are close to saturated. Low permeabilities were encountered. Four adsorption isotherms from samples from one of the Phillips wells gave Langmuir volumes ranging from 19.8 to 25 cm³/g on an as-received basis.



- ▶ Adsorption work done on samples from Gates coals (Lamberson and Bustin, 1992) and Gething Formation (Ryan and Lane, 2002) confirmed the high adsorption capacity of coals from these formations and documents the changes in capacity related to varying maceral compositions.
- ▶ A number of exploration holes have been drilled in the last few years. Most of the data from these holes are still confidential but will start to become public by 2004.
- Potential for coalbed gas exists in the older Minnes Group but little research has been done on its coal measures.

Key References (see Appendix I, References)

Dawson, F.M., Marchioni, D.L., Anderson, T.C. and McDougall, W.J. (2000): An Assessment of Coalbed Methane Exploration Projects in Canada; *Geological Survey of Canada*, Bulletin 549.

Lamberson, M.N. and Bustin, R.M. (1992): Coalbed Methane Characteristics of the Gates Formation Lithotypes, Northeastern British Columbia; *Canadian Coal and Coal Bed Methane Geoscience Forum*, Proceedings, pages 275-284.

Legun, A.S. (2002): Overview of Coalbed Methane Geology in Northeast British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 247-256.

Ryan, B. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website, http://www.em.gov.bc.ca/Mining/Geosurv/coal/coalinbc/coal_bc.htm.

Ryan, B.D. and Lane, B. (2002): Adsorption Characteristics of Coals with Special References to the Gething Formation, Northeast British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 331-346.

Ryan, B. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, Nov. 2003, pages 32-40.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4.

Ziegler, W.H. and Pocock, S.A.J. (1960): The Minnes Formation; *Edmonton Geological Society*, Second Annual Field Trip Guide Book, Rock Lake, pages 43-67.

East Kootenay Coalfields



The southeast corner of the province, often referred to as the East Kootenay region, contains three separate coalfields known respectively as Crowsnest, Elk Valley and Flathead. Since 1898, these coalfields have produced over 500 million tonnes of mainly coking coal and presently host five active mines.

Coal occurs in the Mist Mountain Formation of the Jurassic-Cretaceous Kootenay Group. The formation averages 500 to 600 metres in thickness and contains from 4 to 30 plus seams; cumulative coal thickness ranges up to over 60 metres. Seams range in rank from high to low-volatile bituminous.

Coalbed gas resource assessments for the East Kootenay coalfields indicate potential for more than 14 Tcf.


Elk Valley Coalfield


Coal

- The coal resource to a depth of 1500 metres in the Elk Valley Coalfield is estimated to be 19 billion tonnes (Johnson and Smith, 1991).
- There are three mines in the Elk Valley Coalfield. In the south, the Line Creek mine produces medium-volatile hard coking coal and lesser amounts of thermal coal. In the northern part of the coalfield, the Greenhills and Fording River mines produce medium and high-volatile coking coal from a large number of seams through a thick Mist Mountain Formation section.
- The coalfield contains two north-trending synclines separated by a major normal fault. The Bourgeau Thrust defines the west edge of the coalfield.
- Seams are folded and thrust faulted in both synclines. Thicker seams are developed low in the section but tend to host major thrusts, consequently some are extensively sheared.

Coalbed Gas

- There is an estimated coalbed potential gas resource of 7.7 Tcf in the Elk Valley Coalfield (Johnson and Smith, 1991).





Map of Elk Valley Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm


- In 1981, CANMET desorbed samples from three holes over a depth range of 0 to 400 metres (Feng et al., 1981). They found that seams above 200 metres contained less than 2 cm³/g of gas but below 200 metres gas contents ranged from 5 to 11 cm³/g (as-received basis).
- In 1991-92, Norcen drilled four holes and a limited production test well in the northern part of the Elk Valley Coalfield. The holes were located on the east limb of a syncline where the Mist Mountain Formation is 650 metres thick and contains nineteen seams with a cumulative thickness of 54 metres. Gas contents averaged 13.25 cm³/g (air-dried basis) for coals deeper than 250 metres. There were indications of moderate CO₂ contents in the gas desorbed from some samples.
- Suncor drilled a single hole in the core of the Alexander Creek Syncline south of the Greenhills mine in 1998. Some of the seams had high CO₂ concentrations and the company did not follow up on the exploration.
- Fording Coal Limited drilled a single hole on the mine site in 1993. This 533-metre deep hole intersected 44 cumulative metres of coal between 300 to 520 metres depth. Gas contents ranged from about 3 to 12.6 cm³/g on an air-dried basis, ash varied from 4 to 60 per cent. Generally, measurements indicated permeability in the range of 1 to 6 mD for the seams (Dawson et al., 2000).
- Since 2001, Encana has drilled seventeen holes in the Elk Valley Coalfield and is presently pumping on two pilots. Data will start to become public in 2004.

Crowsnest Coalfield

Coal

- The Crowsnest Coalfield has a coal resource of over 25 billion tonnes and includes the Elkview and Coal Mountain mines (Ryan, 2003). The coalfield extends from the town of Sparwood to 20 kilometres south of the town of Fernie. This 600 square kilometre area is underlain by the Jura-Cretaceous Mist Mountain Formation, which contains from 30 to 60 cumulative metres of high to low-volatile bituminous coal.

| Product coal dry basis | Medium- Volatile | High- Volatile |
|---------------------------|-----------------------|-------------------|
| Volatile matter (%)..... | 21 - 28 | 32 |
| Fixed carbon (%)..... | 64 - 69 | 62 |
| Ash (%) | 8 - 9.8 | 6 |
| Sulfur (%) | 0.3 - 0.7 | 0.4 - 0.8 |
| Btu/lb..... | 13,750 - 14,200 | 14,200 - 15,050 |
| Mj/kg | 32 - 33 | 33 - 35 |
| Hardgrove Index..... | > 80 | >60 |
| Rmax% | 1.1 - 1.35 | 0.8 - 1.1 |



Map of Crowsnest Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

- The structure is a large basin cored by the younger Elk Formation and rimmed by the older Mist Mountain Formation. Coal rank varies around the perimeter of the coalfield and downdip into the basin.
- The Coal Mountain mine, located in an eastern outlier of the coalfield, produces high-volatile A bituminous weak coking coal; most of the coal comes from a single seam near the base of the Mist Mountain Formation. The Elkview mine, located in the northern end of the coalfield, produces medium-volatile hard coking coal, mainly from the bottom four seams in the formation.


Coalbed Gas

- The Crowsnest Coalfield has a potential coalbed gas resource of over 6 Tcf (Ryan, 2003).
- In 1990-91, three companies drilled a total of seven holes in the field. These holes, drilled in the southern part of the coalfield, were drilled to test coal thicknesses and gas contents and did not test for production.
- Mobil/Chevron drilled two holes, one of which penetrated 491 metres of Mist Mountain Formation and intersected seven major coal zones with a cumulative coal thickness of 54 metres. Dawson et al. (2000) reported that the coal is fragmented or sheared. Gas contents ranged from 1.36 to 16.56 cm³/g. Based on comparison of measured gas contents on a dry ash free basis to adsorption isotherm results, samples were saturated to between 25 and 63 per cent of maximum gas capacity. Permeability appeared to be low to moderate.
- Gulf Canada drilled two holes. Results were not encouraging mainly because the holes were drilled too close to coal subcrop. Saskoil drilled three holes. Two holes when combined intersected about 500 metres of the Mist Mountain Formation with about 63 metres of coal. The coal was moderately gassy with gas contents ranging from 0.4 to 11.9 cm³/g (air-dried basis).
- Mobil/Chevron drilled three holes in 2003/2004. Data will remain confidential for 3 years.

Flathead Coalfield

Coal

- Coal occurs in a number of outliers of the Mist Mountain Formation, the largest of which is Sage Creek near the US border.
- The Mist Mountain Formation ranges from 198 to 259 metres in thickness; coal occurs in five horizons having an aggregate thickness of approximately 30 metres.



- The coal resource available for coalbed gas exploration in the Flathead Coalfield is about 1 billion tonnes (Ryan, 2003).

Coalbed Gas

- The coal resource available for coalbed gas exploration is about 1 billion tonnes with a potential coalbed gas resource of 0.4 Tcf (Ryan, 2003).

Key References (see Appendix I, References)

Dawson, F.M., Marchioni, D.L., Anderson, T.C. and McDougall, W.J. (2000): An Assessment of Coalbed Methane Exploration Projects in Canada; *Geological Survey of Canada*, Bulletin 549.

Feng, K.K., Cheng, K.C. and Augsten, R. (1981): Methane Desorption of Fording Coal from Greenhills Multiple Seams; Division Report, *CANMET*, ERP/MRL 81-67(J).

Johnson, D.G.S. and Smith, L.A. (1991): Coalbed Methane in Southeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Petroleum Geology, Special Paper 1991-1.


Pearson, D.E. and Grieve, D.A. (1985): Rank Variation, Coalification Pattern and Coal Quality in the Crowsnest Coalfield, British Columbia; *Canadian Institute of Mining and Metallurgy Bulletin*, September, pages 39-36.

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/Mining/Geolsurv/coal/coalinbc/coal_bc.htm

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.


Ryan, B.D. (2004): The CBM Resource of Some Prospective Areas of the Crowsnest Coalfield; *B.C. Ministry of Energy and Mines*, in press, Oil and Gas Emerging Opportunities and Geosciences Branch.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.



Stratigraphic column - Mist Mountain Formation, Greenhills mine, Elk Valley Coalfield (Dawson, 1995).

Vancouver Island Coalfields



Along the coast of British Columbia coal deposits occur on Vancouver Island, the Queen Charlotte Islands to the north, and some of the Gulf Islands. Coal rank ranges from anthracite to lignite, with most being high-volatile bituminous. Host rocks range from Jurassic to Tertiary though most coal is found in the Upper Cretaceous Nanaimo Group. The major Upper Cretaceous coalfields on Vancouver Island are the Comox and Nanaimo fields. The Comox Coalfield includes the Quinsam underground mine, which is the only operating coal mine on the island. There are at least two coal-bearing formations within the Nanaimo Group. The first is the basal Comox Formation which is the source of coal in the Comox Coalfield, and the Extension and Protection formations in the Nanaimo Coalfield.

Estimates of the coal resources on Vancouver Island range from a low of 800 million tonnes to a high of 6920 million tonnes for an average of 3860 million tonnes.

The coalbed gas resource for Vancouver Island is estimated to be 0.3 to 1.6 Tcf. A gas pipeline passing through the Comox and Nanaimo coalfields enhance the development potential. There are a number of smaller coalfields with limited coalbed gas potential - these are the Suquash, Cowichan, Alberni, Qualicum and Quatsino coalfields. North of Vancouver Island, coal-bearing rocks on the Queen Charlotte Islands of Jura-Cretaceous and Tertiary ages have low coalbed gas potential.

Nanaimo Coalfield

Coal

- The coalfield, which covers about 400 square kilometres, produced over 50 million tonnes of high-volatile A bituminous coal in the 100 years preceding 1953.

The producing area, in which the Douglas, Newcastle, and Wellington seams were mined, covered about 170 square kilometres. Air-dried basis analyses for the three seams is reported below.

| | Douglas | Wellington | Newcastle |
|--------------------------|-------------|------------|-----------|
| Moisture (%)..... | 1.6 | 1.9 | 1.6 |
| Volatile matter (%)..... | 43.2 | 39.4 | 39.7 |
| Fixed carbon (%)..... | 45.5 | 45.7 | 47.7 |
| Ash (%) | 9.2 | 11.7 | 10.1 |
| Sulphur (%) | 1.2 | 1.3 | 0.9 |
| Btu/lb | 13,160 | 12,470 | 12,830 |
| Mj/kg..... | 30.61 | 29.01 | 29.84 |
| Hardgrove Index..... | 67 | 67 | 67 |
| FSI | ND | 3 - 4 | ND |
| Rmax% all seams | 0.64 - 0.72 | | |


(Ryan, 2002)

- Coal occurs in the Upper Cretaceous Extension and Protection formations, in three non-overlapping seams called the Douglas, Wellington and Newcastle. Remaining coal resources are estimated to range from 10 to 100 million tonnes (Ryan, 2003).
- Seams are broken by numerous normal faults which disrupt an easterly regional dip.


Coalbed Gas

- Coalbed gas opportunities in the Nanaimo Coalfield are probably limited because of past underground mining of the Douglas and Wellington seams.
- Recent work indicates there is also coal in the Comox Formation within the Nanaimo Coalfield and there may be some coalbed gas potential in this deeper formation. Cumulative coal thickness in the Douglas coal zone ranges from 3 to 10 metres - a single intersection in the Comox Formation was 3 metres thick. Most of the coalbed gas potential will be in the eastward extension of the Douglas Seam and the deeper Comox coals if they are extensive.
- A gas pipeline passes through the coalfield and enhances development potential.
- The potential coalbed gas resource of the Nanaimo Coalfield is estimated to be in the range of 0.4 Tcf (Ryan, 2003).

Douglas Coal Zone, Nanaimo Coalfield



Representative stratigraphic column (Cathyl-Bickford, Wilson and Hoffman, 1992).



Map of Nanaimo Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Comox Coalfield

Coal

- The coalfield covers an area of about 1100 square kilometres and produced 18 million tonnes of high-volatile A and B bituminous coal from 1888 to 1953.

| | |
|--------------------------|------------|
| Moisture (%)..... | 3.0 |
| Volatile matter (%)..... | 36.5 |
| Fixed carbon (%)..... | 47 |
| Ash (%) | 13.5 |
| Sulphur (%) | 0.25 |
| Btu/lb | 11,880 |
| Mj/kg..... | 27.6 |
| Hardgrove Index..... | 48 |
| FSI | low |
| Rmax% | 0.6 - 0.88 |


(Ryan, 2002)

- Coal is found in the lowermost Comox Formation, which dips eastward with the regional dip complicated by broad folds, thrusts and normal faults; generally deformation is more intense along the western margin of the coalfield.
- The Comox Formation around the town of Courtenay contains four major seams with a cumulative thickness ranging from 2 to 8 metres.
- The Comox Coalfield includes the Quinsam underground mine, which is the only operating coal mine on the island.


Coalbed Gas

- The coalfield is divided into a number of sub-areas for the purposes of estimating coalbed gas resources. In the north, some desorption data was collected from holes drilled on the Quinsam mine site where the rank is high-volatile B bituminous and cumulative coal thickness is about 7.5 metres. Ryan and Dawson (1994) collected seven samples from two shallow holes - gas content of samples collected from depths ranging from 100 to 150 metres ranged from 0.44 to 1.632 cm³/g on a dry ash free basis. The sub-area covers an area of 155 square kilometres and is generally fault-bounded. The area of the coalfield around Campbell River is about 300 square kilometres and the potential coalbed gas resource is about 130 Bcf (Ryan, 2003).
- South of Campbell River and north of Courtenay, the coalfield covers an area of about 250 square kilometres, and has a potential coalbed gas resource of about 130 Bcf (Ryan, 2003). There has not been much surface exploration or drilling in this area and there is little information on coal thicknesses.

North End of Comox Coalfield




Representative stratigraphic column (Ronaghan, 1980).



Map of Comox Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm



- The area of the Comox Coalfield around the town of Courtenay covers about 200 square kilometres. There is a history of mining in the area and gas emission data from the mines indicates gas contents ranging from 7.8 to 11.7 cm³/g on a dry ash free basis at depths that average 250 metres. The Comox Formation contains four major seams with a cumulative thickness ranging from 2 to 8 metres. Three coalbed gas exploration holes and a natural gas well were drilled in the Courtenay area in 2000 and 2001. Twelve samples for desorption were collected from the three coalbed gas holes and the results indicate gas contents ranging from 2.2 to 7.4 cm³/g on an as-received basis (or 3.6 to 12 cm³/g on a dry ash free basis) for samples collected over a depth range of 238 to 255 metres (Ryan, 2002).
- The area of the Comox Coalfield south of Courtenay covers about 330 square kilometres of which about 160 square kilometres is within the Tsable River area. Coal in the area, which has a rank of high-volatile A bituminous, is in the Comox Formation and cumulative coal thickness is about 8.3 metres contained in up to four seams. Gas was reported in a few deep holes drilled in the early 1900s. In 1996, 13 samples were desorbed covering depths from 126 to 376 metres. Gas contents on an as-received basis range from 1.6 to 5.5 m³/tonne (Ryan, 1997). Gas contents increase consistently with depth but samples were under saturated. These concentrations are high enough to make the area attractive for its potential coalbed gas resource. Resource calculations for part of the southern end of the Comox Coalfield indicate a potential of about 0.45 Tcf (Ryan, 2003). A gas pipeline passes through the coalfield and enhances the development potential.

Suquash Coalfield

Coal

- The coalfield covers an area of 120 square kilometres of Nanaimo Group rocks.
- The coal-bearing section is at least 360 metres thick; coal is in the upper 200 metres of the section in approximately nine zones that contain about four metres of coal in total. Coal rank is high-volatile B to A bituminous.

| Quality On An Air-Dried Basis | |
|-------------------------------|-------------|
| Moisture (%)..... | 6 |
| Volatile matter (%)..... | 23 |
| Fixed carbon (%)..... | 25.4 |
| Ash (%) | 45.3 |
| Sulphur (%) | 2.2 |
| Btu/lb | 11,580 |
| Mj/kg..... | 26.94 |
| Hardgrove Index..... | 43 |
| FSI..... | 1.5 |
| Rmax% | 0.63 - 0.81 |

(Ryan, 2002)

- If the coalfield is underlain by, on average, two metres of coal, then the coal resource would be about 300 million tonnes (Ryan, 2003). If the coal-bearing stratigraphy extends to the northeast under Malcolm Island, it provides an area of 300 square kilometres and a potentially larger coal resource.


Coalbed Gas

- If a coal resource of 300 million tonnes is assumed then the estimated potential coalbed gas resource is 45 Bcf (Ryan, 2003).

Graham Island Coalfield

Coal

- Coal deposits on the Queen Charlotte Islands have not been fully mapped because of thick vegetation, lack of outcrop and complex geology. All the deposits are on the larger northern Graham Island and are either Tertiary or Jura-Cretaceous.
- Preliminary exploration by a number of companies in recent years has failed to identify sufficient resources to justify continued development.
- Tertiary lignites are exposed in the northeast coastal areas and the older anthracites and bituminous deposits outcrop in the southwestern part of the island.



Representative stratigraphic column, Suquash Coalfield (Gardner, 1984).

The variation in rank of the older deposits is attributed to the presence of younger volcanic rocks in the area.

| | Lignite | HV Bituminous | Anthracite |
|--------------------------|-----------|---------------|------------|
| Moisture (%)..... | 18.5..... | 1.2..... | 1.8..... |
| Volatile matter (%)..... | 44.2..... | 36.2..... | 4.7..... |
| Fixed carbon (%)..... | 34.7..... | 46.5..... | 85.7..... |
| Ash (%)..... | 2.3..... | 16.1..... | 6.6..... |
| Sulphur (%)..... | 0.3..... | 1.0..... | 0.9..... |
| Btu/lb..... | ND..... | 11,235..... | |
| Mj/kg..... | ND..... | 26.13..... | |

(Ryan, 2002)

Coalbed Gas

- The coal-bearing rocks generally have low coalbed gas potential.

Key References (see Appendix I, References)

Cathyl-Bickford, C.G. (1991): Coal Geology and Coal bed Methane Potential of Comox and Nanaimo Coalfields, Vancouver Island, British Columbia; *Rocky Mountain Association of Geologists*, pages 155-162.

Cathyl-Bickford, C.G., Wilson, R. and Hoffman, G.L. (1992): Geology, Mineability and Coalbed Gas Potential of the Douglas Coal Zone in the Nanaimo Coalfield, Vancouver Island; *Canadian Coal and Coal Bed Methane Geoscience Forum*, Proceedings, pages 293-309.

Dawson, F.M. (1995): Coalbed Methane: A Comparison Between Canada and the United States; *Geological Survey of Canada*, Bulletin 489.

Energy Marketing Update (1992): What is Coal Bed Methane; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Energy Market Update, Vol.4, No.1.

Gardner, S.L. (1984): Geological Reconnaissance of Vancouver Island Coal Areas - Suquash Coal Basin and Outliers, Northern Vancouver Island, British Columbia; Coal Assessment Report 778, *B.C. Ministry of Energy and Mines*.

Proudlock, P. (1990): An Overview of Albertan and Canadian Demethanation Experience, Technology Advances and Development Potential; in Coalbed Methane in Alberta - What's it all about; *Alberta Geological Survey*, Information Series No.108.

Ronaghan, R.J. (1980): Geology of the Coal Reserve in the Quinsam Mining Block; Coal Assessment Report 821, *B.C. Ministry of Energy and Mines*.

Ryan, B.D. and Dawson, F.M. (1994): Coalbed Methane Canister Desorption Techniques; in Geological Fieldwork 1993, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-1, pages 45-256.

Ryan, B.D. (1997): Coalbed Methane in the Comox Formation, Tsable River Area, Vancouver Island; in Geological Fieldwork 1996, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1998-1, pages 353-363.


Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/Mining/Geosurv/coal/coalinbc/coal_bc.htm

Ryan, B.D. (2002): Note on Desorption Results of Comox Formation Coals from the Courtenay Area, Vancouver Island, British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 319-330.

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.


Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.

Klappan and Groundhog Coalfields



Anthracite coal deposits of the Klappan and Groundhog coalfields occur within the Bowser Basin in a remote region of rugged mountainous terrain in northwestern British Columbia. The basin, which encompasses an area of approximately 50,000 square kilometres, is defined by the outcrop extent of up to 3500 metres of sediments of the Bowser Lake Group of Middle Jurassic to Lower Cretaceous age. The coal is contained in the Jura-Cretaceous Currier Formation which is up to 1100 metres thick and contains up to 25 seams, ranging in thickness up to 7 metres. The area is extensively folded with the regional structure dominated by a northwest trending synclinorium.

The Klappan Coalfield which is in the northern part of the Bowser Basin, was extensively explored in the period 1982 to 1986 with the hope of developing a major anthracite export mine. An estimated 37 million tonnes of anthracite is contained in the basin, primarily in the Klappan Coalfield. Though no coalbed gas wells have been drilled in the Bowser Basin, a potential of 8 Tcf of contained coalbed gas is conservatively estimated.



Map of Klappan-Groudhog Coalfields

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm


Coal

- Coal in the northern 10 per cent of the Bowser Basin is found in the Upper Jurassic to Lower Cretaceous Currier Formation in the Klappan Coalfield and in its equivalent in the south, the Prudential Formation in the Groundhog Coalfield.
- The Bowser sediments are extensively folded with the regional structure which is controlled by a prominent northwest trending synclinorium (the Biernes Synclinorium).
- Potentially economic coal seams up to 7 metres thick are present in the lower third of the Currier Formation. In the Mount Klappan area the formation is 900 to 950 metres thick and contains a cumulative coal thickness that ranges up to 54 metres.
- Coal-bearing rocks underlie approximately 2000 square kilometres around the synclinorium.
- The coal section is often folded into asymmetric chevron folds. The gently dipping limbs of these folds often represent undeformed structural domains on the order of 1 square kilometre.
- Coal rank ranges from semi-anthracite to meta-anthracite.
- The Klappan Coalfield was extensively explored in the period 1982 to 1986 by Gulf Canada Corporation with the hope of developing a major anthracite export mine. Fortune Minerals Limited acquired the property in 2002 and is reviewing development potential.
- The estimated potential coal resource in the Klappan and Groundhog coalfields is 37 billion tonnes and much of this is in the Currier Formation within the synclinorium (Ryan and Dawson, 1993).

Coalbed Gas


- An adsorption isotherm for a single sample of drill core coal indicates that the adsorption potential to a depth of about 1400 metres is high.
- The anthracitic rank coals are the best at retaining methane at shallow depth. The high rank has an important implication for improved permeability and gas recovery.
- An estimated potential coalbed gas of 8 Tcf is calculated using a conservative gas content value of 5 cm³/g (Ryan and Dawson, 1993). The resource assessment is restricted to the area underlain by the Currier Formation within the coalfields.
- Potentially economic locations for wells occur on the relatively undeformed flat-dipping limbs of asymmetric chevron folds.

Jura-Cretaceous Currier Formation (Bowser Lake Group)



| Raw | Clean Coal | Air-Dried Basis |
|---------------------|------------|-----------------|
| Moisture (%) | 2.0 | 1.0 |
| Volatile matter (%) | 8.0 | 6.5 |
| Fixed carbon (%) | 54.0 | 85.5 |
| Ash (%) | 36.0 | 7.0 |
| Sulphur (%) | 0.5 | 0.5 |
| Btu/lb | 8,800 | 13,320 |
| Mj/kg | 20.47 | 31.0 |
| Hardgrove Index | | 38 |
| Rmax (%) | 2.8 - 4.4 | |

(Ryan, 2002)



Key References (see Appendix I, References)


Dawson, F.M. (1995): Coalbed Methane: A Comparison Between Canada and the United States; *Geological Survey of Canada*, Bulletin 489.

Ryan, B.D. and Dawson, M.F. (1993): Coal and Coalbed Methane Resource Potential of the Bowser Basin, Northern British Columbia, (104H/104A); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1993-31.

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/mining/geosurv/coal/coalinbc/coal_bc.htm.


Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, Nov. 2003, pages 32-40.

Telkwa Coalfield



The Telkwa Coalfield in central British Columbia extends for about 50 kilometres along the Bulkley River from north of the town of Smithers to south of the village of Telkwa. Two coal-bearing units containing up to 21 metres of coal occur in the Lower Cretaceous Skeena Group. Thermal coal and small quantities of anthracite were mined in the coalfield in the early part of the century and more recently, near Telkwa, the coalfield has been intensively explored by a number of companies with the intention of developing an open-pit thermal-coal mine. A total surface mineable coal resource is estimated at 180 million tonnes and the potential resource available for coalbed gas is about 860 million tonnes.

A gas pipeline that connects Prince Rupert and Kitimat with pipelines from the northeast and southwest passes through the coalfield and is a significant infrastructure advantage. Desorption results indicate that coal in the coalfield retains gas at shallow depths. An estimate of the coalbed gas resource of the Telkwa Coalfield is 0.13 Tcf.




Map of Telkwa Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm


Coal

- Two coal-bearing units separated by a marine mudstone unit are contained in the Lower Cretaceous Skeena Group. The upper unit contains at least 8 seams with cumulative coal thickness up to 14 metres. The lower unit contains a single coal zone with cumulative coal thickness up to 7 metres. The seams are folded into open folds and broken by reverse and normal faults.
- Coal rank ranges from high-volatile bituminous A to anthracite, though most of the coal is in the range high-volatile A to medium-volatile bituminous.
- Close to 400,000 tonnes of thermal coal were produced from 1918 to 1980. Since 1978 there has been intensive exploration in the area culminating in plans to develop a 1 to 1.5 million tonnes per year open-pit thermal coal mine.
- The total surface mineable coal resource is estimated to be 180 million tonnes and south of Telkwa, 20 to 50 million tonnes have been identified as potentially surface mineable. The potential resource available for coalbed gas is about 860 million tonnes.
- The coalfield has historically been explored as a source of thermal coal but the wide range in rank means that there is potential for metallurgical coal.



| Estimated Shipped Quality | |
|---------------------------|-----------|
| Moisture (%) | 10 |
| Volatile matter (%) | 25.6 |
| Fixed carbon (%) | 65 |
| Ash (%) | 7.7 |
| Sulphur (%) | 1.0 |
| Btu/lb | 10,950 |
| Mj/kg | 25.6 |
| FSI | 0.0 - 5.5 |
| Hardgrove Index | 40 - 70 |
| Rmax% average | 1.0 |

(Ryan, 2002)



Coalbed Gas

- Four samples collected over a depth range of 64 to 120 metres have gas contents that range from 3.75 to 4.49 cm³/g on a dry ash-free basis. Most of the samples appear to be saturated based on the results of adsorption isotherms (Ryan and Dawson, 1994).
- Three adsorption isotherms derived from core samples yielded an average Langmuir volume of approximately 12.5 cm³/g (ash free basis) (Dawson, 1995).
- Seams have moderate to good permeability and at depths ranging from 29 to 158 metres; values range from 0.5 to 50 milli Darcies (Ryan and Dawson, 1994).
- Interburden permeability is on average greater than that of the coal.
- Proximity to the Bulkley and Telkwa rivers suggest sufficient hydrostatic pressures exist to have retained significant volumes of gas at depth.
- A coalbed gas resource estimate for the Telkwa deposit is 0.13 Tcf (Ryan, 2003).
- A gas pipeline passes through the centre of the coalfield connecting the field to local or continental markets.

Key References (see Appendix I, References)

Palsgrove, R.J. and Bustin, R.M. (1989): Stratigraphy and Sedimentology of the Lower Skeena Group, Telkwa Coalfield, Central British Columbia (93L/11); *in* Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 449-454.


Ryan, B.D. and Dawson, F.M. (1994): Potential Coal and Coalbed Methane Resource of the Telkwa Coalfield Central British Columbia (93L/11); *in* Geological Fieldwork 1993, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-1, pages 225-243.

Dawson, F.M. (1995): Coalbed Methane: A Comparison Between Canada and the United States; *Geological Survey of Canada*, Bulletin 489.

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/mining/geolsurv/coal/coalinbc/coal_bc.htm.

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Hat Creek Coalfield



The Hat Creek Coalfield is located 20 kilometres west of Cache Creek in the south-central part of the province. The coalfield contains one of the thickest Tertiary coal deposits in the world with an enormous amount of low rank coal concentrated in a small area. It consists of two poorly exposed coal deposits. The northern, Number 1 deposit covers 3.5 square kilometres and the larger Number 2 deposit, 3 kilometres to the south, covers 25 square kilometres. The Number 1 deposit has been explored as a potential open-pit mine.

The Tertiary section in the area of the deposits is 1500 metres thick and is divided into three units. The middle Hat Creek Formation is about 500 metres thick of which about 65 per cent is coal. Estimates of the coal resource in the whole coalfield exceed 10 billion tonnes.

If gas contents are 1 to 3 cm³/g, because of the large coal tonnages there could be a sizable coalbed gas resource of over 0.5 Tcf concentrated in a small area.

Coal

- The Tertiary section in the area of the deposits is 1500 metres thick and is divided into three units. The middle Hat Creek Formation is about 500 metres thick of which about 65 per cent is coal.
- The Number 1 deposit has been explored as a potential open pit mine. Reserves in this deposit to a depth of 200 metres are over 500 million tonnes of lignite A to sub-bituminous C coal. The deposit comprises two south plunging half-synclines truncated on the southeast end by northeast trending gravity faults. Dips average about 25 degrees.
- The Number 2 deposit occurs within a graben that is bounded by north trending normal faults. The resource of the Number 2 deposit to a depth of 460 metres is estimated to be over 2 billion tonnes.
- Total resource in the basin is over 10 billion tonnes.

Coalbed Gas

- The low rank of the coal means the amount of thermogenic methane generated will be small to nil and the ability of the coal to adsorb methane will also be low. However, low rank coals can have high meso-porosities compared to higher rank coals and therefore can retain moderate amounts of free gas, much of which would likely be of biogenic origin.
- Even if gas contents are only of 1 to 3 cm³/g because of the large coal tonnages (over 10 billion tonnes) there could still be a sizable coalbed gas resource at Hat Creek of over 0.5 Tcf concentrated in a small area (Ryan, 2003).

Key References (see Appendix I, References)

Bustin, R.M. (1999): Free Gas Storage in Matrix Porosity: A Potentially Significant Coalbed Resource in Low Rank Coals; in Proceedings International Coalbed Methane Symposium 1999, May 3-7th, *University of Alabama*, pages 197-214.


Church, B.N. (1977): Geology of the Hat Creek Coal Basin; in *Geology in British Columbia 1975*, B.C. Ministry of Energy, Mines and Petroleum Resources, pages 99-118.

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in *Geological Fieldwork 2002*, B.C. Ministry of Energy and Mines, Paper 2003-1, pages 213-236.


COAL MEASURES

Total thickness - 445 metres
 Net coal thickness - 286 metres




| Typical Raw As-Received Coal Quality | |
|--------------------------------------|-----------|
| Moisture (%)..... | 23.5 |
| Volatile matter (%)..... | 24.8 |
| Fixed carbon (%)..... | 25.1 |
| Ash (%) | 26.6 |
| Sulphur (%)..... | 0.2 - 0.7 |
| Btu/lb | 5800 |
| Mj/kg..... | 13.49 |
| Hardgrove Index..... | 58 |
| Rmax%..... | 0.4 |

(Ryan, 2002)



Representative stratigraphic column, Hat Creek Deposit (Suncor CBM Prospectus, 2000).


Princeton Coalfield



The Tertiary Princeton Coalfield is located in the south-central part of the province and is centred on the town of Princeton. It covers an area of about 170 square kilometres and is approximately 24 kilometres long and 4 to 7 kilometres wide. The coalfield is composed of a number of northwest trending synclines with faults defining the western and eastern edges. There were thirteen small underground mines and one on surface that operated in the central part of the coalfield up until about 1961. From 1919-40, about 2.1 million tonnes of coal were produced.

In the area south of Princeton there are four major coal zones, numbered from one at the top to four at the base of about 530 metres of the coal-bearing Allenby Formation. Most of the coal resource, which is high-volatile B bituminous in rank, occurs near the middle of the formation within the southern half of the coalfield, south of the town of Princeton. There is at least a 110 million tonne coal resource.

Based on work to date, Petrobank estimates that the potential gas-in-place resource is 250-500 Bcf.




Map of Princeton and Tulameen Coalfields

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Coal

- In the area south of Princeton there are four major coal zones, numbered from one at the top to four at the base in about 530 metres of section in the Tertiary Allenby Formation.



Generalized stratigraphic column (Dolmage and Campbell, 1975).

- The lowest zone (Number 4) ranges in thickness from 10 to 50 metres and is described as mainly dirty coal with a clean coal seam up to 2 metres thick near the top. The next (Number 3 zone) is 140 metres higher in the section and is about 30 metres thick. It is reported to contain up to two workable seams. The Number 2 zone contains 7.5 metres of coal in a 26-metre section and is 200 metres above the Number 3 zone. The Number 1 zone is 100 metres above the Number 2 zone and contains from 2 to 3 metres of coal.

- There were thirteen small underground mines and one surface mine that operated in the central part of the coalfield up until about 1961. About 2.1 million tonnes of high-volatile coal were produced prior to 1961.

| Typical Dry Basis Analyses | |
|----------------------------|---------------|
| Volatile matter (%)..... | 33.3 - 34.4 |
| Fixed carbon (%)..... | 40.5 - 42.6 |
| Ash (%) | 23 - 26.1 |
| Sulphur (%) | 0.75 - 0.83 |
| Btu/lb | 9941 - 10,336 |
| Mj/kg | 23.1 - 24.0 |
| FSI | ND |
| Hardgrove Index | 39 - 45 |
| Rmax%..... | 0.52 |

(Ryan, 2002)

- In 2002, three coal exploration holes were drilled in the Princeton Coalfield and based on encouraging results the coalbed gas tenure was obtained by Petrobank. Since then the company has conducted two seismic surveys and as of early 2004 plans to drill a multi-hole pilot.

Coalbed Gas

- Based on work to date, Petrobank estimates that the potential gas-in-place resource is 250-500 Bcf (Petrobank Energy and Resources Ltd. 2003 Annual Report).

Key References (see Appendix I, References)


Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website, www.em.gov.bc.ca/Mining/Geosurv/coal/coalinbc/coal_bc.htm.

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in *Geological Fieldwork 2002*, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.

Tuya River Coalfield




The Tuya River Coalfield is located between the communities of Dease Lake and Telegraph Creek in northwestern BC and straddles the drainage of Tuya River and its tributaries Little Tuya River and Mansfield Creek. These three drainages have incised meandering canyons up to 200 metres deep in an area of otherwise subdued relief.

The coal-bearing rocks are not younger than early Eocene and not older than Paleocene and may be equivalent to the Tango Formation of the Sustut Group. The coal zone is about 100 metres thick and contains up to 30 metres of high-volatile bituminous B coal.

The earliest recorded description of coal in the Tuya River area was in 1904 when seams up to 12 metres thick were described adjacent to Tuya River. The basin was explored in detail in the period 1979 to 1980, when 10 cored-holes were drilled and a number of hand trenches dug.

The basin covers approximately 150 square kilometres, potentially containing over 600 million tonnes of coal or about 400 million tonnes to a depth of 1600 metres. The coalbed gas resource is estimated to be 66 Bcf.




Map of Tuyá Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Coal

- The stratigraphic succession of the Tuya River coalfield contains two units of Paleocene to Eocene age, possibly equivalent to the Tango Formation of the Sustut Group. The upper unit, which is at least 300 metres thick, is composed of volcanic-pebble conglomerate, sandstones and volcanic rocks. The lower unit is 200 to 300 metres thick and contains a single coal-bearing zone. The zone is about 100 metres thick and includes from 5 to 30 metres of coal. The structure of the basin is that of an open, northerly-plunging syncline complicated by smaller scale faults and folds.
- To date, exploration consists of mapping and drilling in the western half of the basin by PetroCanada (Reid, 1980; De Nys, 1980) and mapping of the eastern half by Esso Minerals Canada (Vincent, 1979). Ten cored holes were drilled and a number of hand trenches dug.
- The coal, which has a rank of high volatile bituminous B, is hard, well-cleated and contains a moderate amount of amber.
- The basin contains about 400 million tonnes of coal within 1600 metres of surface (Ryan, 2003).
- A subsidiary basin or extension of the Tuya River Coalfield outcrops 20 kilometres southwest of Tuya River in the lower reaches of the Tahltan River. Lignite coal zones ranging up to 4 metres in thickness occur.



| Typical Average Raw Analysis (on an as-received basis) | |
|---|--------------|
| Moisture (%)..... | 12.4 |
| Volatile matter (%)..... | 30.7 |
| Fixed carbon (%)..... | 37.8 |
| Ash (%) | 19.1 |
| Sulphur (%) | 0.5 |
| Btu/lb | 7,740 |
| Mj/kg..... | 18 |
| Hardgrove Index..... | 53 |
| Rmax% | 0.6 - 0.8 |
| | (Ryan, 2002) |

Coalbed Gas

- The Tuya River Coalfield was mapped and the coalbed gas resource appraised in 1991 (Ryan 1991).
- A potential coalbed gas resource of up to 66 Bcf may exist in the basin (Ryan, 1991). Most of this resource would be biogenic methane because of the low rank.

Key References (see Appendix I, References)

De Nys, F.J.G. (1980): Thundercloud Coal Project Technical Report; PetroCanada Exploration, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 243.

Reid, J.L. (1980): Thundercloud Coal Project Technical Report, PetroCanada Exploration, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 242.


Ryan, B.D. (1991): Geology and Potential Coal and Coalbed Methane Resource of the Tuya River Coal Basin; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 419-427.

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/mining/geolsurv/coal/coalinbc/coal_bc.htm

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Vincent, B.D. (1979): Geological Mapping, Tuya River Property, British Columbia: Esso Minerals Canada; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 246.

Tulameen Coalfield



The Tulameen Coalfield is located in the south-central part of the province, 20 kilometres northwest of Princeton. It forms an elliptical sedimentary basin 5 by 4 kilometres in size and contains two well-developed thick seams of high-volatile B bituminous rank. The area underlain by coal is about 10 square kilometres. Underground mining from 1919 to 1940, produced about 2 million tonnes.

The Tertiary stratigraphy in the coalfield is assigned to the Allenby Formation and is subdivided into five units that consist of a lower conformable volcanic unit, three sedimentary units and an uppermost non-conformable volcanic unit. The middle sedimentary unit, which is about 130 metres thick, contains shales and two major coal seams. The upper sedimentary unit consists of conglomerates and sandstones and is at least 400 metres thick. The rocks are folded into a broad syncline. Based on the synclinal form of the coal horizon there is a potential coal resource of at least 300 million tonnes.

There is a potential coalbed gas resource of 50 Bcf in the coalfield.

Coal

- The coal-bearing section contains two coal seams though they tend to be very dirty with partings of shale and bentonite. The lower seam is 7 to 8 metres thick and the upper seam 15 to 21 metres thick. The seams are thought to underlie the whole coalfield.
- Underground mining carried out in the area began in 1919 and ended in 1940; the mines extracted about 2 million tonnes. Renewed interest in the 1950s led to surface mining that extracted a further 150,000 tonnes of coal. There was exploration in the coalfield in the 1970s and 1980s and a major exploration program in 1998. In 1999, a bulk sample was test-washed with favourable results.
- At present, Compliance Energy Corp. has started mining on a small scale and is trucking coal to local markets; the company has expressed interest in the coalbed gas potential.
- Based on the synclinal form of the coal horizon there is a potential coal resource of at least 300 million tonnes (Ryan, 2003).

Coalbed Gas

- A potential coalbed gas resource of 50 Bcf is estimated for the coalfield (Ryan, 2003).

Key References (see Appendix I, References)

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/Mining/Geosurv/coal/coalinbc/coal_bc.htm.


Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.


Williams, V.E. (1978): Coal Petrology of the Tulameen Coalfield, South Central British Columbia, M.Sc. Thesis, Western Washington University; in Coal Assessment Report 832, *B.C. Ministry of Energy and Mines*, 83 pp.

Tulameen Coalfield




| LEGEND | |
|-------------------|-----------------------------|
| | CHILCOTIN GROUP (basalt) |
| PRINCETON GROUP | |
| Allenby Formation | |
| | Summer Creek sandstone |
| | Vermillion Bluffs shale |
| | Hardwicke sandstone |
| | Cedar Formation (volcanics) |
| | Borehole |
| | Adit |
| | Geological contacts |
| | Coal seam |
| | Fault |
| | Syncline |

| Typical Dry Basis Analyses | |
|----------------------------|-------------|
| Volatile matter (%) | 37.4 |
| Fixed carbon (%) | 42.52 |
| Ash (%) | 20.22 |
| Sulphur (%) | 0.44 - 0.66 |
| Btu/lb | 10,828 |
| Mj/kg | 25.2 |
| FSI | ND |
| Hardgrove Index | 39 - 45 |
| Rmax% | 0.62 - 0.86 |
| (Ryan, 2002) | |




Representative stratigraphic column. The coal-bearing middle member of the Tertiary Allenby Formation is approximately 130 metres thick and consists of interbedded shales, mudstone, tuffs, minor sandstone and coal (Williams, 1978).

Merritt Coalfield



The Merritt Coalfield is located in the south-central part of the province and comprises several isolated Tertiary sedimentary areas which in part underlie the town of Merritt and adjacent areas. The coal-bearing sedimentary rocks are assigned to the Coldwater Formation. The main area, which is under and south of Merritt, covers about 80 square kilometres. A smaller area to the east is about 25 square kilometres in size and has not been extensively explored. The Merritt area, which contains high-volatile C to A bituminous coal, was explored in 1945 and 1960 and about 2.4 million tonnes of coal were produced.

There appear to be seven seams with a cumulative coal thickness of 22 metres in a 250-metre section. It is difficult to estimate the resource in the area but if a cumulative coal thickness of 10 metres extends through the whole coalfield, then there could be over 800 million tonnes and a moderate coalbed gas resource of over 50 Bcf under the town of Merritt.



Map of Merritt Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Coal

- The Merritt Coalfield contains high-volatile C to A bituminous coal. From 1906 to 1963, underground mines produced 2.4 million tonnes based on exploration that outlined about 10 million tonnes of underground mineable coal. Overall, the Merritt Coalfield has not been extensively explored.
- There appear to be seven seams with a cumulative coal thickness of 22 metres in a 250-metre section in the Tertiary Coldwater Formation.
- It is difficult to estimate the resource in the area but if a cumulative coal thickness of 10 metres extends through the whole coalfield, then there could be over 800 million tonnes (Ryan, 2003).

Coalbed Gas

- There could be a moderate coalbed gas resource of over 50 Bcf under the town of Merritt (Ryan, 2003).
- A company has acquired freehold coal rights in the area and hopes to drill coal exploration holes that will provide some information on the coalbed gas potential.

Key References (see Appendix I, References)

Matheson, A. (1992): Subsurface Thermal Coal Sampling Survey, Merritt Coal Deposits, South-Central British Columbia; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 427-432.




Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/Mining/Geosurv/coal/coalinbc/coal_bc.htm

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.

Merritt Coalfield




Typical Air-Dried Analysis

| | |
|---------------------|-------------|
| Moisture (%) | 5.0 |
| Volatile matter (%) | 34.0 |
| Fixed carbon (%) | 52.0 |
| Ash (%) | 9.0 |
| Sulphur (%) | 0.5 |
| Btu/lb | 12,500 |
| Mj/kg | 29.08 |
| FSI | 3 |
| Hardgrove Index | 57 |
| Rmax% | 0.64 - 0.86 |

(Ryan, 2002)


Bowron River Coalfield



The Bowron River Coalfield is located in the east-central part of the province, 50 kilometres east-southeast of Prince George. The coalfield is 2.5 kilometres wide and 15 kilometres long. Coal was first reported in 1871, however, it was not until 1910-11 that development work started with the construction of an adit and survey of the area.

The lower 85 metres of the more than 700 metre thick Tertiary sedimentary section is coal-bearing. Seams dip at 20 to 60 degrees to the northeast. The section contains at least three seams with a cumulative thickness of 8.5 metres. The coal, which is high-volatile B bituminous in rank, is characterized by a high amber content (8 per cent). Considering only the lower seam, there is an estimated potential resource of 400 million tonnes down to a depth of 1200 metres.

Any coalbed gas potential will depend on the presence of biogenic methane though the presence of amber may help initiate generation of thermogenic methane at a lower rank. There could be a coalbed gas resource of about 40 Bcf.




Map of Bowron River Coalfield

PDF files of Regional Coalbed Gas Maps available at: www.em.gov.bc.ca/subwebs/oilandgas/resource/cbg/maps.htm

Coal

- The lower 85 metres of the more than 700 metre thick Tertiary sedimentary section is coal bearing.



Predominantly flaser-bedded quartz arenites with rippled shale streaks containing occasional horizons exhibiting discrete burrows and other bioturbation. Poorly sorted bands of grit and conglomerate up to two metres occur intermittently. Coalified plant material, coal spar and root structures are scattered throughout. Coal lenses, bands and seams are abundant. Amber occurs in the coal, podzols and carbonaceous shales. Slickensiding is common.

Simplified stratigraphic column. Diamond-drillhole GSB-90-3 is sited along strike of the coal zone, close to the western fault and the southwest bank of the Bowron River (Matheson and Sadre, 1991).

- The coal section dips at 20 to 60 degrees to the northeast and contains at least three seams with a cumulative thickness of 8.5 metres. Coal rank is high-volatile B bituminous and is characterized by a high amber content (8 per cent).
- There has been some underground exploration but no mining and no exploration since 1990.

| Average Dry Basis Analysis | |
|----------------------------|--------|
| Volatile matter (%)..... | 36.7 |
| Fixed carbon (%)..... | 42 |
| Ash (%) | 24.6 |
| Sulphur (%) | 1.39 |
| Btu/lb | 11,000 |
| Mj/kg..... | 25.59 |
| FSI | 1 - 3 |
| Hardgrove Index..... | 58 |
| Rmax%..... | 0.65 |

(Ryan, 2002)

- Considering only the lower seam, Matheson and Sadre (1991) estimated a potential resource of 400 million tonnes down to a depth of 1200 metres.

Coalbed Gas

- Any coalbed gas potential will depend on the presence of biogenic methane though the presence of amber may help initiate generation of thermogenic methane at a lower rank.
- There could be a coalbed gas resource of about 40 Bcf (Ryan, 2003).

Key References (see Appendix I, References)

Matheson, A. and Sadre, M. (1991): Subsurface Coal Sampling Survey, Bowron River Coal Deposits, Central British Columbia; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 391-397.


Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/Mining/Geolsurv/coal/coalinbc/coal_bc.htm

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.

Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4, pages 29-68.

Coal River Coalfield



The Coal River Coalfield is located in northern British Columbia between the Yukon border and the Liard River and is crossed by the Alaska Highway in the south. About 10 kilometres north of the highway, Coal River cuts through an area of Tertiary rocks exposing a single seam of lignite. The complete seam is not exposed but partial seam thicknesses range up to 8 metres.


Boulders of coal were first reported at the mouth of Coal River (in the Liard River) in 1891 but it wasn't until the early 1940s that a Geological Survey of Canada field-crew located the source. The coalfield was examined in 1990 by the Provincial geological survey in order to further delineate the deposit.

The basin covers an area of about 35 square kilometres and contains an estimated lignite resource of 100 million tonnes with a possible coalbed gas resource of 6 Bcf.

Coal

- In 1891, transported lignite boulders were reported near the mouth of Coal River (in the Liard River) by R.G. McConnell of the Geological Survey of Canada (GSC). In the early 1940s, M.Y. Williams (GSC) tracked the boulders to their source, approximately 10 kilometres up from the mouth of Coal River. The area was mapped in 1958 and 1960 by H. Gabrielse (GSC) and further examined in 1989 and 1990 (Ryan, 1996). The deposit has not been drilled.
- Coal River cuts through an area of Tertiary rocks exposing a single seam of lignite. The complete seam is not exposed but partial seam thicknesses range up to 8 metres.
- The lignite is overlain by clay which is at least 10 metres thick and is a near white, plastic clay. A darker plastic clay underlies the lignite.
- Tertiary outcrops are restricted to the riverbanks; trees, swamp and a burn zone cover the rest of the area. The area around the river is marked by large crescent-shaped slumps, possibly where younger sediments have slid on the underlying clay layer.
- A water-well drilled near where the Coal River crosses the Alaska Highway intersected 15 metres of coal at a depth of 15 metres. This may or may not be the same seam that outcrops 10 kilometres up Coal River.
- The lignite is cleated with two sets generally developed.
- Based on thickness data, the possible extent of the basin and by using a conservative 5 metre lignite thickness, a potential resource estimate of about 100 million tonnes exists (Ryan, 2002).

| Average Air-Dried Analysis | |
|--------------------------------|-------|
| Moisture (%) as-received | 9.4 |
| Moisture (%) air-dried..... | 13.8 |
| Volatile matter (%)..... | 41.18 |
| Fixed carbon (%)..... | 29.94 |
| Ash (%) | 4.84 |
| Sulphur (%) | 0.15 |
| Btu/lb | 9785 |
| Mj/kg..... | 22.76 |
| Rmax%..... | 0.2 |
| (Ryan, 2002) | |



Representative stratigraphic column of the Coal River lignite occurrence.

Coalbed Gas

- The rank is too low for generation of thermogenic methane, however, the lignite may contain reasonable quantities of biogenic methane.
- A lignite resource of about 100 million tonnes could contain a coalbed gas resource of about 6 Bcf depending on its ability to retain free gas and its ability to adsorb gas (Ryan, 2003).

Key References (see Appendix I, References)

Ryan, B.D. (1991): Geology and Potential Coal and Coalbed Methane Resources of the Tuya River Coal Basin; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 419-429.

Ryan, B.D. (1996): Lignite Occurrences on the Coal River, Northern British Columbia (94M/10); in Geological Fieldwork 1995, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1996-1, pages 271-275.

Ryan, B.D. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website www.em.gov.bc.ca/mining/geosurv/coal/coalinbc/coal_bc.htm.

Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.

Ryan, B.D. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.

REFERENCES

SELECTED GENERAL REFERENCES

- Bickford, C.G.C. and Kenyon, C. (1988): Coalfield Geology of Eastern Vancouver Island; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 441-450.
- British Columbia Coals - Quality and Resources (1992); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Information Circular 1992-18.
- British Columbia Coal Quality Catalog (1992); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Information Circular 1992-20.
- Bustin, R.M. (1999): Free Gas Storage in Matrix Porosity: A Potentially Significant Coalbed Resource in Low Rank Coals; in Proceedings International Coalbed Methane Symposium 1999, May 3-7th, *University of Alabama*, pages 197-214.
- Campbell, D.D., Jory, L.T. and Saunders, C.R. (1977): Geology of the Hat Creek Coal Deposits; *Canadian Mining and Metallurgical Bulletin*, Vol. 70, No. 782, pages 99-108.
- Cathyl-Bickford, C.G. (1991): Coal Geology and Coalbed Methane Potential of Comox and Nanaimo Coalfields, Vancouver Island, British Columbia; *Rocky Mountain Association of Geologists*, pages 155-162.
- Coal Assessment Report 163, Crowsnest Resources Ltd., 1982; *B.C. Ministry of Energy and Mines Library*.
- Coal in British Columbia (1986); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-3.
- Dawson, F.M. (1995): Coalbed Methane: A Comparison Between Canada and the United States; *Geological Survey of Canada*, Bulletin 489.
- Dawson, F.M., Marchioni, D.L., Anderson, T.C. and McDougall, W.J. (2000): An Assessment of Coalbed Methane Exploration Projects in Canada; *Geological Survey of Canada*, Bulletin 549.
- Dolmage Campbell & Associates Ltd. (1975): Coal Resources of British Columbia; *British Columbia Hydro and Power Authority*.
- Dowling, D.B. (1915): Coal Fields of British Columbia; *Geological Survey of Canada*, Memoir 69.
- Dickson, J. (1941): Coal Analyses of British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 14.
- Energy Marketing Update (1992): What is Coal Bed Methane; *B.C. Ministry of Energy, Mines and Petroleum Resources*; Energy Market Update, Vol.4, No.1.
- Feng, K.K., Cheng, K.C. and Augsten, R. (1981): Methane Desorption of Fording Coal from Greenhills Multiple Seams; Division Report, *CANMET*, ERP/MRL 81-67(J).
- Gardner, S.L. (1999): Coal Resources and Coal Mining on Vancouver Island; *B.C. Ministry of Energy and Mines*, Open File 1999-8.
- Grieve D.A. (1990): British Columbia Coal Specifications; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Information Circular 1990-5.
- Grieve D.A. (1992): Relationships Between Coal Quality Parameters in British Columbia Coals; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 397-404.
- Grieve D.A. (1992): British Columbia Coals - Quality and Resources; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Information Circular 1992-18.
- Grieve D.A. (1992): British Columbia Coal Quality Catalog; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Information Circular 1992-20.
- Grieve D.A. (1992): Phosphorous in British Columbia Coking Coals; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1992-20.
- Grieve, D.A. and Holuszko, M.E. (1990): Trace Elements, Mineral Matter and Phosphorous in British Columbia Coals; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 361-370.
- Grieve, D.A., Holuszko, M.E. and Goodarzi, F. (1995): British Columbia Coal Quality Survey, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 96.
- Holuszko, M.E. and Grieve, D.A. (1991): Washability Characteristics of British Columbia Coals; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 371-379.
- Holuszko, M.E. (1994): Washability Characteristics of British Columbia Coals; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-2.
- Kenyon, C.E. (1985): Coalfile; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 399-402.
- Kenyon, C.E. (1986): Coalfile; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 235-236.
- Kenyon, C.E. (1987): Coalfile; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 431-432.
- Lamberson, M.N. and Bustin, R.M. (1992): Coalbed Methane Characteristics of the Gates Formation Lithotypes, Northeastern British Columbia; *Canadian Coal and Coal Bed Methane Geoscience Forum*, Proceedings, pages 275-284.
- Matheson, A., Grieve, D.A., Holuszko, M.E. and Goodarzi, F. (1994): Selected Thermal Coal Basins of British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-3.
- Matheson, A. (1986): Coal in British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-3.
- Matheson, A. (1989, revised): Producing and Developing Thermal Coal Properties in British Columbia; *B.C.*

Ministry of Energy, Mines and Petroleum Resources, Open File 1988-23.

- Matheson, A. and Van Den Bussche, B. (1990): Subsurface Coal Sampling Survey, Central British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 445-448.
- Papic, M.M., Warren, L.H. and Woodley, R.M. (1977): Hat Creek Utilization; *Canadian Mining and Metallurgical Bulletin*, November 1977, pages 99-105.
- Pearson, D.E. and Grieve, D.A. (1985): Rank Variation, Coalification Pattern and Coal Quality in the Crowsnest Coalfield, British Columbia; *Canadian Institute of Mining and Metallurgy Bulletin*, September, pages 39-36.
- Proudlock, P. (1990): An Overview of Albertan and Canadian Demethanation Experience, Technology Advances and Development Potential; in Coalbed Methane in Alberta - What's it all about; *Alberta Geological Survey*, Information Series No.108.
- Read, P.B. (1987): Industrial Minerals In Some Tertiary Basins, Southern British Columbia; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 247-254.
- Ryan, B.D. (1992): An Equation for Estimation of Maximum Coalbed-Methane Resource Potential; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 393-396.
- Ryan, B.D. and Price J.T. (1993): The Predicted Coke Strength after Reaction Values of British Columbia Coals, With Comparisons to International Coals; in Geological Fieldwork 1992, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1993-1, pages 507-516.
- Ryan, B.D. and Dawson, F.M. (1994): Coalbed Methane Canister Desorption Techniques; in Geological Fieldwork 1993, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-1, pages 245-256.
- Ryan, B.D. (1996): Controls on Coal Washing Difficulty, (82G/15 93L/11); in Geological Fieldwork 1995, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1996-1, pages 259-269.
- Ryan, B.D. and Grieve, D.A. (1996): Source and Distribution of Phosphorus in British Columbia Coal Seams; in Geological Fieldwork 1995, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1996-1, pages 277-294.
- Ryan, B.D., Gransden, J. and Price, J. (1998): Fluidity of Western Canadian Coals and Its Relationship to other Coal and Coke Properties; in Geological Fieldwork 1997, *B.C. Ministry of Employment and Investment*, Paper 1998-1, pages 27-1 to 27-17.
- Ryan, B.D., Leeder, R., Price, J.T. and Gransden, J.F. (1999): The Effect of Coal Preparation on the Quality of Clean Coal and Coke; in Geological Fieldwork 1998, *B.C. Ministry of Energy and Mines*, Paper 1999-1, pages 247-275.
- Ryan, B. (2002): Coal in British Columbia; *B.C. Ministry of Energy and Mines*, Website, http://www.em.gov.bc.ca/Mining/Geolsurv/coal/coalinc/coal_bc.htm.
- Ryan, B. (2003): A Summary of Coalbed Methane Potential in British Columbia; *Canadian Society of Exploration Geophysicists Recorder*, Vol.28, No.9, November 2003, pages 32-40.
- Ryan, B.D. (2003): Pseudovitrinite: Possible Implications for Gas Saturation in Coals and Surrounding Rocks; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 203-212.
- Ryan, B.D. (2003): Overview of the Coalbed Methane Potential of Tertiary Coal Basins in the Interior of British Columbia; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 213-236.
- Ryan, B.D. (2003): Cleat Development in Some British Columbia Coals; in Geological Fieldwork 2002, *B.C. Ministry of Energy and Mines*, Paper 2003-1, pages 237-256.
- Schroeter, T.G. (1985): Mineral and Coal Potential, Northwestern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 58.
- Schroeter, T.G., White, G.V. and Koo, J. (1986): Coal in Northwestern British Columbia - An Overview; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-5.
- Smith, G.G. (1989): Coal Resources of Canada; *Geological Survey of Canada*, Paper 89-4.
- Thompson, B.J. (1985): Canada-British Columbia Coal Data Project; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 403-404.
- Thompson, B.J. (1986): Canada-British Columbia Coal Data Project; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 237-238.
- Van den Bussche, B.G. and Grieve, D.A. (1990): Phosphorus in British Columbia Coking Coals; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 427-430.

MINISTRY OF ENERGY AND MINES REFERENCES FOR INDIVIDUAL COALFIELDS

Bowron River

- Linds, R.K. (1982): Bowron River Progress Report; in Geological Fieldwork 1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1982-1, pages 227-228.
- Matheson, A. and Sadre, M. (1991): Subsurface Coal Sampling Survey, Bowron River Coal Deposits, Central British Columbia; in Geological Fieldwork

1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 391-397.

Coal River

Ryan, B.D. (1991): Geology and Potential Coal and Coalbed Methane Resources of the Tuya River Coal Basin; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 419-429.

Ryan, B.D. (1996): Lignite Occurrences on the Coal River, Northern British Columbia (94M/10); in Geological Fieldwork 1995, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1996-1, pages 271-275.

Comox

Bickford, C.G.C., Hoffman, G. and Kenyon, C. (1989): Coal Measures Geology of the Quinsam Area; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 559-563.

Bickford, C.G.C., Hoffman, G. and Kenyon, C. (1990): Geological Investigations in the Coal Measures of the Oyster River, Mount Washington and Cumberland Area, Vancouver Island; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 431-437.

Cathyl-Bickford, C.G. (1992): Geology and Energy Resource Potential of The Tsable River and Denman Island; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 419-426.

Cathyl-Bickford, C.G. and Hoffman, G.L. (1998): Geological maps of the Nanaimo and Comox Coalfields, *B.C. Ministry of Energy and Mines*, Open File 1998-7.

Cathyl-Bickford, C.G. (2001): Lithostratigraphy of the Comox and Trent River Formations in the Comox Coalfield, Vancouver Island (92F/7, 10, 11, 14); in Geological Fieldwork 2000, *B.C. Ministry of Energy and Mines*, Paper 2001-1, pages 363-370.

Eastwood, G.E.P. (1977): Quinsam Area; in Geology in British Columbia 1977-1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 69-75.

Eastwood, G.E.P. (1978): Quinsam Area, Vancouver Island; in Geological Fieldwork 1977, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1978-1, pages 55-56.

Eastwood, G.E.P. (1979): Quinsam Area, Vancouver Island; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 66-67.

Holuszko, M.E., Matheson, A. and Grieve, D.A. (1993): Pyrite Occurrences in Telkwa and Quinsam Coal Seams; in Geological Fieldwork 1992, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1993-1, pages 527-546.

Kenyon, C., Cathyl-Bickford, C.G. and Hoffman, G. (1991): Quinsam and Chute Creek Coal Deposits (NTS (92/13,14); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-3.

Matheson, A.H. (1990): Subsurface Coal Sampling Survey, Quinsam Area, Vancouver Island, British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 439-443.

Ronaghan, R.J. (1980): Geology of the Coal Reserve in the Quinsam Mining Block; Coal Assessment Report 821, *B.C. Ministry of Energy and Mines*.

Ryan, B.D. and Dawson, F.M. (1994): Coalbed Methane Desorption Results from the Quinsam Coal Mine and Coalbed Methane Resource of the Quinsam Coalfield, British Columbia, Canada (92F/13,14); in Geological Fieldwork 1993, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-1, pages 215-224.

Ryan, B.D. (1995): Calcite in Coal from the Quinsam Coal Mine, British Columbia, Canada, Origin, Distribution and Effects on Coal Utilization; in Geological Fieldwork 1994, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1995-1, pages 243-260.

Ryan, B.D. (1997): Coalbed Methane in the Comox Formation in the Tsable River Area, Vancouver Island; in Geological Fieldwork 1996, *B.C. Ministry of Employment and Investment*, Paper 1997-1, pages 353-363.

Ryan, B.D. (2002): Note on Desorption Results of Comox Formation Coals from the Courtenay Area, Vancouver Island, British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 319-330.

Van der Flier-Keller, E. and Dumais, S. (1988): Inorganic Matter Content and Specialized Element Potential of the Nanaimo and Comox Coalfields, Vancouver Island (92G,F,K); in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 435-439.

East Kootenay

Bustin, R.M. (1981): Preliminary Investigations as to the Effects of Shear on Coal Quality in Southeastern British Columbia; in Geological Fieldwork 1980, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-1, pages 133-148.

Geology of Crowsnest Coalfield (1981), Southern Part; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 42.

Gigliotti, F.B. and Pearson, D.E. (1978): Preliminary Geological Map of Crowsnest Coalfield, Northeast Part; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 31.

Grieve, D.A. and Pearson, D.E. (1985): Geology and Rank Distribution of the Elk Valley Coalfield; in Geology in British Columbia 1977-1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 17-23.

- Grieve, D.A. and Pearson, D.E. (1982): Geology of the Greenhills Range, Elk Valley Coalfield; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 51.
- Grieve, D.A. and Ollerenshaw, N.C. (1982): Stratigraphy of the Elk Formation in Southeastern British Columbia; in Geological Fieldwork 1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1982-1, pages 51-53.
- Grieve, D.A. and Elkins, P.R. (1986): Correlation and Comparison of Two Coal-Bearing Zones Between Ewin Pass and Bare Mountain, Elk Valley Coalfield, Southeastern British Columbia; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 17-24.
- Grieve, D.A. and Fraser, J.M. (1983): Line Creek and Crown Mountain Areas, Elk Valley Coalfield; in Geological Fieldwork 1982, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-1, pages 21-26.
- Grieve, D.A. and Kilby, W. (1985): Structural Modelling of Parts of the Northern Dominion Coal Block (Parcel 73), Southeastern British Columbia; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 53-56.
- Grieve, D.A. and Kilby, W. (1986): Flathead Ridge Coal Area, Southern Dominion Coal Block (Parcel 82), Southern British Columbia; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 25-36.
- Grieve, D.A. and Price, R.A. (1987): Geological Setting of the South Half of the Elk Valley Coalfield; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 63.
- Grieve, D.A. and Fraser, J.M. (1985): Geology of the Elk Valley Coalfield, Southern Half (Kilmarnock Creek to Alexander Creek); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 60.
- Grieve D.A. and Ollerenshaw, N.C. (1989): Stratigraphy of the Elk Formation in the Fernie Basin, Southeastern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-2.
- Grieve D.A. and Kilby, W. (1989): Geology and Coal Resources of the Dominion Coal Block, Southeastern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-4.
- Grieve, D.A. and Goodarzi, F. (1994): Trace Elements in Coals of the East Kootenay and Peace River Coalfields, British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1994-15.
- Grieve, D. (1981): Elk Valley Coalfield; in Geological Fieldwork 1980, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-1, pages 71-72.
- Grieve, D. (1981): Flathead Coalfield; in Geological Fieldwork 1980, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-1, pages 65-70.
- Grieve, D. (1982): Mount Banner Area, Elk Valley Coalfield; in Geological Fieldwork 1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1982-1, pages 46-50.
- Grieve, D.A. (1984): Tonsteins: Possible Stratigraphic Correlation Aids in East Kootenay Coalfields; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 36-41.
- Grieve, D. (1985): Geology of the Mount Banner Area, Elk Valley Coalfield; in Geology in British Columbia 1977-1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 24-33.
- Grieve, D. (1986): Crowsnest Coalfield; in Exploration in British Columbia 1985; *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages B33-B38.
- Grieve, D. (1987): Coal Rank Distribution, Flathead Coalfield, Southeastern British Columbia; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 361-364.
- Grieve, D. (1987): Subsurface Coal Rank Profiles Ewin Pass to Bare Mountain, Elk Valley Coalfield, Southeastern British Columbia; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 351-360.
- Grieve, D. (1989): Numerical Depositional Modelling of the Elk Valley Coalfield; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 535-541.
- Grieve, D. (1993): Geology and Rank Distribution of the Elk Valley Coalfield, Southeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 82.
- Holuszko, M.E. (1992): Washability of Peace River and East Kootenay Coals; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 405-417.
- Holuszko, M.E. (1993): Washability of Lithotypes from a Selected Seam in the East Kootenay Coalfield, Southeast British Columbia; in Geological Fieldwork 1992, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1993-1, pages 517-525.
- Johnson, D.G.S. and Smith, L.A. (1991): Coalbed Methane in Southeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Petroleum Geology, Special Paper 1991-1.
- Morris, R.J. and Grieve, D.A. (1988): Elk Valley Coalfield, North Half; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 427-434.
- Morris, R.J. and Grieve, D.A. (1990): Geology of the Elk Valley Coalfield, North Half (Henrietta Creek to Elk Lakes); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 68.
- Newmarch, C.G. (1953): Geology of the Crowsnest Coal Basin With Special Reference to the Fernie Area; *British Columbia Department of Mines*, Bulletin 33.
- Pearson, D.E. (1977): East Kootenay Coalfield; in Geological Fieldwork 1976, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1977-1, pages 77-82.
- Pearson, D.E. and Duff, P.M.D. (1976): Studies in the East Kootenay Coalfields; in Geological Fieldwork 1975,

B.C. Ministry of Energy, Mines and Petroleum Resources, Paper 1976-1, pages 93-98.

- Pearson, D.E., Gigliotti, F.B., Ollerenshaw, N.C. and Grieve, D.A. (1977): Preliminary Geological Map of Crowsnest Coalfield, Northwest Part; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 24.
- Pearson, D.E. and Grieve, D.A. (1978): Preliminary Geological Map of the Crowsnest Coalfield, West Part; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 27.
- Pearson, D.E. and Grieve, D.A. (1979): Crowsnest Coalfield; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 61-65.
- Pearson, D.E. and Grieve, D.A. (1980): Elk Valley Coalfield (82J/2); in Geological Fieldwork 1979, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1980-1, pages 91-96.
- Pearson, D.E. and Grieve, D.A. (1980): Coal Measures in the Crowsnest Coalfield; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 42.
- Ryan, B.D. and Khan, M. (1998): Maceral Affinity of Phosphorus in Coals from the Elk Valley Coalfield, British Columbia; in Geological Fieldwork 1997, *B.C. Ministry of Employment and Investment*, Paper 1998-1, pages 28-1 to 28-19.
- Ryan, B.D. and Takkinen, M. (2000): In Situ fracture Porosity and Specific Gravity of Highly Sheared Coals from Southeast British Columbia; in Geological Fieldwork 1999, *B.C. Ministry of Energy and Mines*, Paper 2000-1, pages 359-371.
- Ryan, B.D. (2000): A Note on Difficult to Wash Coals from Southeast British Columbia; in Geological Fieldwork 1999, *B.C. Ministry of Energy and Mines*, Paper 2000-1, pages 373-382.
- Ryan, B.D. (2004): The CBM Resource of Some Prospective Areas of the Crowsnest Coalfield; *B.C. Ministry of Energy and Mines*, in press, Oil and Gas Emerging Opportunities and Geosciences Branch.

Graham Island

- Sutherland-Brown, A. (1968): Geology of the Queen Charlotte Islands, British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 54.

Hat Creek

- Church, B.N. (1976): Geology of the Hat Creek Basin; in Geological Fieldwork 1975, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1976-1, pages 104-108.
- Church, B.N. (1977): Geology of the Hat Creek Coal Basin; in Geology in British Columbia 1975, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 99-118.

Church, B.N. (1980): Hat Creek Bocanne-Buchite; in Geological Fieldwork 1979, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1980-1, pages 97-100.

Church, B.N. (1981): Further Studies of the Hat Creek Coal Deposit; in Geological Fieldwork 1980, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-1, pages 73-78.

Klappan and Groundhog

- Buckham, A.F. and Latour, B.A. (1950): The Groundhog Coalfield, British Columbia; *Geological Survey of Canada*, Bulletin 16.
- Cookenboo, H.O. and Bustin, R.M. (1990): Stratigraphy of Coal Occurrences in the Bowser Basin; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 473-477.
- Cookenboo, H.O. and Bustin, R.M. (1991): Coal-Bearing Facies in the Northern Bowser Basin; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 415-418.
- Gilchrist, R.D. (1979): Groundhog Coalfield; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 84-85.
- Koo, J. (1984): The Telkwa, Red Rose and Klappan Coal Measures in Northwestern British Columbia; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 81-90.
- Koo, J. (1986): Geology of the Klappan Coalfield in Northwestern British Columbia; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 225-228.
- Koo, J. (1986): Geology of the Klappan Coalfield in Northwest British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1986-3.
- Ryan, B.D. and Dawson, F.M. (1993): Coal and Coalbed Methane Potential of the Bowser Basin, Northern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1993-31.
- Schroeter, T.G. and White, G.V. (1985): The Sustut Coal Measures; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 285-290.

Merritt

- Matheson, A. (1992): Subsurface Thermal Coal Sampling Survey, Merritt Coal Deposits, South-Central British Columbia; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 427-432.

Nanaimo

- Bickford, C.G.C. (1989): Geology, Mining Conditions and Resource Potential of the Wellington Coal Bed, Georgia Basin; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 553-558.
- Cathyl-Bickford, C.G. and Hoffman, G.L. (1991): Geology and Coal Resources of the Nanaimo Group in the Alberni, Ash River, Cowie Creek and Parksville Areas, Vancouver Island; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 381-385.
- Cathyl-Bickford, C.G., Wilson, R. and Hoffman, G.L. (1992): Geology, Mineability and Coalbed Gas Potential of the Douglas Coal Zone in the Nanaimo Coalfield, Vancouver Island; *Canadian Coal and Coal Bed Methane Geoscience Forum*, Proceedings, pages 293-309.
- Cathyl-Bickford, C.G. and Hoffman, G.L. (1998): Geological maps of the Nanaimo and Comox Coalfields, *B.C. Ministry of Energy and Mines*, Open File 1998-7.
- Kenyon, C. and Bickford, C.G.C. (1989): Vitrinite Reflectance Study of Nanaimo Group Coals of Vancouver Island; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 543-552.
- Van der Flier-Keller, E. and Dumais, S. (1988): Inorganic Matter Content and Specialized Element Potential of the Nanaimo and Comox Coalfields, Vancouver Island (92G,F,K), in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 435-439.

Peace River

- Broatch, J. (1982): An Investigation of The Palynology of the Peace River Coalfield, Northeastern British Columbia; in Geological Fieldwork 1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1982-1, pages 113-121.
- Broatch, J. (1986): Palynological Zonation and Correlation of the Peace River Coalfield, Northeastern British Columbia; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 321-326.
- Broatch, J. (1987): Palynological Zonation and Correlation of the Peace River Coalfield, Northeastern British Columbia, An Update; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 379-384.
- Broatch, J.C. (1988): A Summary of the Results of a Palynological Investigation of British Columbia's Northeast Coalfield; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 451-458.
- Carmichael, S.M.M. (1982): Depositional Environments and Paleocurrent Trends in the Gates Member, Northeast Coalfield; in Geological Fieldwork 1981, *B.C.*

Ministry of Energy, Mines and Petroleum Resources, Paper 1982-1, pages 244-258.

- Cunningham, J.M. and Sprecher, B. (1992): Peace River Coalfield Digital Mapping Program; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 441-449.
- Cunningham, J.M. and Sprecher, B. (1992): Geology of the Lemoray Creek and Carbon Creek Areas, Northeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1992-12.
- Duff, P. McL.D. (1978): Correlation Studies in the Peace River Coalfield; in Geological Fieldwork 1977, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1978-1, pages 60-61.
- Duff, P. McL.D. (1979): Correlation Studies in the Peace River Coalfield; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 86-87.
- Duff, P. McL.D. and Gilchrist, R.D. (1981): Correlation of Lower Cretaceous Coal Measures, Peace River Coalfield; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-3.
- Flynn, B.P., McMechan, R.D. and McMechan M. (1977): Geology of the Southeastern Peace River Coalfield; in Geological Fieldwork 1976, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1977-1, page 83.
- Flynn, B.P., McMechan, R.D. and McMechan M. (1977): Wapiti River Area, Peace River Coalfield; in Geology in British Columbia 1976, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 137-147.
- Gilchrist, R.D. (1979): Burnt River Area; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 79-83.
- Gilchrist, R.D. (1980): Wolverine-Hasler Map-Area; in Geological Fieldwork 1979, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1980-1, pages 101-102.
- Gilchrist, R.D., Flynn, B.P. and Hauser, R.I. (1978): Mount Spieker Area; in Geological Fieldwork 1977, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1978-1, pages 57-59.
- Gilchrist, R.D. and Flynn, B.P. (1978): Coal Resources, Peace River Coalfield, Northeastern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 33.
- Grieve, D.A. and Goodarzi, F. (1994): Trace Elements in Coals of the East Kootenay and Peace River Coalfields, British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1994-15.
- Holuszko, M.E. (1992): Washability of Peace River and East Kootenay Coals; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 405-417.
- Hughes, J.E. (1964): Jurassic and Cretaceous Strata of the Bullhead Succession in the Peace and Pine River Foothills; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 51.

- Hughes, J.E. (1967): Geology of the Pine Valley, Mount Wabi to Solitude Mountain, Northeastern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 52.
- Hunter, D.J. and Cunningham, J.M. (1991): Burnt River Mapping and Compilation Project; in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 407-414.
- Hunter, D.J. and Cunningham, J.M. (1991): Geology of the Burnt River and Gwillum Lake (Southwest Half) Areas, Northeastern British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1991-4.
- Jahans, P.C. (1992): Pine Valley Mapping and Compilation Project; in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 433-440.
- Jahans, P.C. (1993): Geology of the Pine River Area; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1993-22.
- Karst, R.H. (1979): Vitrinite Reflectance As A Correlation Tool in the Carbon Creek Coal Measures; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 73-78.
- Karst, R.H. (1979): Coal Property Examinations, Peace River Coalfield; in Geological Fieldwork 1978, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1979-1, pages 67-72.
- Karst, R.H. (1981): Correlation of the Lower Cretaceous Stratigraphy of Northeastern British Columbia From Foothills to Plains; in Geological Fieldwork 1980, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1981-1, pages 79-90.
- Karst, R. and White, G.V. (1980): Coal Rank Distribution within the Bluesky-Gething Stratigraphic Horizon of Northeast B.C.; in Geological Fieldwork 1979, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1980-1, pages 103-108.
- Kilby, W. (1984): Tonsteins and Bentonites in Northeast British Columbia; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 95-107.
- Kilby, W. (1984): The Character of the Bluesky Formation in the Foothills of Northeastern British Columbia; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 108-112.
- Kilby, W. (1985): Tonstein and Bentonite Correlations in Northeast British Columbia; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 257-277.
- Kilby, W. (1986): Biaxial Reflecting Coals in the Peace River Coalfield; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 127-138.
- Kilby, W. (1986): Some Chemical and Mineralogical Characteristics of Tonsteins and Bentonites in Northeast British Columbia; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 139-150.
- Kilby, W. (1989): Tectonically Altered Coal Rank, Boulder Creek Formation, Northeastern British Columbia; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 565-570.
- Kilby, W.E. and Hunter, D.J. (1990): Tumbler Ridge, Northeast British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 455-459.
- Kilby, W.E. and Johnston, F.T. (1988): Kinuseo Mapping and Compilation Project; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 463-470.
- Kilby, W.E. and Johnston, F.T. (1988): Bedrock Geology of the Kinuseo Creek Area, Northeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1988-21.
- Kilby, W.E. and Johnston, F.T. (1988): Bedrock Geology of the Kinuseo Falls Area, Northeast British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1988-22.
- Kilby, W.E. and Oppelt, H.P. (1985): Numerical Depositional Modelling - Gething Formation Along the Peace River; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 233-250.
- Kilby, W.E. and Wrightson, C.B. (1987): Bullmoose Mapping and Compilation Project; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 373-378.
- Kilby, W.E. and Wrightson, C.B. (1987): Sukunka River; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1987-7.
- Lamberson, M.N. and Bustin, R.M. (1989): Petrology, Sedimentology and Geochemistry of Gates Formation Coals, Northeastern British Columbia; in Geological Fieldwork 1988, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1989-1, pages 571-576.
- Lamberson, M.N., Bustin, R.M., Kalkreuth, W. and Pratt, K.C. (1990): Lithotype Characteristics and Variation in Selected Coal Seams of the Gates Formation, Northeastern British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 461-468.
- Legun, A. (1983): Stratigraphy and Sedimentology Notes on the Bullhead Mountain - Peace River Canyon, Carbon Creek Area, Northeastern British Columbia; in Geological Fieldwork 1982, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-1, pages 93-98.
- Legun, A. (1984): Butler Ridge Map-Area, Peace River District; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 123-130.
- Legun, A. (1984): Stratigraphic and Depositional Relationships Between the Bluesky Marker Unit, Gething Marine Tongue, and Upper Coal Measures of the Gething Formation; in Geological Fieldwork 1983,

- B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 117-122.
- Legun, A. (1984): Geology of the Butler Ridge Area; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Preliminary Map 57.
- Legun, A. (1985): Eastern Limit of Upper Coal Measures of the Gething Formation (Chamberlain Member), Peace River Coalfield; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 233-250.
- Legun, A. (1985): Geology of the West Carbon Creek Area; in Geological Fieldwork 1984, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1985-1, pages 227-232.
- Legun, A. (1986): Geology of the Carbon Creek Area; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 155-160.
- Legun, A. (1987): A Geological Update of the Carbon Creek and Butler Ridge Areas; in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 365-368.
- Legun, A. (1987): Relation of Gething Formation Coal Measures to Marine Paleoshorelines, (93P, 93I); in Geological Fieldwork 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1987-1, pages 369-372.
- Legun, A. (1988): Coal Trends in the Gething Formation - An Update; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 459-462.
- Legun, A. (1987): Geology of the Carbon Creek Area (0930/15); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1987-21.
- Legun, A. (1988): Geology and Coal Resources of the Carbon Creek Map Area; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-3.
- Legun, A. (1990): Stratigraphic Trends in the Gething Formation (93P/1 to 8 and 93I/14,15); *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1990-33.
- Legun, A.S. (2002): Overview of Coalbed Methane Geology in Northeast British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 247-256.
- Legun, A.S. (2003): Coalbed Methane Geology of the Peace River District, Northeastern British Columbia; *B.C. Ministry of Energy and Mines*, Geoscience File 2003-2, 2 sheets.
- Mathews, W.H. (1946): Geology and Coal Resources of the Carbon Creek-Mount Bickford Map Area; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 24.
- McKechnie, N.D. (1955): Coal Reserves of the Hasler Creek-Pine River Area, British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Bulletin 36.
- Oppelt, H.P. (1986): Stratigraphy and Character of the Bluesky Formation; in Geological Fieldwork 1985, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1986-1, pages 161-166.
- Ryan, B.D. (1990): Preliminary Survey of the Coal Resources of the Upper Cretaceous Rocks, Northeastern British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 469-471.
- Ryan, B.D. (1997): Coal Quality Variations in the Gething Formation, Northeast British Columbia (93O,J,I); in Geological Fieldwork 1996, *B.C. Ministry of Employment and Investment*, Paper 1997-1, pages 373-397.
- Ryan, B.D. and Lane, B. (2002): Adsorption Characteristics of Coals with Special References to the Gething Formation, Northeast British Columbia; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 331-346.
- Ryan, B.D., Fournier, M. and Ditttrick, M. (2002): Selenium Concentrations in Mine Refuse and Mist Mountain Rocks; Evaluation of Variations Laterally and Over Time; in Geological Fieldwork 2001, *B.C. Ministry of Energy and Mines*, Paper 2002-1, pages 347-364.
- Stott, D.F. and Gibson, D.W. (1980): Minnes Coal, Northeastern British Columbia; in Current Research, Part C, *Geological Survey of Canada*, Paper 80-1C, pages 135-137.
- White, G.V. (1983): The Bluesky - A Stratigraphic Marker in Northeastern British Columbia; in Geological Fieldwork 1982, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-1, pages 85-88.
- Wrightson, C.B. (1988): Geology of Duke and Honeymoon Pit Areas, Monkman Coal Deposit, Northeast British Columbia; in Geological Fieldwork 1987, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1988-1, pages 471-480.
- Ziegler, W.H. and Pocock, S.A.J. (1960): The Minnes Formation; Second Annual Field Trip Guide Book, Rock Lake, *Edmonton Geological Society*, pages 43-67.

Princeton

- McMechan, R.D. (1976): Princeton Basin; in Geological Fieldwork 1975, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1976-1, pages 99-103.
- McMechan, R.D. (1983): Geology of the Princeton Basin; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-3.

Suquash

- Gardner, S.L. (1984): Geological Reconnaissance of Vancouver Island Coal Areas - Suquash Coal Basin and Outliers, Northern Vancouver Island, British Columbia; Coal Assessment Report 778, *B.C. Ministry of Energy and Mines*.
- Kenyon, C. (1991): The Suquash Coalfield, Vancouver Island; in Geological Fieldwork 1990, *B.C. Ministry of*

Energy, Mines and Petroleum Resources, Paper 1991-1, pages 387-390.

Telkwa

- Holuszko, M.E., Matheson, A. and Grieve, D.A. (1993): Pyrite Occurrences in Telkwa and Quinsam Coal Seams; in Geological Fieldwork 1992, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1993-1, pages 527-546.
- Koo, J. (1983): Telkwa Coalfield, West-Central British Columbia; in Geological Fieldwork 1982, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-1, pages 113-122.
- Koo, J. (1984): The Telkwa, Red Rose and Klappan Coal Measures in Northwestern British Columbia; in Geological Fieldwork 1983, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1984-1, pages 81-90.
- Matheson, A.H. and Van Den Bussche, B. (1990): Subsurface Coal Sampling Survey, Telkwa Area, Central British Columbia (93L/11); in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 445-448.
- Malott, M.L. (1990): Telkwa Coal North (093L 152); in Exploration in British Columbia 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 175-179.
- Palsgrove, R.J. and Bustin, R.M. (1991): Stratigraphy, Sedimentology and Coal Quality of the Lower Skeena Group, Telkwa Coalfield, Central British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-2.
- Palsgrove, R.J. and Bustin, R.M. (1990): Stratigraphy and Sedimentology of the Lower Skeena Group, Telkwa Coalfield, Central British Columbia; in Geological Fieldwork 1989, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1990-1, pages 449-454.
- Ryan, B.D. and Dawson, M.F. (1994): Potential Coal and Coalbed Methane Resource of the Telkwa Coalfield, Central British Columbia; in Geological Fieldwork 1993, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1994-1, pages 225-243.
- Ryan, B.D. and Ledda, A. (1998): A Review of Sulphur in Coal: With Specific Reference to the Telkwa Deposit, North-Western British Columbia; in Geological Fieldwork 1997, *B.C. Ministry of Employment and Investment*, Paper 1998-1, pages 29-1 to 29-21.
- Ryan, B.D. (1990): Geology and Potential Coal and Coalbed Methane Resources of the Seaton Coal Basin (93M/3,7); in Exploration in British Columbia 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 153-167.
- Ryan, B.D. (1991): Density of Coals from the Telkwa Coal Property, Northwestern British Columbia, (93L/11); in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 399-406.

- Ryan, B.D. (1992): Coal Rank Variations in the Telkwa Coalfield, Central British Columbia (93L/11); in Geological Fieldwork 1991, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1992-1, pages 451-460.
- Ryan, B.D. (1993): Geology of the Telkwa Coalfield, British Columbia; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Open File 1993-21.
- Ryan, B.D. (1997): Coal Rank Variations in the Cabinet Creek Area, Telkwa Coalfield, Central British Columbia (93L/11); in Geological Fieldwork 1996, *B.C. Ministry of Employment and Investment*, Paper 1997-1, pages 365-372.
- Telkwa Coal (1978): in Geology in British Columbia 1977-1981, *B.C. Ministry of Energy, Mines and Petroleum Resources*, page 129.
- White, G.V. (1987): Guess Creek Coal Deposit; in Exploration in British Columbia 1986, *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages B65-B70.

Tulameen

- Church, B.N. and Brasnett, D. (1983): Geology and Gravity Survey of the Tulameen Coal Basin; in Geological Fieldwork 1982, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1983-1, pages 47-55.
- Evans, S.H. (1978): Tulameen Coal Basin; in Geological Fieldwork 1977, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1978-1, pages 83-85.
- Evans, S.H. (1985): Geology of the Tulameen Coal Basin; in Geology in British Columbia 1977-1981; *B.C. Ministry of Energy, Mines and Petroleum Resources*, pages 76-87.
- Williams, V.E. (1978): Coal Petrology of the Tulameen Coalfield, South Central British Columbia, M.Sc. Thesis, Western Washington University; in Coal Assessment Report 832, *B.C. Ministry of Energy and Mines*, 83 pp.

Tuya River

- De Nys, F.J.G. (1980): Thundercloud Coal Project Technical Report; PetroCanada Exploration, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 243.
- Reid, J.L. (1980): Thundercloud Coal Project Technical Report, PetroCanada Exploration, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 242.
- Ryan, B.D. (1991): Geology and Potential Coal and Coalbed Methane Resources of the Tuya River Coal Basin, in Geological Fieldwork 1990, *B.C. Ministry of Energy, Mines and Petroleum Resources*, Paper 1991-1, pages 419-429.
- Panteleyev, A., Bailey, Bloodgood, M.A. and Hancock, K.D. (1996): Geology and Mineral Deposits of the Quesnel River - Horsefly Map Area, Central Quesnel Trough,

British Columbia; *B.C. Ministry of Employment and Investment*, Bulletin 97, 156 pages.

Shives, R.B.K., Holman, P.B., Rebolledo, L., Hetu, R.J. (1995): Airborne geophysical survey, Mount Polley area, B.C.; *Geological Survey of Canada*, Open File 2802, 82 pages.

Vincent, B.D. (1979): Geological Mapping, Tuya River Property, British Columbia: Esso Minerals Canada; *B.C. Ministry of Energy, Mines and Petroleum Resources*, Coal Assessment Report 246.