




Geological Survey Branch

## MINERAL POTENTIAL MAP 1992-5

# METALLIC MINERAL POTENTIAL OF THE CHUCHI LAKE MAP AREA

NTS 93N/2E


#### BY K. BELLEFONTAINE AND J. NELSON



#### INTRODUCTION

The Chuchi Lake map area is underlain by the Takla Group and is intruded by numerous alkaline intrusions. The overall favourable mineral potential of the Takla Group and the close proximity to the Mt. Milligan (93N/1) and Chuchi (93N/7) alkaline porphyry deposits suggests a promising metallogenic environment. The entire area is designated as having moderate mineral potential with high potential zones scattered throughout. Small prospects of skarn and polymetallic veins also occur in the area.

#### CLASSIFICATION OF MINERAL POTENTIAL



#### QUALITATIVE DESCRIPTIONS OF METALLIC MINERAL POTENTIAL CLASSIFICATIONS

6	Extremely High	Known deposits (developed prospects) with significant identified resources in the ground. Very favourable supporting data from all applicable areas; very high degree of confidence in designation. Continued exploration highly probable; potential for mine developement is high.
5	Very High	Known prospects, developed prospects, past producing mines (with undefined resources in the ground) in highly favourable geological environment. Very good supporting data from all three criteria, high degree of confidence in designation. Future exploration expected.
4	High	Known or indicated resources in highly favourable geological environment. Very good supporting geological and geochemical data. Resources or deposits not necessarily identified; however, positive indications that deposit forming processes were active in the area. Moderate to high degree of confidence in designation. Future exploration expected.
3	Moderate	Favourable geologiacl environment for resource occurrence. Very favourable geological data from one or more criteria; but significant occurrences lacking. Moderate degree of confidence in designation. Future exploration likely, particularly if near an area of higher designation. Good potential for upgrading classification.
2	Moderate	Moderately favourable environment for resource accumulation as indicated by one line of evidence (ie. favourable geology, mineralization or anomalous geochemistry with no other supporting data). Areas generally lack sufficient prospecting to identify resources. Moderate to low degree of confidence in designation. Future reconnaissance exploration to be expected. Good potential for upgrading classification.
1	Low	Current data indicates that the existence of resources is unlikely. Moderate to high degree of confidence in designation. Future exploration for deposit types considered is unlikely.
U	Unknown	Current data is either outdated or insufficiently detailed for a reasoned determination of mineral potential. High degree of confidence in designation. Future exploration could occur

Description

SYMBOLS		ANOMALOUS THRESHOLDS				
	Developed Prospect/Deposit (Major Resources)	Zn	128 ppm 80 ppm	As Sb	17 ppm 1 ppm	ly s
$\blacksquare$	Prospect/Deposit (Minor Resources)		9 ppm 0.5 ppm	Au	9 ppb	
	Showing	Ag	0.5 ppm			

### MINERAL DEPOSIT MODELS

Deposit Type	Commodities	Geological Indicators	Example
Alkaline Porphyries	Cu, Au	Crowded porphyritic alkaline intrusions; potassic and propylitic alteration; probable structural control.	Mt. Milligan (93N/1) 400 million tonnes 0.2% Cu, 0.48 g/t Au
Polymetallic Veins	Au, As, Sb, Pb, Zn	Quartz veins (+/-epithermal) near intrusions and faults.	Wit (93N/1) 20 million tonnes 7% Pb-Zn
Skarns	Cu, Au	Limey sediments in close proximity to alkaline intrusions; skarn alteration.	Dem Lake (93K/16)

#### DEFINITION OF ANOMALOUS THRESHOLDS (RGS)

Regional Geochemical Survey (RGS) data for mapsheet 93N includes Zn, Cu, Pb, Ag, As, and Sb. This data set was complimented by additional samples from mapsheet 93N/2 that were analysed for Au. Anomalous threshold values were established statistically by using the PROBPLOT program (Stanley, 1978). Anomalous thresholds are either a) the lower (2%) threshold for the first identifiably distinct population at or above the 95th percentile or b) the average of this threshold with the upper threshold for the next lower population; whichever option is most geologically reasonable.

#### REFERENCES

REFERENCES

McLaren, G. (1990): A Mineral Resource Assessment of the Chilko Lake Planning Area, British Columbia Ministry of Energy, Mines and Petroleum Resources, Bulletin 81.

Nelson, J., Bellefontaine, K., Rees, C. and MacLean, M. (1992a): Regional Geological Mapping in the Nation Lakes Area (93N/2E, 93N/7E), British Columbia Ministry of Energy, Mines and Petroleum Resources, Geological Fieldwork, 1991, Paper 1992-1, pages 103-118.

Nelson, J., Bellefontaine, K., MacLean, M. and Rees, C., (1992b): Geology and Mineral Potential of the Chuchi Lake East Half (93N/2) and Klawii Lake East Half (93N/7) Map Areas, British Columbia Ministry of Energy, Mines and Petroleum Resources, Open File 1992-4.

55°00 Stanley, C. R. (1987): PROBPLOT: An Interactive Program to Fit Mixtures to Normal (and Log Normal) Distributions with Maximum Likelihood Optimitzation Procedures; Association of Exploration Geochemists, Special Volume 14.