

## COPPER-GOLD-SILVER DEPOSITS TRANSITIONAL BETWEEN SUBVOLCANIC PORPHYRY AND EPITHERMAL ENVIRONMENTS

By Andre Panteleyev

**KEYWORDS:** Economic geology, porphyry copper, epithermal, copper, gold, silver, magmatic, hydrothermal, acid-sulphate, high sulphidation, advanced argillic, steam heated, alunite, kaolinite, pyrophyllite, enargite.

### INTRODUCTION

The many and varied types of intrusion-related mineral deposits in circum-Pacific volcanic arcs, including gold-rich porphyry and related epithermal types, have been discussed by Berger and Henley (1989), Sillitoe (1989, 1990, 1991a, b and in press), Hedenquist *et al.* (1990) and Sillitoe and Camus (1991). The similarity in geological environments of the many described porphyry copper, copper-gold, copper-molybdenum and epithermal precious metal deposits with those in the Canadian Cordillera is evident, but the scarcity of documented acid-sulphate high-sulphidation, advanced argillic-type epithermal deposits and mineralization in British Columbia is surprising. This lack of deposits is probably only apparent and not a geologic reality. It appears to be largely due to a lack of recognition and study of acid-sulphate-type deposits and their environments except for rare cases, for example, Clapp (1915), Bradford (1985) and Diakow *et al.* (1991).

A new project has been initiated to study the inter-relationships of subvolcanic porphyry copper deposits and genetically related epithermal mineralization in British Columbia. Of particular interest are deposits with hydrothermal alteration of the kaolinite-alunite-quartz-pyrite-bearing **acid-sulphate** type (Hayba *et al.* 1985, Heald *et al.* 1987), also known as **high sulphidation** (Hedenquist, 1987) or a special case of **advanced argillic** (Meyer and Hemley, 1967). Similar mineralization and alteration suites have been described as: alunite-kaolinite (pyrophyllite), enargite-gold, enargite massive sulphide, high sulphur, Nansatsu-type, epithermal quartz-alunite, alunitic quartzite (Russian terminology), volcanic-hosted copper-arsenic-antimony, Roseki clay or acidic zone (Japanese terminology) and hot spring gold-silver.

### BACKGROUND

The spatial proximity of magmatic hydrothermal and some epithermal mineralization has been postulated in a number of geologic models (Sillitoe, 1983, 1988, 1989, 1991a; Mutschler *et al.*, 1985; Bonham, 1986, 1988; Panteleyev, 1986). This re-emphasizes Lindgren's concepts of a continuum in ore-forming hydrothermal environments in volcanic settings, from hydrothermal systems dominated by magmatic fluids at depth, to largely geothermal meteoric-groundwater systems near the surface. As stated by Henley (1991):

- "magmatic vapour from crystallizing plutons is critical to [mineralization in] the epithermal environment much as described for porphyry copper-molybdenum deposits."
- "in volcanic terranes the distinction of epithermal from porphyry-type environments of mineralization becomes largely one of convenience for exploration than one of reality." and "... a practical understanding of the relationship between magmatism and structural evolution is critical to the future of [epithermal] exploration ...".

Some epithermal deposits are positioned above or marginal to subvolcanic intrusion-related porphyry-type mineralization. Although the relationship between porphyry copper deposits and adularia-sericite-type epithermal deposits is considered by some to be speculative, the genetic connection with acid-sulphate-type deposits is well established (Henley and Ellis, 1983; Henley, 1991; Sillitoe, 1991a, b). According to Sillitoe (1989, 1991a), the latter typically occur above porphyry copper mineralization, albeit they are laterally offset in some districts by structural channeling.

Acid-sulphate alteration with associated copper-gold-silver mineralization is characterized by zoned hydrothermal mineral assemblages containing abundant silica as quartz, chalcedony and opaline silica (cristobalite), kaolin (including kaolinite, dickite and halloysite), pyrophyllite, and alunite/natroalunite. Locally, white mica (sericite/illite), mixed-layer clays, andalusite, and rarely diaspore, corundum and dumortierite are present, commonly in zonal arrangement. In some deposits late-stage barite, gypsum, anhydrite, jarosite and native sulphur are common as well as minor boehmite, phillipsite, tourmaline and accessory topaz, rutile and zunyite. The principal ore minerals, in addition to abundant pyrite and/or hematite, are gold, electrum, chalcocopyrite, copper sulphosalts (enargite - famatinite/luzonite), tetrahedrite/tennantite, bornite, chalcocite and covellite. Elevated values of gold, silver, arsenic and antimony are common in copper ores.

The advanced argillic alteration with its siliceous and aluminous mineral assemblages, commonly with the sulphate minerals alunite, gypsum and jarosite and rarely native sulphur, is a product of strongly oxidized, sulphur-rich, acidic hydrothermal fluids. It generally occurs at late stages of mineralization during declining hydrothermal activity. The acid-sulphate, advanced argillic alteration originates in three ways (Hayba *et al.*, 1985; Reye *et al.* 1989, Sillitoe, in press): from magmatic-hydrothermal fluids in which magmatic volatiles are evolved at depth from crystallizing magma and interact with surrounding groundwater as described by Henley and McNabb (1978) - the porphyry environment; near surface where steam-heated fluids occur above the water table due to vapour separation (boiling) caused by depressurization of a ascending hydrothermal flu-

ids – the epithermal environment; and by oxidation of sulphides in the supergene environment.

## INTRUSION-RELATED EPITHERMAL PRECIOUS METAL DEPOSITS IN BRITISH COLUMBIA

This new project proposes to examine, describe and study prospective environments for these deposits in British Columbia. MINFILE has five listings for enargite-bearing deposits; three of these in the Taku-Sutlahine River area are related zones within a large hydrothermal alteration system

TABLE 2-1-1  
BRITISH COLUMBIA DEPOSITS WITH ACID-SULPHATE, ADVANCED ARGILLIC ALTERATION CONTAINING KAOLINITE±PYROPHYLLITE±ALUNITE AND/OR ENARGITE\*

Location/Property Name	MINFILE	Published References
<b>Sutlahine River area</b>		
1. *THORN, DAISY, INK, Camp Creek	104K 031, 116	
2. *KAY, LIN, LIN 1-8	104K 030	
<b>Iskut River Area</b>		
3. *Johnny Mountain/REG	104B 107	
4. Treaty Glacier	104B 078	Alldrick and Britton, 1991; J.F.H. Thompson, personal communication, 1991
<b>Toodoggone River area</b>		
5. AL, Alberts Hump, Bonanza	94E 078, 79, 85, 91, 99	Diakow <i>et al.</i> , 1991
6. Brenda (Jan alunite)	94E 107	
7. SHAS (Shasta)	94E 050	
8. Silver Pond	94E 069	
<b>Central B.C.</b>		
9. Equity Silver mine	93L 001	Cyr <i>et al.</i> , 1984; Wojdak and Sinclair, 1984; 'deep acid-sulphate' – Sillitoe, 1991a
<b>Taseko River/Mt. McClure area</b>		
10. Empress	92O 033	Bradford, 1985 Company reports, Westpine Metals Ltd.
11. *Taylor-Windfall	92O 028	
<b>Vancouver Island</b>		
12. Expo (Hushamu, Mt. MacIntosh, HEP, Pemberton Hills)	92L 185, 240, 78, 308	Company reports, Moraga Resources Ltd., BHP-Utah Mines Ltd.; P.G. Dasler, personal communication, 1991
13. Red Dog	92L 200	Company reports, Moraga Resources Ltd., Crew Natural Resources Ltd.
14. Wanokana	92L 272	Company reports, Acheron Resources Ltd., P.G. Dasler, personal communication, 1991
15. Island Copper mine	92L 138, 158, 273	Cargill <i>et al.</i> , 1976; Perello, 1987; Company reports
16. Kyuquot Sound (Easy Inlet)	92L 072, 117	Clapp, 1915; MacLean, 1988
<b>Southern B.C.</b>		
17. Riverside	92INW087	MacLean, 1988
18. Pyro	92HSE131	MacLean, 1988

Note: \* denotes occurrences containing enargite. Deposits with MINFILE numbers are summarized in MINFILE from various published and unpublished sources — mainly company reports and assessment reports filed with the Ministry.

at the Thorn and adjoining properties. Pyrophyllite has been documented at four locations (MacLean, 1988). These and other deposits mentioned in various reports, mainly company reports, are listed in Table 2-1-1.

During the summer of 1991, the Thorn property and a number of northern Vancouver Island occurrences considered to be the highest priorities were briefly examined. The presence of widespread enargite in quartz veins is confirmed at the Thorn property within a large area of pyritic, intensely leached, limonitic (jarositic), sericite alteration. On northern Vancouver Island silica-clay 'caps' and pyrophyllite-bearing breccias form extensive alteration zones in Bonanza Group volcanic rocks. More detailed work in these and other areas is planned for 1992.

## SUMMARY

Zones of acid-sulphate high-sulphidation, advanced argillic alteration can host or overlie major precious metal and copper deposits. The discovery of even one major bonanza deposit in this environment can be a major economic bonanza. For example, at El Indio, Chile in 1975, the recognized economic potential of veins and breccias led to the discovery of ore reserves containing 140 tonnes of gold, 771 tonnes of silver and 0.4 million tonnes of copper in a 150 by 500-metre area (Jannas *et al.*, 1990). In addition, acid-sulphate advanced argillic alteration zones are near-surface features that can mark buried bulk-mineable porphyry copper-gold deposits such as the Island Copper orebody (Cargill *et al.*, 1976; Perello, 1987) or enargite-type copper-gold-silver deposits such as the Lepanto deposit in the Philippines and the related, newly discovered, Lepanto Far Southeast deposit with combined metal content of 526 tonnes gold and 3.45 million tonnes copper (Sillitoe, 1991a).

The target areas for this study are both areas of past exploration or newly discovered areas of interest in which hydrothermal clay-silica zones of acid-sulphate advanced argillic alteration have not been recognized or evaluated. The relationships of intrusive rocks, hydrothermally altered zones, structural controls for ore and mineralization might best be studied at a regional or district scale. As stated by Sillitoe 1991, page 202, in a summary of intrusion-related gold deposits: "An appreciation of the variety of gold deposit types in intrusion-centered systems, the geological parameters that controlled them, their mutual interrelationships and the resultant metal zoning patterns provide a cogent framework for gold exploration in volcano-plutonic arcs." With regard to recent discoveries in long-active mining districts, he further states: "No less than 25 of the 33 newly discovered [post-1979] deposits . . . are located in old mining districts. Furthermore, given that districts possess radii as great as 8 km, individual exploration targets can be large."

An invitation is extended to readers with personal knowledge of acid-sulphate, high-sulphidation, advanced argillic deposits in British Columbia to share information with the writer in order to inventory the deposits and identify the favourable geological environments.

## ACKNOWLEDGMENTS

Keith Mounjoy provided affable, competent assistance in organizing this project and during the course of fieldwork. Gerald Carlson and J.R. Woodcock, consulting geologists, Vancouver, provided the writer with numerous reports and shared personal observations about the Thorn property. Information about various northern Vancouver Island properties and advice about site visits were offered by Peter G. Dasler and David Pawliuk of Daiwan Engineering Limited on behalf of Moraga Resources Limited and Arne O. Birke-land on behalf of Keewatin Engineering Incorporated. Access to EXPO property diamond-drill core and generous discussions at the mine-site were offered by John Fleming, Island Copper mine. Constructive comments about the content and style of this report were received from Dave Lefebure and John Newell.

## SELECTED BIBLIOGRAPHY

- Alldrick, D.J. and Britton, J.M. (1991): Sulphurets Area Geology (Parts of 104A/5W, 12W; 104B/8E, 9E); *B.C. Ministry of Energy, Mines and Petroleum Resources, Open File 1991-21*.
- Berger, B.R. and Henley, R.W. (1989): Advances in the Understanding of Epithermal Gold-Silver Deposits, with Special Reference to the Western United States; in *The Geology of Gold Deposits, the Perspective in 1988, Economic Geology*, Monograph 6, pages 405-423.
- Bonham, H.F., Jr. (1986): Models for Volcanic-hosted Epithermal Precious Metal Deposits; A Review; in *Proceedings Symposium 5: Volcanism, Hydrothermal Systems and Related Mineralization, International Volcanological Congress*, Auckland, pages 13-17.
- Bonham, H.F., Jr. (1988): Models for Volcanic-hosted Precious Metal Deposits; A Review; in *Bulk Mineable Precious Metal Deposits of the Western United States*, Schafer, R.W., Cooper, J.J. and Vikre P.G., Editors, *Geological Society of Nevada*, Reno, pages 259-271.
- Bradford, J.A. (1985): Geology and Alteration in the Taseko River Area, Southwestern British Columbia; unpublished B.Sc. thesis, *The University of British Columbia*, 122 pages.
- Cargill, D.G., Lamb, J., Young, M.J. and Rugg, E.S. (1976): Island Copper; in *Porphry Deposits of the Canadian Cordillera*, Sutherland Brown, A., Editor, *Canadian Institute of Mining and Metallurgy*, Special Volume 15, pages 206-218.
- Clapp, C.H. (1915): Alunite and Pyrophyllite in Triassic and Jurassic Volcanics at Kyuquot Sound, British Columbia; *Economic Geology*, Volume 10, Number 1, pages 70-88.
- Cyr, J.B., Pease, R.B. and Schroeter, T.G. (1984): Geology and Mineralization at Equity Silver Mine; *Economic Geology*, Volume 79, pages 947-968.
- Diakow, L.J., Panteleyev, A. and Schroeter, T.G. (1991): Jurassic Epithermal Deposits in the Toodoggone River Area, Northern British Columbia - Examples of Well-preserved Volcanic-hosted, Precious Metal Mineralization; *Economic Geology*, Volume 86, pages 529-554.
- Hayba, D.O., Bethke, P.M., Heald, P. and Foley, N.K. (1985): Geologic, Mineralogic, and Geochemical Characteristics of Volcanic-hosted Epithermal Precious-metal Deposits; in *Geology and Geochemistry of Epithermal Systems*, Berger, B.R. and Bethke, P.M., Editors, *Reviews in Economic Geology*, Volume 2, pages 129-167.
- Heald, P., Foley, N.K. and Hayba, D.O. (1987): Comparative Anatomy of Volcanic-hosted Epithermal Deposits: Acid Sulfate and Adularia-Sericite Types; *Economic Geology*, Volume 82, pages 1-26.
- Hedenquist, J. W. (1987): Mineralization Associated With Volcanic-related Hydrothermal Systems in the Circum-Pacific Basin; in *Transactions of the Fourth Circum-Pacific Energy and Mineral Resources Conference*, Singapore, Horn, M.K., Editor, *American Association Petroleum Geologists*, pages 513-524.
- Hedenquist, J.W., White, N.C. and Siddeley, G., Editors (1990): *Epithermal Gold Mineralization of the Circum-Pacific; Geology, Geochemistry, Origin and Exploration*, Volumes I and II; *Journal of Geochemical Exploration*, Volume 35, 446 pages and Volume 36, 474 pages.
- Henley, R.W. (1991): Epithermal Gold Deposits in Volcanic Terranes; in *Gold Metallogeny and Exploration*, Foster, R.P., Editor, *Blackie and Sons*, pages 131-164.
- Henley, R.W. and Ellis, A.J. (1983): Geothermal Systems Ancient and Modern: A Geochemical Review; *Earth Science Reviews*, Volume 19, pages 1-50.
- Henley, R.W. and McNabb, A. (1978): Magmatic Vapor Plumes and Ground Water Interaction in Porphyry Copper Emplacement; *Economic Geology* Volume 73, pages 1-20.
- Jannas, R.R., Beane, R.E., Ahler, B.A. and Brossnahan, D.R. (1990): Gold and Copper Mineralization at the El Indio Deposit, Chile; in *Epithermal Gold Mineralization of the Circum-Pacific: Geology, Geochemistry, Origin and Exploration, II*, Hedenquist, J.W., White, N.C. and Siddeley, G., Editors, *Journal of Geochemical Exploration*, Volume 36, pages 233-267.
- Meyer, C. and Hemley, J.J. (1967): *Wall Rock Alteration; in Geochemistry of Hydrothermal Ore Deposits*, Barnes, H.L., Editor, *Holt, Rinehart and Winston, Inc.*, pages 166-235.
- MacLean, M. (1988): Talc and Pyrophyllite in British Columbia; *B.C. Ministry of Energy Mines and Petroleum Resources, Open File 1988-19*, 108 pages.
- Mutschler, F.E., Griffen, M.E., Scott Stevens D. and Shannon, S.S. Jr. (1985): Precious Metal Deposits Related to Alkaline Rocks in the North American Cordillera - An Interpretive Review; *Geological Society South Africa, Transactions*, Volume 88, pages 366-377.
- Panteleyev, A. (1986): A Canadian Cordilleran Model for Epithermal Gold-Silver Deposits, *Geoscience Canada, Volume 13*, pages 101-111; and in *Ore Deposit Models*, Roberts, R.G. and Sheahar, P.A., Editors, *Geoscience Canada, Reprint Series 3*, pages 31-44.

- Perello, J.A. (1987): The Occurrence of Gold at Island Copper Mine, Vancouver Island, British Columbia; unpublished M.Sc. thesis, *Queen's University*, Kingston, 85 pages.
- Rye, R.O., Bethke, P.M. and Wasserman, M.D. (1989): Diverse Origins of Alunite and Acid-Sulfate Alteration: Stable Isotope Systematics; *United States Geological Survey*, Open-File Report 89-5, 33 pages.
- Sillitoe, R.H. (1983): Enargite-bearing Massive Sulfide Deposits High in Porphyry Copper Systems; *Economic Geology*, Volume 78, pages 348-352.
- Sillitoe, R.H. (1988): Gold and Silver Deposits in Porphyry Systems; in *Bulk Mineable Precious Metal Deposits of the Western United States*, Schafer, R.W., Cooper, J.J. and Vikre, P. G., Editors, *Geological Society of Nevada*, Reno, pages 233-257.
- Sillitoe, R.H. (1989): Gold Deposits in Western Pacific Island Arcs: The Magmatic Connection; in *The Geology of Gold Deposits: The Perspective in 1988*, Keays, R.R., Ramsay, W.R.H. and Groves, D.I., Editors, *Economic Geology*, Monograph 6, pages 274-291.
- Sillitoe, R.H. (1990): Gold-rich Porphyry Copper Deposits of the Circum-Pacific Region – An Updated Overview; in *Pacific Rim Congress 90*, Gold Coast, Queensland, 1990, Proceedings, Volume 2, *Australian Institute of Mining and Metallurgy*, pages 119-126.
- Sillitoe, R.H. (1991a): Intrusion-related Gold Deposits; in *Gold Metallogeny and Exploration*, Foster, R.P, Editor, *Blackie and Sons*, pages 165-209.
- Sillitoe, R.H. (1991b): Gold Metallogeny in Chile; *Economic Geology*, Volume 86, pages 1187-1205.
- Sillitoe, R.H. (in press): Gold-rich Porphyry Copper Deposits: Occurrence Model and Exploration Implications; in *Proceedings 8th IAGOD Symposium, IUGS/IAGOD Deposit Modeling Program Conference*, Ottawa, 1990.
- Sillitoe, R.H. (in press): Epithermal Models, Geometric Controls and Shallow Features; in *Proceedings 8th IAGOD Symposium, IUGS/IAGOD Deposit Modeling Program Conference*, Ottawa, 1990.
- Sillitoe, R.H. and Camus, F., Editors (1991): Gold Deposits of the Chilean Andes; *Economic Geology*, Volume 86, pages 1153-1345.
- Wojdak, P.J. and Sinclair, A.J. (1984): Equity Silver-Copper-Gold Deposit: Alteration and Fluid Inclusion Studies; *Economic Geology*, Volume 79, pages 969-990.